REFERENCE ONLY



JNIVERSITY OF LONDON THI	ESIS
--------------------------	------

Degree PhD Year 2006 Name of Author L1, Cheryl Suet-Yee

COPYRIGHT

This is a thesis accepted for a Higher Degree of the University of London. It is an unpublished typescript and the copyright is held by the author. All persons consulting the thesis must read and abide by the Copyright Declaration below.

COPYRIGHT DECLARATION

I recognise that the copyright of the above-described thesis rests with the author and that no quotation from it or information derived from it may be published without the prior written consent of the author.

LOAN

Theses may not be lent to individuals, but the University Library may lend a copy to approved libraries within the United Kingdom, for consultation solely on the premises of those libraries. Application should be made to: The Theses Section, University of London Library, Senate House, Malet Street, London WC1E 7HU.

REPRODUCTION

University of London theses may not be reproduced without explicit written permission from the University of London Library. Enquiries should be addressed to the Theses Section of the Library. Regulations concerning reproduction vary according to the date of acceptance of the thesis and are listed below as guidelines.

- Before 1962. Permission granted only upon the prior written consent of the A. author. (The University Library will provide addresses where possible).
- 1962 1974. In many cases the author has agreed to permit copying upon B. completion of a Copyright Declaration.
- C. Most theses may be copied upon completion of a Copyright 1975 - 1988. Declaration.
- 1989 onwards. Most theses may be copied. D.

This thesis comes within category D.



This copy has been deposited in the Library of

LCI

This copy has been deposited in the University of London Library, Senate House, Malet Street, London WC1E 7HU.

Viral and Host Gene Expression Patterns in Human Herpesvirus Type-7 Infection of T-Cells

Cheryl Suet-Yee Li

Submitted to the University of London for the degree of Doctor of Philosophy

December 2005

Viral Genomics and Bioinformatics Group Department of Immunology and Molecular Pathology Windeyer Institute of Medical Sciences University College London UMI Number: U593060

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



UMI U593060 Published by ProQuest LLC 2013. Copyright in the Dissertation held by the Author. Microform Edition © ProQuest LLC. All rights reserved. This work is protected against unauthorized copying under Title 17, United States Code.



ProQuest LLC 789 East Eisenhower Parkway P.O. Box 1346 Ann Arbor, MI 48106-1346

ABSTRACT

Human herpesvirus type 7 (HHV-7), with its high prevalence and apparently nonpathogenic nature, is believed to be a virus very well adapted to humans. It is therefore useful to study the transcription patterns of the virus and the host to understand the life cycle and pathogenesis (or the lack of it) of HHV-7 infection. In order to obtain the global pattern of host and viral gene expression, we have made a DNA microarray containing probes that target all 86 predicted HHV-7 open reading frames and approximately 1000 cellular genes. This is the first description of an integrated hostpathogen microarray for HHV-7.

The microarrays were used to follow the gene expression patterns over a 72-hour time course infection of a CD4⁺ T-cell line (Sup-T1) by HHV-7 *in vitro*, which revealed an overall increase of HHV-7 transcripts over time, as well as the coordinated transcription profiles for groups of viral genes with similar functions. HHV-7 genes predicted to be immediate-early transcription factors were the first genes to be expressed, followed by genes involved in viral DNA replication, and at later stages, the genes encoding structural proteins and proteins of packaging functions were expressed. Temporal classification was assigned for 68 HHV-7 ORFs, of which a selection was confirmed by RT-PCR in the presence of metabolic inhibitors of *de novo* protein synthesis and viral DNA replication. Both gene expression profiling and inhibitor treatment demonstrated that the lytic gene expression of HHV-7 was temporally regulated, consistent with the classification of viral gene expression into immediate-early, early and late transcripts in other herpesviruses.

ACKNOWLEDGEMENTS

This is probably my favourite page in this thesis, when I finally have the chance to breathe and to reflect on the past few years. Throughout this PhD, there have been difficult and wonderful times, sometimes simultaneously, and it has been a remarkable experience indeed. I would like to thank many people who have provided help towards the production of this thesis, and who have shown me kindness thoughout the process.

First of all, I am truly grateful for having Paul Kellam as my supervisor. Thank you Paul, for your guidance, patience, encouragement, and for sharing your vision and expertise in genomic science with all of us in VGB. I thank Mary Collins, Robin Weiss, Aíne McKnight, Ari Fassati and Vince Emery for their invaluable suggestions and inspirations. The HHV-7 virus work could not have happened without the help of Duncan Clark from the Royal Free Virology Department, and the production of the microarrays would be impossible without the contributions from Ed Tsao and Jane Rasaijaah. I also thank Nicola Gilbert, Liz Thomson, Carolyn McElvaney and John Cheshire for their generous help all these years. This PhD would be challenging but no fun without all my colleagues in the VGB group, who provided a vibrant environment for research. I am thankful to the MRC for funding this PhD project.

Finally, thanks to my parents for their understanding and patience to a daughter who never calls, visits only once a year, and for their support in my choice of career. Last but not least, thanks to Jacob, for sharing everything with me.

Thanks be to God. May His name be praised.

CONTENTS

Abstract	2
Acknowledgements	
Contents	
Figures	
Tables	
Abbreviations	13
CHAPTER 1 Introduction	
1.1 Human Herpesvirus 7	
1.1.1 A brief history of human herpesvirus type-7	
1.1.2 Human herpesvirus	
1.1.3 Morphology and culture in vitro	
1.1.4 Genome	
1.1.5 Gene expression	
1.1.6 Cell entry and Receptor	
1.1.7 Cell tropism & Persistence	
1.1.8 Transmission & Prevalence	
1.1.9 Disease association & Immune response	
1.1.10 Cellular interaction	
1.1.11 Life cycle	
1.1.11.1 Latency/Persistence	
1.1.11.2 Lytic infection	
1.2 DNA Microarrays	
1.2.1 The rise of the genome, transcriptome, and proteome	
1.2.2 Principles of DNA microarray technology	
1.2.2.1 Microarray platforms	
1.2.2.2 Microarray and gene expression analysis	45
1.2.3 Analysis of microarray data	
1.2.3.1 Data filtering and transformation	
1.2.3.2 Data normalisation	
1.2.3.3 Identification of differentially expressed genes	

1.2.3.4 Expression patterns recognition by clustering analysis	51
1.2.3.5 Confirmation of microarray results	53
1.2.4 Applications of microarrays in host-pathogen interactions	55
1.2.4.1 Microarray studies of microbial gene expression	55
1.2.4.2 Microarray studies of viral gene expression	56
1.2.4.3 Microarray studies of host gene expression during viral infection	57
1.3 Aim and Scope of thesis	61

CHAPTER 2 Materials & Methods	62
2.1 Buffers and Solutions	63
2.2 Mammalian cell culture	63
2.2.1 Cell lines	63
2.2.2 Cell growth media	63
2.2.3 Thawing	64
2.2.4 Passaging cells	64
2.2.5 Freezing cells	64
2.3 Culturing HHV-7	65
2.3.1 Virus strain	65
2.3.2 Virus infection	65
2.3.3 Virus culture and maintenance	65
2.3.4 Stock preparation	65
2.3.5 Virus quantification	66
2.3.5.1 Observation of cytopathic effect (CPE)	66
2.3.5.2 Immunofluorescence assay (IFA)	66
2.3.5.3 Infectious units count (IUC)	67
2.3.5.4 TCID ₅₀ determination	67
2.3.6 Flow cytometry	68
2.3.7 Time course infection of SupT1 cells by HHV-7	68
2.3.8 Metabolic inhibitor assay for temporal gene expression mapping	69
2.4 Producing HHV-7 array probes	70
2.4.1 Array probe primers	70
2.4.2 HHV-7 DNA preparation	70
2.4.3 Initial amplification of viral DNA	70
2.4.4 Agarose electrophoresis of PCR product	71

2.4.5 Cloning array probes	. 71
2.4.5.1 Competent cell preparation	. 71
2.4.5.2 Ligation of DNA into pGEM-T easy vector	. 72
2.4.5.3 Transformation	. 72
2.4.6 Identification of array probes	. 72
2.4.6.1 Growth of bacterial clones	. 72
2.4.6.2 Screening for viral insert-containing clones by PCR	. 73
2.4.6.3 Purifying plasmid DNA from positive clones	. 73
2.4.6.4 Verifying sequence of viral insert	. 74
2.5 Production of human gene probes	. 74
2.5.1 PCR optimisation	. 74
2.5.2 Preparing lymphochip clones for human array probes	.75
2.6 Production of custom-made microarrays	. 76
2.6.1 Array PCR to amplify viral and human probes	. 76
2.6.2 Human and viral gene probes purification	. 76
2.6.3 Microarray spotting and processing	.77
2.7 Measuring transcript abundance using custom-made microarrays	.77
2.7.1 RNA extraction from TRIzol	. 77
2.7.1.1 Total RNA extraction	. 77
2.7.1.2 DNase treatment of total RNA	. 78
2.7.1.3 Messenger RNA purification	. 78
2.7.2 RNA extraction from RLT lysate	. 79
2.7.2.1 Total RNA extraction	. 79
2.7.2.2 Sense-strand RNA amplification	. 80
2.7.3 Reference mix	. 81
2.7.4 Fluorescent labelling of RNA	. 81
2.7.5 Hybridisation onto microarray	. 82
2.8 Array scanning and data collection	. 83
2.9 Array data processing and normalisation	. 83
2.10 Clustering analysis	. 84
2.11 Significance Analysis of Microarrays (SAM)	. 84
2.12 Reverse transcription-PCR (RT-PCR)	. 85
2.12.1 RT-PCR reaction	. 85
2.12.2 Gel purification	. 85

2.12.3 Sequence analysis of gel bands
CHAPTER 3 Production of HHV-7 Microarrays
3.1 Introduction
3.1.1 Reasons for HHV-7 microarray production
3.2 Results
3.2.1 Production of HHV-7 array probes
3.2.1.1 Primer design and initial PCR amplification from genomic DNA90
3.2.1.2 Cloning of array probes96
3.2.1.3 Sequence verification of clones
3.2.1.4 Amplification and purification of HHV-7 array probes 101
3.2.2 Optimisation of array probe amplification103
3.2.2.1 Optimisation of PCR reaction103
3.2.2.2 Robotic liquid handling104
3.2.3 Production of HHV-7-human microarrays107
3.2.3.1 Production of human gene probes
3.2.3.2 Microarray printing110
3.2.4 Characteristics of HHV-7-human microarrays 112
3.2.4.1 Effects of Dye-bias and RNA preparation
3.2.4.2 Data filtering and Signal-to-Noise Ratio threshold
3.2.4.3 Sensitivity of HHV-7 probes119
3.2.4.4 Specificity of microarrays
3.2.4.5 Reproducibility of microarrays
3.2.4.6 Use of HHV-7-human microarray in gene expression profiling 127
3.2.5 Data Normalisation 128
3.2.5.1 Effects of Total Intensity Normalisation
3.2.5.2 Effects of Lowess Normalisation
3.3 Summary
CHAPTER 4 Infectivity of Human Herpesvirus type-7 (HHV-7) in SupT1 cells 133
4.1 Introduction
4.1.1 Infectivity of HHV-7 134

4.1.4 Scope of chapter 4	137
4.2 Results	138
4.2.1 Maintaining HHV-7 infection in SupT1 cells	138
4.2.2 Morphology of infected cells	139
4.2.3 Quantification of HHV-7 virus	142
4.2.3.1 Quantification by Tissue Culture Infectious Dose ₅₀ (TCID ₅₀)	142
4.2.3.2 Quantification by Infectious units count (IUC)	143
4.2.3.3 Summary of Quantification	149
4.2.4 Optimisation of HHV-7 culturing to achieve a high multiplicity of	f infection
	150
4.2.4.1 Production of high tire virus	150
4.2.4.2 Effect of Spinoculation & Polybrene on Inoculation efficiency.	152
4.2.5 RNA amplification	157
4.2.5.1 Reasons for RNA amplification	157
4.2.5.2 Sense-strand amplification	157
4.3 Summary	163
CHAPTER 5 Gene Expression Profile of HHV-7	
5.1 Introduction	165
5.1.1 Classification of herpesvirus gene expression	165
5.1.2 Temporal regulation of HHV-7 gene expression	
5.2 Results	
5.2.1 HHV-7 infection of SupT1 cells	169
5.2.2 HHV-7 gene transcription increased during infection	172
5.2.3 HHV-7 expression profile during lytic infection of SupT1 cells	
5.2.3.1 Clustering (100%-presence) of HHV-7 genes	178
5.2.3.2 Clustering (80%-presence) of HHV-7 genes	181
5.2.3.3 HHV-7 genes of similar functions were co-expressed	188
5.2.3.4 Immediate-early transactivators	190
5.2.3.5 DNA replication and metabolism	191
5.2.3.6 Capsid assembly and Virion structure	193
5.2.3.8 Classification of HHV-7 genes from clustering pattern	197
5.2.3.9 Predicting functions for previously uncharacterised genes	198

5.2.4 Temporal classification of HHV-7 genes using metabolic inhibitors	200
5.2.4.1 Immediate-early and Early genes	200
5.2.4.2 Early and Late genes	201
5.2.4.3 Comparison of expression profiling vs metabolic inhibitor assay	206
5.2.5 Temporal classification of HHV-7 genes by RT-PCR	211
5.2.5.1 RT-PCR conditions	211
5.2.5.2 Classification of HHV-7 gene expression by RT-PCR	212
5.2.6 Host gene expression	217
5.3 Summary and Discussion	225

CHAPTER 6 Conclusions and Future Directions	
6.1 HHV-7 gene expression	
6.2 HHV-7 culturing	231
6.3 Host genes expression by HHV-7	

Appendix A Sequence alignments of HHV-7 array probes	233
Appendix B Identities of Lymphochip clones on custom-made microarrays	257
Appendix C Sequence alignments of splice variants of HHV-7	271
References	273

FIGURES

Figure 1.1 HHV-7 genome architecture
Figure 1.2 Herpesvirus conserved gene blocks
Figure 1.3 Temporal gene expression in herpesviruses
Figure 1.4 Principle of microarrays
Figure 3.1 Diagram of HHV-7 microarray production93
Figure 3.2 HHV-7 ORFs and positions of array probes
Figure 3.3 PCR amplification of HHV-7 fragments
Figure 3.4 Screening array clones for HHV-7 inserts
Figure 3.5 Re-amplified and purified HHV-7 array probes
Figure 3.6 PCR optimisation and DNA polymerase enzymes batch test 105
Figure 3.7 PCR Amplification and purification of human array probes 108
Figure 3.8 Format of HHV-7 host-pathogen microarray111
Figure 3.9 Effects of Dye-bias on array signal distribution
Figure 3.10 Data filtering and SNR threshold determination
Figure 3.11 Sensitivity of HHV-7 probes
Figure 3.12 Specificity of HHV-7 probes
Figure 3.13 Reproducibility of HHV-7 probes
Figure 3.14 Tree structure of clustering analysis of selected microarrays
Figure 3.15 Effects of normalisation example I (array S14)
Figure 3.16 Effects of normalisation example II (array S20)131
Figure 4.1 Morphology of HHV-7 infected cells
Figure 4.2 IFA and FACS of HHV-7 infected cells
Figure 4.3 Quantification of HHV-7 by IFA
Figure 4.4 Quantification of HHV-7 by FACS 146
Figure 4.5 Configuration of FACS 147
Figure 4.6 Summary of quantification methods 148
Figure 4.7 Effect of Spinoculation & Polybrene on HHV-7 inoculation efficiency 156
Figure 4.8 Diagram of the sense strand RNA amplification procedure
Figure 4.9 Bioanalyer analysis of total and amplified RNA

Figure 5.1 IFA confirmation of viral infection	170
Figure 5.2 Percentage of infected cells increased with time	171
Figure 5.3 Abundance of HHV-7 transcripts increased with time	174
Figure 5.4 Cluster analysis of time course (100%-presence)	180
Figure 5.5 Cluster analysis of time course (80%-presence)	183
Figure 5.6 Mean expression ratios by cluster	187
Figure 5.7 Mean expression ratios by function	189
Figure 5.8 Mean expression ratios of selected genes	195
Figure 5.9 Sequence alignment of HHV-7 U55A and U55B	196
Figure 5.10 Analysis of metabolic inhibitor assay by clustering	202
Figure 5.11 Analysis of metabolic inhibitor assay by SAM	204
Figure 5.12 Mean expression ratios of IE/E and L genes	205
Figure 5.13 RT-PCR trial and optimisation	214
Figure 5.14 RT-PCR amplification of selected HHV-7 transcripts	215
Figure 5.15 Clustering analysis of host gene expression	222
Figure 5.16 SAM analysis of host gene expression	223
Figure 5.17 Signal and pass rate of HHV-7 and cellular probes	224
Figure 5.18 HHV-7 genome map with temporal classification	227

TABLES

Table 1.1 Human herpesviruses - classification and disease associations
Table 1.2 Strains of HHV-7 21
Table 1.3 Predicted functions of HHV-7 ORFs and betaherpesvirus homologues 26
Table 1.4 Microarray studies of host gene expression in viral infections 59
Table 2.1 Constituents of buffers and solutions 63
Table 2.2 Cell lines 63
Table 2.3 Conditions for DNA polymerases in batch test 75
Table 3.1 HHV-7 array primers
Table 3.2 Cloning and sequence verification of HHV-7 array probes 99
Table 3.3 DNA polymerase batch test 104
Table 3.4 Test hybridisations for the host-pathogen microarrays 112
Table 3.5 Comparison of SNR for a subset of HHV-7 probes 122
Table 3.6 Reproducibility of microarrays – Pearson correlation coefficients
Table 4.1 Results using different quantification methods of HHV-7 stocks
Table 4.2 Production of high titre virus stocks 151
Table 4.3 Effects of spinoculation and polybrene on HHV-7 stock 020723153
Table 4.4 Effects of spinoculation and polybrene on various HHV-7 virus stocks 155
Table 4.5 Example arrays hybridised to messenger RNA or amplified RNA159
Table 5.1 Selected publications on temporal mapping of herpesvirus gene expression
Table 5.2 Number of HHV-7 transcripts detected increased with time
Table 5.3 Classification of HHV-7 genes in this thesis and previous studies
Table 5.4 Temporal classification by RT-PCR results compared with microarray results
and previous publication
Table 5.5 Significantly diffrentiated cellular genes in HHV-7 infected SupT1 cells 219

ABBREVIATIONS

°C	Degree Celsius
2d-ael	2-dimentional gel
Acc. No	Accession number
	Analysis of variance
aRNA	Amplified RNA
B-coll	Bone marrow-derived lymphocyte
	Bacillus of Calmette and Guérin
	Bovine bernesvirus
	Basic local alignment search tool
	BLAST puckeotide
DLAJIN	BLAST HUCIEOlide
DSA C terminue	Corboxyd (COOH) torminuo
Clerminus	Carboxyi (COOH)-terminus
	Cord blood mononuclear cell
	Complementer (DNA
CUNA	Complementary DNA
	Chronic faugue syndrome
CGH	Comparative genome hybridisation
CHX	Cycloneximide
Cm ²	Squared centimetre
CMV	Cytomegalovirus
CO ₂	Carbon dioxide
CPE	Cytopathic effect
dATP	Deoxyadenosine triphosphate
dCTP	Deoxycytidine triphosphate
ddATP	Dideoxyadenosine triphosphate
ddCTP	Dideoxycytidine triphosphate
ddGTP	Dideoxyguanosine triphosphate
ddUTP	Dideoxyuridine triphosphate
dGTP	Deoxyguanosine triphosphate
dITP	Deoxyinosine triphosphate
DC	HHV-7 strain DC
DMEM	Dulbeccos modified Eagle's media
DMSO	Dimethyl sulphoxide
DNA	Deoxyribonucleic acid
DNase	Deoxyribonuclease
dNTP	Deoxynucleotide triphosphate
dpi	Day post infection
DR	Direct repeat
DTT	Dithiothreitol
dTTP	Deoxythymidine triphosphate
dUTPase	Deoxyuridine triphosphatase
E	Early
E. coli	Escherichia coli
EBNA	Epstein-Barr nuclear antigen
EBV	Epstein-Barr virus
EDTA	Ethylenediaminetetraacetic acid
EHV	Fauine hernesvirus
FACS	Eluorescence activated cell sorting
FBS	Foetal bovine serum
FCS	Foetal calf serum
FDR	False discovery rate
FITC	Fluorescein isothioovanata
ESC	Forward souther
n	Grom
ษ aR	Glycoprotoin P
эс aH	Giveenatein H
9'' 0	Giveoprotein I
gu al	Giycoprotein J Chycoprotein L
yr	Giycoprotein L

gM	Glycoprotein M
Ğ-protein	Guanine nucleotide binding proteins
h	Hour
HCL	Hierarchical clustering
HCMV	Human cytomegalovirus
HHV	Human herpesvirus
HIV	Human immunodeficiency virus
hni	Hour post infection
HD\/	Human papilloma virus
	Hernes simplex virus
	Infected cell polypeptides
IE	Immediate early
	Immunofluorescence assav
IDTC	Isopropylthio-D-galactoside
	Infectious units count
	In vitro transcription
JI	
	Kapacila acrooma associated horpesvirus
NORV I	Lata
LD	
	Log Dase 2
LOWESS	Local weighted scatterplot smoothing
M. IUDERCUIOSIS	
M/A	Scatter plot of Cy5/Cy3 ratio versus geometric mean
mA	
MCMV	Murine cytomegalovirus
mCP	Minor capsid protein
MCP	Major capsid protein
MeV	Multiexperiment viewer
μg	Microgram
MHV	Murine gammaherpesvirus
MIAME	Minimal information about a microarray experiment
MIDAS	Microarray Data Analysis System
MIE	Major immediate-early
min	Minute
μΙ	Microlitre
ml	Millilitre
μM	Micromolar
mM	Millimolar
umol	Micromole
MOI	Multiplicity of infection
mRNA	Messenger RNA
na	Nanogram
nm	Nanometre
N-terminal	Amino (NH ₂)-terminal
ORP	Origin-hinding protein
	Ontical density
ORE	Open reading frame
orilyt	Origin of lutic replication
D/S	popioillin and strontomyoin
	Phoenbonocostic coid
	Priosphonoacelic aciu Desinherel bloed trimphenite
	Phoenhoto huffered antina
	Phymerece seline
	Polymerase chain reaction
	Prosphonotormic acid
	Picomole Disata multiplica tuka
	Photomultiplier tube
	Milyridsis rosea
FIFE	roiytetratiuoroetnyiene

RK	HHV-7 strain RK
RNA	Ribonucleic acid
RNase	Ribonuclease
RPMI	Roswell Park Memorial Institute
RT-PCR	Reverse transcription-PCR
S	Second
SAM	Significance analysis of microarrays
SARS-CoV	Severe acute respiratory syndrome-associated coronavirus
SDS	Sodium dodecyl sulphate
SNP	Single nucleotide polymorphism
SNR	Signal-to-noise ratio
SOM	Self-organising map
SSC	Side scatter
SSC	Saline sodium citrate
SSPE	Saline-sodium phosphate-EDTA
TAE	Tris-acetate-EDTA
T-cell	Thymus-derived lymphocyte
TCID ₅₀	Tissue culture infectious dose ₅₀
TE	Tris-EDTA
TIGR	The Institute for Genome Research
TNF	Tumour necrosis factor
TP	Transport protein
TRAIL	TNF-related apoptosis-inducing ligand
TRAIL-R	TRAIL receptor
T-RNA	Total RNA
U	Unique region
UK	United Kingdom
UL	Unique long region
US	Unique short region
UV	Ultra-violet
V	lg variable region
V	Volt
v/v	Volume by volume
v-fos	Viral homologue of the fos family of transcription factors
vIRFs	Viral interferon regulatory factors
VZV	Varicella zoster virus
w/v	Weight by volume
X-gal	5-bromo-4-chloro-3-indolylD-galactoside

CHAPTER 1

INTRODUCTION

1.1 Human Herpesvirus 7

1.1.1 A brief history of human herpesvirus type-7

Human herpesvirus type-7 (HHV-7) was discovered in 1990, when Frenkel and colleagues observed abnormal cytopathic effect (CPE) in primary T-cells, obtained from the peripheral blood of a healthy individual, RK, under T-cell activating conditions. A previously unidentified virus was isolated from the culture and typical herpesvirus virions were revealed by electron microscopy. The virion had a diameter of 170 nm, consisting an electron dense cylindrical core, capsid, tegument, and envelope (Frenkel et al., 1990; Secchiero et al., 1994). Restriction pattern and DNA hybridisation analysis showed the virus to be distinct from the six known human herpesviruses at the time (Frenkel et al., 1990), and was therefore designated the seventh human herpesvirus. HHV-7 was classified as a member of the Roseolovirus genus of the betaherpesvirus subfamily. Soon after, laboratory cultivation of HHV-7 in a continuous cell line (SupT1) was reported (Berneman et al., 1992a), enabling easier production of the virus and advancing research, including the identification of a cellular receptor (Lusso et al., 1994) and deciphering the complete genome sequence (Megaw et al., 1998; Nicholas, 1996).

1.1.2 Human herpesvirus

To date, 8 human herpesviruses have been identified. They share some common features, for example they are large viruses of 120 - 300 nm in diameter, with a complex virion structure composed of an envelope, a tegument and an icosahedral capsid that encase a relatively long (120 - 250 kb), linear, double stranded DNA genome. The herpesvirus genomes consist of 7 blocks of core genes, which are preserved in all members albeit arranged in different order (Roizman, 2001b). All herpesviruses encode a large number of enzymes involved in nucleic acid metabolism, DNA synthesis, and protein processing. The expression of these genes in productive infection results in progeny virus release and the death of host cell. All herpesviruses can establish a latent infection in the host. Viral genomes exist as closed circular molecules in the host cell

during latency, and viral gene expression is minimised (Roizman, 2001b). Nonetheless, each herpesvirus has unique properties and some viruses are more similar than others. Herpesviruses are classified into three subfamilies, alphaherpesvirus, betaherpesvirus and gammaherpesvirus, based on biological properties (Table 1.1). The closest relative of HHV-7 is HHV-6, discovered in 1986 (Salahuddin et al., 1986), with amino acid sequence similarities ranging from 41 to 75% between products of the core gene homologues (Nicholas, 1996). These two viruses belong to the genus of Roseolovirinae.

Table 1.1 Human he	pesviruses -	 classification 	and disease	associations
--------------------	--------------	------------------------------------	-------------	--------------

Common name of virus and subfamily	Human herpesvirus nomenclature	Disease association
Alphaherpesviruses		
Herpes simplex virus type 1 (HSV-1)	HHV-1	Oropharygeal herpes (cold sore)
Herpes simplex virus type 2 (HSV-2)	HHV-2	Genital herpes
Varicella zoster virus (VZV)	HHV-3	Varicella (chicken pox)
		Herpes zoster
Betaherpesviruses		
Human cytomegalovirus (HCMV)	HHV-5	Mononucleosis, Retinitis
		Complications in
		immunocompromised/transplant
Human herpesvirus 6A	HHV-6A	Exanthem subitum (rare)
Human herpesvirus 6B	HHV-6B	Exanthem subitum
		Transplant complications
Human herpesvirus 7	HHV-7	Exanthem subitum
Gammaherpesviruses		
Epstein-Barr virus (EBV)	HHV-4	B-cell lymphoma
,		Nasopharyngeal carcinoma
Kaposi's sarcoma-associated virus	HHV-8	Kaposi's sarcoma (KS)
(KSHV)		Primary effusion lymphoma
		Multicentric Castleman's disease

1.1.3 Morphology and culture in vitro

HHV-7 has been cultured in vitro in the CD4⁺ immature T-cell line SupT1 (Berneman et al., 1992a), activated CD4⁺ T-cells purified from peripheral blood lymphocytes (Secchiero et al., 1994), peripheral blood mononuclear cells (PBMCs) (Berneman et al., 1992b) and cord blood mononuclear cells (CBMCs) (Ablashi et al., 1998). HHV-7 infection has a 4-day cycle in cord blood lymphocytes (Black et al., 1997). Infection in SupT1 is best maintained in medium supplemented with 5% foetal calf serum (FCS) (Black et al., 1997), probably because a lower FCS concentration (optimum for cells 10%) facilitates virus propagation by retarding cell division. Light microscopy of infected SupT1 and primary T-cells reveal refractile, dense, vacuolated cells and giant multinucleated syncytia (Ablashi et al., 1998; Secchiero et al., 1997a), indicative of viral induced cytopathological effect (CPE). HHV-7 nucleocapsids are observed in the nucleus and cytoplasm of infected cells (Ablashi et al., 1998), and mature virions are seen in extracellular space (Secchiero et al., 1994; Secchiero et al., 1997a). Infected SupT1 also elongate and loosely attach to the culture flask. These flask-attached SupT1 cells contain HHV-7 nucleocapsids and are capable of passing on infection (Ablashi et al., 1998). During the course of in vitro infection, viral load increases with time while the viable cell count decreases (Secchiero et al., 1997a). The infected, multinucleated cells die by necrosis, while apparently uninfected, small cells in the same culture undergo apoptosis (Secchiero et al., 1998b; Secchiero et al., 1997a). Both apoptosis and necrosis of these cells were inhibited by PFA (phosphonoformic acid), a viral DNA synthesis inhibitor. Ablashi et al suggested HHV-7 is a cell-associated virus hence only released by cell lysis such as freeze-thawing (Ablashi et al., 1998), whereas Secchiero et al reported extracellular virus particles by electron microscopy (Secchiero et al., 1994). Various strains of HHV-7 hav been described in the literature, summarised in Table 1.2.

Table 1.2 Strains of HHV-7

Strain	Genome	Source	Titre (TCID ₅₀)	Particles/ml	Reference ^b
AL		PBMCs of healthy adult			1
CZ		Infant with febrile syndrome			2
H7-2		PBMC/saliva of children	10 ³	1.8×10 ⁹	3
H7-3		PBMC/saliva of children	10 ³	1.7×10 ⁹	3
H7-4		PBMC/saliva of children	10 ⁴ -10 ⁵	3.8×10 ⁹	3
JB	Same restriction pattern to JI	PBMC of adult with CFS ^a	10 ³	1.5×10 ⁹	1, 3
JHC		CBMC of adult	10 ³	3.9×10 ⁹	3
JI	144861 bp	PBMC of adult with CFS ^a		a an an far a sharin	1, 4, 5
KHR		PBMC/saliva of children (Japan)	10 ⁴ -10 ⁵	1.9×10 ⁹	3
RK	153080 bp	PBMC of healthy adult			6, 7

* CFS, chronic fatigue syndrome

^b 1, Secchiero et al., 1994; 2, Menegazzi et al., 1999; 3, Ablashi et al., 1998; 4, Berneman et al., 1992b; 5, Nicholas, 1996; 6, Frenkel et al., 1990, 7, Megaw et al., 1998

1.1.4 Genome

Two strains of HHV-7, JI (Nicholas, 1996) and RK (Megaw et al., 1998) have been fully sequenced and reveal double stranded DNA genomes of approximately 145 and 153 kb, respectively. The HHV-7 genome consists of a central unique region (U) flanked by terminal direct repeats (DR) on either end (Figure 1.1). The genome contains reiterated sequences in the DR regions (T1 and T2) and in the unique region (R1 and R2). These vary in length and copy number between strains, hence account for the difference in overall genome length. Otherwise the genomes of the two strains are highly conserved, differing only by 0.1% (179 bp). Genome analyses have identified the presence of conserved genes, arranged in 7 core gene blocks in the unique region (Figure 1.1), a feature conserved among herpesviruses (Figure 1.2). These core genes encode products involved in regulatory functions, DNA replication and repair, nucleotide metabolism and virion assembly. Amino acid sequences of the core genes have the highest identity (41-75%) to HHV-6, and they are in the same order and orientation as HCMV and HHV-6. The genome of HHV-7 is predicted to have at least 86 open reading frames (ORFs), some with predicted functions, by sequence homology to HHV-6 (Gompels et al., 1995) and HCMV (Chee et al., 1990) (Table 1.3). An extra 14 ORFs has been suggested (Nicholas, 1996) but later considered to be either introns of adjacent ORFs or unlikely to encode protein (Megaw et al., 1998). Of the 86 ORFs there are 84 different genes; 2 ORFs in the 3'-end DR region (DR1' and DR6') are identical to their counterparts in the 5'-end (DR1 and DR6). Up to 11 genes in HHV-7 may be spliced (Megaw et al., 1998). Two regions in the genome (U17-U19 and U86-U90) display conserved layout and splicing pattern to the two IE loci in HCMV (Pellett, 2001), although the amino acid sequences are diverged. 8 ORFs in the HHV-7 genome (DR6, U2, U3, U7, U8, U17, U25, U95) are members of the HCMV US22 gene family of duplicated and diverged genes. Except for 7 genes specific to the roseoloviruses (U20, U21, U23, U24, U26, U100), the rest of the HHV-7 ORFs all have counterparts in HCMV. Amongst the roseoloviruses, HHV-7 encodes U53.5 that has no homologue in HHV-6, and a duplicated gene (U55A and U55B) that is present in HHV-6 as a single gene (U55); otherwise there are no known genes unique to HHV-7 and absent in HHV-6. In contrast, HHV-6 encodes at least 5 genes (DR3, U6, U22, U83, U94) that have no homologues in HHV-7. HHV-7 therefore has a smaller genome and encodes fewer

genes than HHV-6 and HCMV. The origin of lytic replication (*oriLyt*) is identified between U41 and U42 (Figure 1.1).



Figure 1.1 HHV-7 (RK) genome arrangement. Shaded region denotes the terminal direct repeats. Reiterated sequences (T1, T2, R1, R2) are labelled and shown as black rectangles. Predicted open reading frames (prefixed by U, DR) are shown as block arrows in the direction of expression, above their gene names. The origin of lytic replication (oriLyt) is indicated by black arrow. Scale in kilobases. The two regions homologous to HCMV immediately early loci are shown as striped rectangles. The 7 herpesvirus core gene blocks and 1 block conserved in betaherpesviruses are labelled and shown as white rectangles.

Alpha

Simplexvirus		
TR. UL12 UL5-156 UL12-21 UL22-25 UL27-29 UL31-39 UL59-56 IR. IR. TR.		
gL, UDG gM HB MCP gH, DNA Pol, CH Trans		
Varicellovirus		
TR. 16-27 20-31 33-37 40-41 42-55 59469R.IRs TRs		
Beta $1 2 4 5 6 7$		
Cytomegalovirus		
TR. 01.22-33 01.45-63 01.59-72 01.73-80 01.85-87 01.89-405 01.112-117 IR. IR.	AT	4
Beta-genes 1 2 3 4 5 6 7 A		4
TR. U12-19 U27-37 U59-41 U12-45 U49-45 U49-45 U49-45 U49-45		
Gamma	Beta-ger	185
Lymphocryptovirus	Block 2	
	Bock 4	
, Rhadinovirus	Block 7	
2 4 5 50 3 100 150	200	230 kb

Figure 1.2 Conserved gene blocks of the Alpha-, Beta-, and Gammaherpesvirinae. Human herpesvirus-1 (HHV-1) and HHV-2 belong to the genus Simplexvirus; HHV-3, Varicellovirus; HHV-4, Lymphocryptovirus; HHV-5, Cytomegalovirus; HHV-6 and HHV-7, Roseolovirus; HHV-8, Rhadinovirus. The seven conserved herpesvirus sequence blocks (block 1 through block 7), plus an additional sequence block (beta-genes) found only in betaherpesviruses, are diagramed. ORFs bounding the conserved sequence blocks are indicated. Black arrows, gene blocks inverted relative to the betaherpesviruses; TRL, IRL, IRS, and TRS, inverted repeats flanking the L or S component, respectively; Hatched arrows, directionality of terminal repeats. Gene product abbreviations: gL, glycoprotein L; UDG, uracil-DNA glycosylase; PK, phosphotransferase; Exo, alkaline exonuclease; gM, glycoprotein M; HP, helicase/primase complex; MCP, major capsid protein; gH, glycoprotein H; AP, protease; DNA Pol, DNA polymerase; gB, glycoprotein B; PP, polymerase-associated processivity

HHV-7 ORF (strain *)		Homologues ^b		Producted function	Ref ^c
(JI)	(RK)	HHV-6	HCMV		1.61
H1				Unlikely to encode protein	2
DR1	DR1 ex1	DR1		US22 gene family	1,2
DR2	DR1 ex2	DR2		US22 gene family	1,2
H2				Unlikely to encode protein	2
DR6	DR6 ex1	DR6		US22 gene family	1,2
DR7	DR6 ex2	DR7		US22 gene family	1,2
НЗ				Unlikely to encode protein	2
H4				Unlikely to encode protein	2
U2	U2	U2	UL23	US22 gene family	1,2
U3	U3	U3	UL24	US22 gene family	1,2
U4	U4	U4	UL27	Related to U7 exon 2	2
U5/7	U7 ex2	U5	UL27	US22 gene family Related to U4	1,2 2
	U7 ex1	U7	UL28	US22 gene family	1,2
U8	U8	U8	UL29	US22 gene family	1,2
U10	U10	U10	UL31	Nuclear phosphoprotein	3
U11	U11	U11	UL32	Structural phosphoprotein	1,2
U12	U12	U12	UL33	G-protein coupled receptor Chemokine receptor	1,2 1
U13	U13	U13		-	1,2
U14	U14	U14	UL35	HCMV UL25/35 gene family, antigenic HHV-7 pp85 Tegument phosphoprotein	4 3
U15	U15	U15		HCMV UL25/35 gene family	1,2
U16	U17 ex2	U16	UL36x2	IE-B, transactivator US22 gene family	1,2 2
U17Ex	U17 ex1	U17Ex	UL36x1	IE-B, transactivator US22 gene family	1,2 2
U17A		U17A		Unlikely to encode protein	2
U18				IE-B, homologue to HCMV IE protein	1
U18	U18	U18	UL37x3	IE-B membrane glycoprotein	2
U19	U19	U19	UL38	IE-B protein	1,2
U20	U20	U20		Probable membrane glycoprotein lg gene family?	1,2 1
U21	U21	U21		Probable membrane olycoprotein	12
U23	U23	U23		Probable membrane glycoprotein EHV-1 gJ homologue	1,2 1
U24	U24	U24		Glycoprotein Contains hydrophobic domain	1
U24A	U24A	U24A		Contains hydrophobic domain	2
U25	U25	U25	UL43	US22 gene family Transactivator	1,2 1
U26	U26	U26		-	1.2
U27	U27	U27	UL44	DNA polymerase processivity factor	1.2
U28	U28	U28	UL45	Ribonucleotide reductase (large subunit)	1.2
U29	U29	U29	UL46	Minor capsid protein (mCP) Component of intercapsomeric triplex	1,2 2
U30				Capsid assembly, myosin	1
1130	1130	1130	111.47	Torumont protoin	,

Table 1.3 Predicted functions of HHV-7 ORFs and betaherpesvirus homologues

HHV-7 (ORF (strain *)	Homoic	aues ^b		
(JI)	(RK)	HHV-6	HCMV	Predicted function	Ref °
<u></u>	 U31	U31	UL48	Large tegument protein	1,2
U32	U32	U32	UL48.5	Capsid protein (located on tips of hexons)	2
U33	U33	U33	UL49	Virion protein	1,2
000	000	•••		Probable virion protein	1
U34	U34	U34	UL50	Membrane-associated phosphoprotein	2
1135	1135	U35	UL51	Role in DNA packaging	2
000	000	000	0201	Probable virian protein	1
U36	U36	U36	UL52	Role in DNA packaging	2
1137	1137	1137	111.53	Putative phosphoprotein	4
007	001	007	0200		1
U38	U38	U38	UL54	Catalytic subunit of DNA polymerase	2
U39	U39	U39	UL55	Envelope glycoprotein B (gB)	1,2
				Transport protein TP	1
U40	U40	U40	UL56	Role in DNA packaging	2
				Major DNA-binding protein	1,2
U41	U41	U41	UL57	Single stranded DNA-binding protein	2
1140	1140	1140	111.60	Transactivator	1
042	042	042	OL09	Post-translational regulator of gene expresson	2
U43	U43	U43	UL70	Component of DNA helicase-primase complex, primase	1,2
U44	U44	U44	UL71	-	1,2
U45	U45	U45	UL72	dUTPase	1
U46	U46	U46	UL73	Membrane protein	2
U47	U47	U47	UL74	Envelope glycoprotein O (Imanishi et al.)	1,2
U48	U48	U48	UL75	Envelope glycoprotein H (gH)	1,2
U49	U49	U49	UL76	Fusion protein	1
U50	U50	U50	UL77	Virion protein	1
				Role in DNA packaging	2
U51	U51	U51	UI 78	G-protein coupled receptor	1,2
			02.0	OpioidR homologue	1
U52	U52	U52	UL79		1,2
U53	U53	U53	UL80	Protease/assembly protein	1,2
				N-terminal protease domain	2
	U53.5			Major capsid scaffold protein	2
U54	U54	U54	UL82/83	Tegument protein transactivator	1
				Virion transactivator	2
U55A	U55A	U55	UL84	Probable replication function	1
				Related to U55B	2
U55B	U55B	U55	UL84	Probable replication function	1
				Related to USSA	2
U56	U56	U56	UL85	Capsid protein	1,2
				component of intercapsomenc inprex	2
U57	U57	U57	UL86	Major capsid protein (MCP)	1,2
	1150				2
058	008	058		-	1,2
008	009	059	UL88	•	1,2
U60	U66 ex2	U60	UL89x2	Late spliced gene (U60/U66) DNA packaging	1,2
1100	1100			r uauve terminase	2
062	062	U62	UL91	•	1,2
003	003	063	UL92	-	1,2
004	004	064	0193	Role in DNA packaging, tegument protein	2

HHV-7 ORF (strain *)		Homologues ^b			Def ⁶
(JI)	(RK)	HHV-6	HCMV	Predicted function	Ret
U65	U65	U65	UL94	Tegument protein	2
U66	U66 ex1	U66	UL89x1	Late spliced gene (U60/U66) DNA packaging Putative terminase	1,2 2
U67	U67	U67	UL95	-	1,2
U68	U68	U68	UL96		1,2
U69 U69	U69	U69	UL97	Phosphotransferase Serine-threonine protein kinase, tegument protein	1 2
U70	U70	U70	UL98	Alkaline exonuclease Deoxyribonuclease, maturation/packaging of DNA	1 2
U71	U71	U71		Myristylated tegument protein	2
U72	U72	U72	UL100	Integral membrane protein (gM) Role in virion envelopment	1,2 2
U73	U73	U73		Origin-binding protein (OBP) Helicase	1,2 2
U74	U74	U74	UL102	Component of DNA helicase/primase complex	1,2
U75	U75	U75	UL103	-	1,2
U76	U76	U76	UL104	Probable virion protein Minor capsid protein, DNA packaging	1 2
U77	U77	U77	UL105	Component of DNA helicase/primase complex Helicase	1,2 2
H5				Unlikely to encode protein	2
U79	U79 ex1	U79	UL112	HCMV replicaton, spliced (UL112/UL113) Probable role in DNA replication	1 2
H6	U79 ex2			HHV-6 U79 homology (C terminus) Probable role in DNA replication	1 2
U80	U79 ex3	U80	UL113	HCMV replicaton, spliced (UL112/UL113) Probable role in DNA replication	1 2
U81	U81	U81	UL114	Uracil-DNA glycosylase	1,2
U82	U82	U82	UL115	Envelope glycoprotein L (gL)	1,2
U84	U84	U84	UL117	Spliced in HCMV	1
U85	U85	U85		Probable membrane glycoprotein, related to OX-2	1,2
U86	U86	U86	UL122	IE-A protein, HCMV IE2 homology	1,2 1
U89	U90 ex3	U89		IE-A transactivator	1,2
U90	U90 ex2	U90		IE-A transactivator	1,2
U91	U91	U91		Probable membrane glycoprotein	2
H7				Unlikely to encode protein	2
U95	U95	U95		US22 gene family, MCMV IE2 homolog	1,2 1
H8	U100 ex10			Envelope glycoprotein gp105	2
U98	U100 ex3	U98		Envelope glycoprotein gp105	2
U99	U100 ex2	U99		Homology to HHV-6 gp82/105	1,2
U100	U100 ex1	U100		Homology to HHV-6 gp82/105	1,2
H1 [.]				Unlikely to encode protein	2
DR1'	DR1' ex1	DR1'		US22 gene family DR1/DR6 homology	1,2 1
DR2'	DR1' ex2	DR2'		US22 gene family	1,2
HZ				Unlikely to encode protein	2
DR6'	DR6' ex1	DR6'		US22 gene family DR1/DR6 homology	1,2 1

HHV-7 ORF (strain *)		Homologues ^b	Des dists d fue stice	
(JI)	(RK)	HHV-6 HCMV		
DR7'	DR6' ex2	DR7'	US22 gene family Transactivator	1,2 1
H3'			Unlikely to encode protein	2
H4'			Unlikely to encode protein	2

^a HHV-7 strain JI, Nicholas *et al* 1996; RK, Megaw *et al* 1998

^b HHV-6, Gompels et al 1995; HCMV, Chee et al 1990

° Reference 1, Nicholas et al 1996; 2, Megaw et al 1998; 3, Menegazzi et al 1999; 4, Pellet and Dominguez 2001

1.1.5 Gene expression

Lytic expression of HHV-7 genes appears to follow the temporal regulated pattern (Menegazzi et al., 1999), common to the herpesviruses (Figure 1.3). Herpesvirus genes can be classified into immediate early (IE or α), early (E or β) or late (L or γ) genes, based on their patterns of regulated expression (Chambers et al., 1999; Honess and Roizman, 1974; Mirandola et al., 1998; Wagner et al., 1972). IE genes are expressed immediately-early during infection independently of *de novo* protein synthesis. They provide transcription regulatory functions for the E genes. Transcription of the E genes rely on the IE genes but not viral DNA replication, and their products are often associated with viral DNA replication functions. L genes often encode structural proteins, and their expression is almost completely dependent on viral DNA replication. A few of the HHV-7 genes have been classified into their kinetic classes using inhibitory drugs that block protein synthesis (cycloheximide and emetine) and DNA replication (phosphonoacetic acid) (Menegazzi et al., 1999). More details on HHV-7 and herpesvirus gene expression are given in Chapter 5.



Figure 1.3 Temporal regulation of herpes simplex virus (HSV) gene expression. After virion binding, internalisation and nuclear entry, the viral DNA circularizes and is transcribed by host RNA polymerase II to give first the IE or α mRNAs. Five of the six HSV IE proteins act to regulate viral gene expression in the nucleus. They transactivate E or β gene transcription. The E proteins are involved in replicating the viral DNA molecule. Viral DNA synthesis stimulates L or γ gene expression. The L proteins are involved in assembling the capsid in the nucleus and modifying the membranes for virion formation. The mature enveloped virion exits the cell. Tegument proteins vhs (virion host shutoff) and VP16 are involved in degrading mRNA and activating IE gene expression respectively. (Reproduced from Roizman and Knipe, 2001)

1.1.6 Cell entry and Receptor

The cell surface CD4 molecule is believed to be the receptor for HHV-7 infection, supported by various experimental findings. It has been demonstrated that, when several non-susceptible cell lines, Jurkat (T-cell line), U937 (monocyte), SAS413 and K562 (myelocytes) were transfected with CD4, they became susceptible to HHV-7 infection, as measured by cell morphology and indirect immunofluorescence (Yasukawa et al., 1997; Zhang et al., 2000). Radiolabelled HHV-7 virus binds specifically to CD4transfected HeLa cells but not to unmodified HeLa cells (Lusso et al., 1994). HHV-7 infection was inhibited in a dose-dependent manner by preincubation with soluble CD4, anti-CD4 monoclonal antibodies, or gp120, the viral ligand for CD4 in human immunodeficiency virus (HIV) infection of CD4⁺ T cells (Lusso et al., 1994; Yasukawa et al., 1997). Flow cytometry studies showed that surface CD4 is reduced in HHV-7infected PBMCs and SupT1 cells, while other T-cell markers CD2, CD3 and CD44 remained unchanged (Lusso et al., 1994; Zhang et al., 2000). Downregulation of receptor molecules (CD4 and CCR5) during infection has also been observed in HIV (Piguet et al., 1999), possibly as a viral strategy to prevent superinfection (Little et al., 1994; Mitchell et al., 2003), and is believed to play an important role in HIV replication (Levesque et al., 2004; Lundquist et al., 2002). The above experimental results therefore conclude that CD4 is essential for HHV-7 infection of CD4⁺ T cells.

To date, the viral ligand for CD4 has not been identified, and the binding site on CD4 has not been mapped. However, clues can be obtained by comparing HHV-7 and HIV-1 interaction with CD4. HHV-7 infection is blocked by a monoclonal antibody which binds to the V3/V4 domains of CD4, as well as by those that bind to the V1 region (Lusso et al., 1994); HIV-gp120 binding to CD4 is via the V1 domain, unaffected by anti-V3/V4 antibodies. Moreover, pre-exposure of live or UV-inactivated HHV-7 inhibits HIV-1 infection in macrophages (Crowley et al., 1996). Therefore the HHV-7 binding site on CD4 is probably different to that of HIV-1, but may span multiple domains so as to block HIV-1 binding. Understanding HHV-7/CD4 interactions may provide clues for further antiviral strategies against HIV-1.

Although CD4 is essential for HHV-7 infection, many CD4⁺ T-cell lines are either only weakly or not at all susceptible to the virus (Yasukawa et al., 1997), suggesting that CD4 alone is not sufficient for HHV-7 infection. The search for an alternative or co-receptor led to the identification of cell surface proteoglycans as potential candidates. Heparin, the natural ligand of heparan sulphate proteoglycans, inhibits HHV-7 binding to SupT1, formation of syncytia and viral replication (Secchiero et al., 1997c), suggesting heparan sulphate may play a role in HHV-7 infection. Several HHV-7 proteins have been proposed as viral ligands for attachment to cell surface proteoglycans, including glycoprotein B encoded by U39 (Secchiero et al., 1997c) and gp65 encoded by U98/100 (Skrincosky et al., 2000). Heparan sulphate proteoglycans have been reported to be the cellular receptors for alphaherpesviruses (Spear and Longnecker, 2003), herpes simplex virus (Campadelli-Fiume et al., 2000), CMV (Compton et al., 1993), VZV (Zhu et al., 1995), and Papillomavirus (Shafti-Keramat et al., 2003).

The role of CXCR4 as a co-receptor has also been studied with inconclusive results. Two studies reported downregulation of surface CXCR4 during HHV-7 infection (Secchiero et al., 1998c; Yasukawa et al., 1999), whereas another study showed that CXCR4 transcription is not modified by HHV-7 and does not act as a co-receptor (Zhang et al., 2000). More research is needed to strengthen our understanding of the molecular interactions during HHV-7 entry of CD4⁺ T cells.
1.1.7 Cell tropism & Persistence

HHV-7 has a restricted cell tropism of T-lymphocytes *in vitro*, demonstrated by double immunofluorescence of infected cord blood mononuclear cells (CBMCs) (Berneman et al., 1992a). SupT1, a CD4⁺ T-lymphoblast cell line derived from non-Hodgkin's T cell lymphoma, is the only cell line shown to support sustained, productive HHV-7 infection *in vitro* (Ablashi et al., 1998; Berneman et al., 1992a), shown by CPE observation, IFA for viral antigen, and electron microscopy. On the other hand, a number of cell lines are shown to be non-susceptible to HHV-7 infection (no viral antigen by IFA and no viral DNA by PCR), including several CD4⁺ and CD4⁻ T-cell lines HSB2, MoLT-3, J-Jhan, MoLT-4, Jurkat (Ablashi et al., 1998); HUT-78, MT-4, (Zhang et al., 2000); EU-T-01, Molt-15, HO8-HT (Yasukawa et al., 1997); B-cell lines Daudi, Raji, MY-LCL, MJ-LCL (Yasukawa et al., 1997); and macrophages (Crowley et al., 1996). In contrast to Crowley et al, it has also been reported that HHV-7 is able to establish productive infection in macrophages up to 2 weeks post inoculation, while viral DNA could be detected for up to a month but not viral mRNA, therefore HHV-7 may establish latent infection in macrophages (Zhang et al., 2001a).

The *in vivo* reservoirs of HHV-7 are not known, but the virus is frequently isolated from peripheral blood (Fernandez et al., 2002) and saliva (Wyatt and Frenkel, 1992). Viral DNA is present in 55 to 100% of salivary gland specimens (Sada et al., 1996; Yadav et al., 1997). HHV-7 has also been detected by immunohistochemistry in other tissues including lung (7/8), skin (7/11), mammary glands (4/7), liver (5/8), kidney (8/10) and tonsils (2/6) (Fujisaki et al., 1998; Kempf et al., 1998). However, only a small proportion of the these samples (lung, 1/8; skin, 2/11; and mammary glands, 2/7) also scored positive for viral DNA in PCR, compared to salivary glands (6/10). Tissue samples of brain, large intestine and spleen were HHV-7-negative.

1.1.8 Transmission & Prevalence

HHV-7 is a highly prevalent virus, with over 90% adults being seropositive to the virus (Clark et al., 1993; Wyatt et al., 1991). Primary infection is thought to occur in early

childhood. Some data showed that, like HHV-6, antibodies to HHV-7 were detected in infants under 2 years old (Clark et al., 1993), whereas others suggest a later seroconversion in 3 to 4 year olds (Wyatt et al., 1991). Another study reported HHV-7 prevalence peaks at 60 % in 11-13 year olds (Yoshikawa et al., 1993). Infectious HHV-7 virus has been readily isolated from saliva of healthy adults (Wyatt and Frenkel, 1992; Yoshikawa et al., 1993), suggesting the saliva plays a role in spreading the virus. It has been shown that viral DNA isolated from the saliva of children shared similar restriction patterns to their parents, providing further proof for the horizontal transmission of HHV-7 via close contact between parent and child (Takahashi 1997). In addition, breast feeding may also be a possible transmission route, as viral DNA has been detected in breast milk, and the seropositivity rate is slightly higher in children who are breast-fed than bottle-fed (Fujisaki et al., 1998).

1.1.9 Disease association & Immune response

The ubiquitous nature of HHV-7 makes it difficult to identify disease associations. Studies have linked the virus to exanthem subitum (ES) (Tanaka-Taya et al., 2000; Torigoe et al., 1995; Ueda et al., 1994), a childhood illness characterised by fever and skin rash, possibly because HHV-7 reactivates HHV-6 (Tanaka-Taya et al., 2000), the primary causal agent of ES (Yamanishi et al., 1988). There are controversial views on the role of HHV-7 in pityriasis rosea (PR, a relapsing rash). One study detected HHV-7 DNA in plasma, PBMC and skin samples from patients with PR, but not in controls (Drago et al., 1997), whereas another study concluded there was no significant difference between the prevalence of HHV-7 in PR skin lesion samples and control samples, either by nested PCR for HHV-7 DNA, or by IFA for antigen (Kempf et al., 1999). HHV-7 may be an opportunistic pathogen responsible for transplant complications, for example delayed neutrophil engraftment in bone marrow recipients (Chan et al., 1997). It may also have associations with CMV disease in liver (Mendez et al., 2001), kidney (Osman et al., 1996) and bone marrow (Chan et al., 1997) transplant patients. At least some of these correlated with HHV-7 viraemia and/or DNA load (Brennan et al., 2000; Chan et al., 1997; Mendez et al., 2001). There are no significant links between HHV-7 and chronic fatigue syndrome (CFS) (Ablashi et al., 1998), and the isolation of HHV-7 from CFS patients may be the consequence of CFS

rather than cause (Secchiero et al., 1994). In summary, to date no disease has been confirmed to be caused by HHV-7, with the virus apparently very well adapted to persistence and transmission in the human host.

There are very few published studies on the host immune response to HHV-7, possibly due to the virus's lack of pathogenicity. However, information can be gathered from the data of serological prevelance studies. Primary infection by HHV-7 leads to the production of HHV-7–specific antibodies in 40% of healthy children by 44 months (Wyatt et al., 1991). Antibody titres to HHV-7 in children are high in the first 2 months (94.4%), possibly due to presence of maternal antibodies, followed by a decline between 6-7 months (30%). A rise in HHV-7 antibodies after 8 months (Clark et al., 1993) was possibly due to the infected infants producing their own antibodies to the virus after the reduction in maternal antibodies. These antibodies are apparently protective against the virus, as sera from healthy individuals and CFS patients who are HHV-7–positive inhibit HHV-7–induced CPE *in vitro* (Secchiero et al., 1994). HHV-7 antibodies are sustained into adulthood, as sera from 24 out of 26 (92%) healthy adults are seropositive for HHV-7 (Wyatt et al., 1991).

In addition to a humoral response, it has also been observed that $CD4^+$ T-cells produce gamma interferon (IFN γ) and show a proliferative response to HHV-7 antigen (Yasukawa et al., 1993). Specificity tests showed that 28% of the T-cell clones reacted against both HHV-6 and HHV-7. Since primary infection by HHV-7 seem to occur after HHV-6 (Wyatt et al., 1991), perhaps the immune system is already equipped for protection against HHV-7 upon primary infection. In addition, during latency, HHV-7 probably evades the immune system by minimising viral gene expression (Menegazzi et al., 1999).

1.1.10 Cellular interaction

It would not be surprising that some of the HHV-7 gene products interact with the host. HHV-7 induces a range of effects in infected cells. Examples include the downregulation of the cellular receptor CD4 (Furukawa et al., 1994; Lusso et al., 1994; Secchiero et al., 1997b) and downregulation of CXCR4 (Secchiero et al., 1998c). HHV-7 infection of CD4⁺ T cells also dysregulates expression and activity of cyclin B and the cyclin-dependent kinase cdc2, leading to cell cycle arrest and polyploidisation of infected cells (Secchiero et al., 1998a). These interactions may be involved in apoptosis regulation and virion production (Secchiero et al., 1998b). A possible immune evasion mechanism has also been observed; HHV-7 infected cells had increased levels of Tumour necrosis factor (TNF)-related apoptosis-inducing ligand (TRAIL) but decreased surface TRAIL receptor TRAIL-R1. As a result while bystander cells were killed by TRAIL-mediated cytotoxicity, HHV-7 infected cells remained resistant (Secchiero et al., 2001). The underlying molecular mechanisms of these effects are not characterised, and the participation of any HHV-7 genes or proteins remains unknown. Although the HHV-7 genome has not been extensively studied, information can be deduced from their homologues in other herpesviruses, especially HHV-6 and HCMV. Genes in the HHV-6 IE-A and IE-B loci, DR6, U3 and U27 are reported to have transcription regulatory functions. It has been shown that viral promoters such as HIV-1 LTR (Geng et al., 1992) and EBV EBNA-2 (Cuomo et al., 1998), in addition to the cellular promoter for CXCR4 (Yasukawa et al., 1999) are affected by HHV-6 regulators. HHV-7 U39 encodes the glycoprotein gB, which binds to cell surface heparin sulphate proteoglycans (Secchiero et al., 1997c), and may play a role in attachment and penetration, as in HHV-6 (Takeda et al., 1996). Glycoproteins gH and gL, encoded by U48 and U82 respectively, may be involved in membrane fusion, at least in HHV-6 (Qian et al., 1993). HHV-6 has also been reported to simultaneously stimulate host protein synthesis and shut down DNA synthesis (Pellett, 2001), thus it is possible that HHV-7 may induce changes of similar scale in the host during infection.

1.1.11 Life cycle

The life cycle of HHV-7 has not been extensively studied, but can be deduced from other properties of the virus, as mentioned in the previous sections. It is widely accepted that HHV-7, like all herpesviruses, persists in the host by a combination of lytic and latent infection, possibly in various sites or tissue types (Black and Pellett, 1999).

1.1.11.1 Latency/Persistence

After acute infection, the virus probably establishes latency in T-cells, as no expression of viral lytic genes were detected by RT-PCR in PBMCs isolated from patients despite having a high viral load (Menegazzi et al., 1999). Moreover, virus can be isolated from activated but not from resting PBLs, suggesting the virus may be reactivated from latency under T-cell activating conditions (Frenkel et al., 1990). Perhaps HHV-7 from activated T-cells is disseminated in the host via blood (Fernandez et al., 2002), and establishes a persistent infection in various other tissues, where viral DNA has been detected (section 1.1.7). Finally, HHV-7 persistence may be maintained by lytic production in salivary glands, as the virus is readily shed in saliva (Kempf et al., 1998; Wyatt and Frenkel, 1992; Yoshikawa et al., 1993).

1.1.11.2 Lytic infection

Lytic infection begins with reactivation or with viral attachment to specific surface receptors and penetration of host cells (section 1.1.6). HHV-7 encodes the necessary genes for replication and virion assembly (section 1.1.4), and the regulated expression of these genes (section 1.1.5) drives the production cycle of the virus. Electron microscopy (Black et al., 1997) detected DNA-containing capsids in the nucleus 3 days after infection. Capsids appear to acquire tegument in the perinuclear space, seen as a pool of electron-dense material surrounding the capsid. Tegumented capsids can be observed in the cytoplasm, and probably acquire the envelope by budding upon the golgi membrane. Matured virions are released into extracellular space after 5 days, causing cell death by apoptosis and necrosis *in vitro* (section 1.1.3).

In conclusion, HHV-7 is an efficient virus, establishing persistent infection in most of its human host population, whilst causing minimal disease, avoiding host immune response, and maintaining the ability to replicate. Studying HHV-7 may therefore reveal more detail about how the human host can tolerate the presence of commensal viruses.

1.2 DNA Microarrays

1.2.1 The rise of the genome, transcriptome, and proteome

In recent years, the advantages in studying gene expression using microarray technology have been widely acknowledged (Harrington et al., 2000; Lockhart and Winzeler, 2000). This technology excelled due to its high throughput nature, reproducibility and relative ease of use. The expansion of microarray technology can also be attributed to its parallel development with three technical advancements in biomedical science in the recent years. These are the rapid generation of genomic data from various genome projects, development in automated precision robotics and the improvement in computational data management.

To date, the complete genome sequences of many organisms have been elucidated, including the human genome (McPherson et al., 2001). Genomes of many pathogens have been sequenced, including 217 bacterial genomes (http://www.tigr.org), 799 virus genomes (http://www.ncbi.nlm.nih.gov/PMGifs/Genomes/10239.html), as well as many protozoan, fungus and helminth species (http://www.sanger.ac.uk). A large number of genome projects are currently ongoing. Many of the completed genome sequences have been annotated for potential sites of expressed open reading frames (ORFs), transcription initiation site, promoter sequences and splicing information. However, genome sequence data alone do not provide information about what the genes do (prediction by sequence homology is limited to the number of genes of which the functions we already know), how their expression is regulated, or what roles they play in the biological processes in coordination with other genes. The conventional methods for gene analysis do not have the capacity to match the vast quantity of data generated, hence there is a need for large scale analysis of genome data. Microarray technology provides a bridge for this gap. This high throughput technology allows many genes to be analysed simultaneously and rapidly.

Early arrays contained only several hundred elements printed on nylon membranes. The production of high density solid substrate microarrays relied upon automated precision robotics technology to deposit tens of thousands of elements onto the substrate surface (DeRisi et al., 1997; Schena et al., 1995). Therefore the advancement in precise and high density robotic deposition facilitated the development of microarray technology.

Good data management systems are required for many stages in microarray experiments (Chicurel, 2002). First of all, accurate genomic data available in genome databases have facilitated the production of array elements of specific sequences (Kellam, 2001). Annotated genomic databases have also been developed, for example, VIDA is a specialist database for viruses (Alba et al., 2001) that organise ORFs from many complete viral genome sequences into homologous protein families. A set of standards - Minimal Information About a Microarray Experiment (MIAME) (Brazma et al., 2001) has been created for the submission of microarray results to the public repository (<u>http://www.ebi.ac.uk/arrayexpress/</u>), providing a interface for scientists to access and compare microarray data. These computational advances therefore played important roles in the development of high-density microarrays.

The high throughput nature of microarrays made them ideal for genome-wide analysis applications. Comparative genome hybridisation (CGH) is a term given for using microarrays in genotyping (Hacia, 1999; Stoughton, 2005) and single nucleotide polymorphism (SNP) studies (Shen et al., 2005). CGH arrays have been developed and extensively applied to investigate genetic diversity in a number of bacterial systems including Helicobacter pylori, Campylobacter jejuni, Yersinia pestis, Mycobacterium tuberculosis, and Streptococcus pneumoniae (Hakenbeck et al., 2001; Hinchliffe et al., 2003; Kato-Maeda et al., 2001; Salama et al., 2000). In a study of 15 virulent Helicobacter pylori strains, 362 out of 1643 genes analyzed were identified as strainspecific (Hakenbeck et al., 2001; Hinchliffe et al., 2003; Kato-Maeda et al., 2001; Salama et al., 2000). A similar genotyping analysis comparing 22 strains of Yersinia pestis and 10 strains of Yersinia pseudotuberculosis identified a number of differences that may relate to the evolutionary pressures on these species in their respective niches (Hinchliffe et al., 2003). An oligonucleotide array containing 82 polymorphic oligonucleotides of the 16S ribosomal RNA gene has been used for genotyping of polymorphisms and identification of mycobacterium species (Troesch et al., 1999). CGH arrays are not limited to bacterial studies. DNA microarrays have been made for genotyping of the amoeba parasite Entamoeba histolytica (MacFarlane et al., 2005a),

and to identify deletions in BCG vaccine strains (Behr et al., 1999). Wang *et al* developed an oligonucleotide (70-mer) microarray to detect 140 different viruses using known sequence data (Wang et al., 2002a).

Perhaps one of the most widely used applications of microarrays is the monitoring of gene expression (section 1.2.4). By means of measuring abundance of all mRNA transcripts in a sample simultaneously, microarrays provide a global picture of gene expression pattern (the transcriptome), which shows not only the expression of many genes individually, but also the coordinated expression pattern of the different genes at a given condition. This often reveals the physiological state of the system, such as stages of cell or tissue development (Pang et al., 2005), or responses to external stimuli such as drug treatments (Gerhold et al., 2001; Gray et al., 1998; Scherf et al., 2000). Transcription studies have helped towards understanding the mechanisms of many diseases, including many cancers such as Hodgkin's lymphoma (Cossman, 2001), hepatocellular carcinomas (Graveel et al., 2001; Okabe et al., 2001), breast cancer (Murphy et al., 2005), colorectal carcinogenesis (Kitahara et al., 2001). However, the accessibility of tumour tissues presents a challenge to such microarray studies, as cancerous cells need to be separated from normal cells in order to give differentiated expression patterns. For example, Kitahara et al used microarrays to compare the transcription profiles of colorectal cancer cells to noncancerous colonic epithelia, separating them using laser-capture microdissection, and amplifying the RNA signal. The differential expression pattern observed indicated that certain processes were accelerated in the cancer cells (metabolism, production of reactive oxygen species), whereas regulation of cell cycle, transcription and mitosis were impared, compared to normal tissues. Microarrays have also offered insights into microbial infections (section 1.2.4) and facilitated identification of disease subphenotypes and diagnosis (Helmberg, 2001; Petrik, 2001). The diversity of microarray applications in gene expression profiling has been extensively reviewed (Heller, 2002; Stoughton, 2005).

Based on the observation that genes with similar function are often co-expressed (Bassett et al., 1999), gene expression profiling can be applied to predict functions of unknown genes that are expressed with characterised ones, facilitating the annotation of microbial genome, for example, in *Saccaromyces cerevisiae* (Eisen et al., 1998). More

details of microarray-based gene expression analysis of microorganisms will be given in section 1.2.4.

Gene expression experiments provide valuable information about the global pattern of transcription in a biological system, but they do not address post-translational modifications of the protein product. Various methods are used in proteome studies, such as mass spectrometry, 2-dimentional (2d-) gel electrophoresis and protein chips (Pandey and Mann, 2000). Proteomic mass spectrometry analysis of Plasmodium falciparum, the protozoan responsible for malaria, has been used to characterise 2400 proteins at four stages of the protozoan's life cycle (Florens et al., 2002). Protein chips containing 35 - 149 protein probes have been developed for several bacteria, such as Salmonella enterica (Cai et al., 2005), Helicobacter pylori (Das et al., 2005), and Neisseria meningitides (Steller et al., 2005). A protein chip has also been produced for yeast, which contains purified proteins over-expressed from 5800 open reading frames of the yeast genome (Zhu et al., 2001). Miller et al. used protein arrays with cellular genes to identify potential prostate cancer biomarkers in prostate cancer serum (Miller et al., 2003). However, although proteomics targets the 'final product' of gene expression, they are more difficult to conduct, and possibly less robust than DNA microarrays (Lockhart and Winzeler, 2000). The 2d-gel approach is limited by dynamic range hence only efficient for the analysis of a subset of proteins (Pandey and Mann, 2000). Production of protein chips is more complex and costly than DNA microarrays, because proteins have a high variety of binding chemistries, lower stability, and often require multimerization and post-translational modifications. A lack of simple amplification processes like PCR also makes it more difficult to generate large quantities of protein (LaBaer and Ramachandran, 2005). Therefore currently protein arrays are of smaller scale and diversity than the DNA microarrays.

1.2.2 Principles of DNA microarray technology

Microarray technology is a large-scale competitive hybridisation technique that measures the relative abundance of many DNA or RNA species in parallel. A microarray essentially contains a large set of nucleic acid probes of known sequences immobilised onto a solid support, known as the substrate. The sample in question (a mixed population of DNA or RNA) is labelled, usually with fluorescence or radioactivity, and applied to the microarray. The different species of DNA or RNA in the sample hybridise to their corresponding array probes of complementary sequences. The detected fluorescence or radioactivity of each hybridised probe is a measure of the relative abundance of that particular gene in the sample. There are many types of microarrays and experimental design varies, but all share the common principle of detecting abundance of many nucleic acids simultaneously using a large number of probes.

1.2.2.1 Microarray platforms

There are three main types (or platforms) of microarrays, based on substrate material and the nature of probes. They differ in a range of features but for simplicity we will call them membrane arrays, spotted arrays and oligonucleotide chips in this thesis. Membrane arrays are the earliest form of microarray, using nylon membrane as the substrate on which only a few hundred DNA probes (PCR product or clones of a expression library) are deposited (Ahn et al., 2002; Jenner et al., 2001). Membrane arrays are usually used with radioactive-labelled samples. On the other hand, both spotted arrays and oligonucleotide chips use glass as substrate, which allows for more densely-arranged probes thus have a higher capacity (10,000s of probes per glass slide) than membrane arrays. Spotted arrays and oligonucleotide chips are both used to detect fluorescent-labelled samples but they differ in the nature of probes and method of probe generation. Spotted arrays are made in a similar manner to membrane arrays. DNA probes are pre-made (by cloning or by PCR) and then deposited or 'spotted' onto the substrate. The probes are generally longer (200-2000 bases), although oligonucleotide probes as small as 50-mer has been described (Kane et al., 2000). In contrast, oligonucleotide chips are made by *in situ* synthesis of oligonucleotide probes 25-mer to 60-mer in length, and usually more than one probe per gene are present on the array (Harrington et al., 2000).

The presence of multiple probes per gene gives oligonucleotide chips the ability to detect splice variants and to cover the entire genome, rather than just the predicted open reading frames (Lockhart et al., 1996; Pease et al., 1994). However, generating oligonucleotide probes in situ is costly, time consuming, and can result in products of variable yield. On the other hand, deposition of probes onto spotted arrays is faster, cheaper and can be easily implemented in non-commercial research laboratories, as in the case of many viral arrays (Chambers et al., 1999; Paulose-Murphy et al., 2001; Stingley et al., 2000). It has been shown that the sensitivity of a 50-mer oligonucleotide chip was comparable to that of spotted arrays with longer probes, but some of these 50mer probes cross-hybridised to sequences that share a 75-80% similarity with the target sequence (Kane et al., 2000), suggestive that the shorter probes may have lower specificity. To improve specificity, short oligonucleotide chips normally contain mismatched sequences as controls (Lockhart et al., 1996). Nonetheless, some laboratories have created spotted arrays with shorter oligonucleotide probes (Chambers et al., 1999; Kane et al., 2000; Okamoto et al., 2000; Stingley et al., 2000), demonstrating the convenience of spotted arrays and the value of shorter probes, especially for applications that require multiple probes per gene, such as single nucleotide polymorphism studies (Okamoto et al., 2000). Many studies have shown that specificity and sensitivity increase with probe size and the optimal probe length is 60 - 70-mer, with only modest increases above this size.

A number of studies have compared microarray data generated using different array platforms with contradictory conclusions. Some publications showed that data obtained with different array platforms were agreeable (Kane et al., 2000; Yuen et al., 2002); others reported little correlation between platforms (Kothapalli et al., 2002). However, recent studies argued that when care was taken to compare relative expression rather than absolute values, the patterns of expression were similar on both platforms (oligonucleotide chips and spotted arrays) even though the relative amplitude of change was greater in one platform than the other (Irizarry et al., 2005; Larkin et al., 2005). It was also reported that different array platforms actually had *less* effect on the observed

expression variations than both biological differences of the samples (Larkin et al., 2005) and the laboratories in which the experiments were conducted (Irizarry et al., 2005). When discrepancies were observed between oligonucleotide chips and spotted microarrays, quantitative RT-PCR results supported neither platform but gave a third pattern (Larkin et al., 2005), suggesting neither platform was more accurate that the other, and the discrepancies might have been due to splice variants. These findings concluded that the choice of array platform will not have significant effect on the accuracy of data obtained, providing careful data analysis was performed. For our study we have chosen to deposit PCR-amplified DNA probes (200 - 2000 bases) onto spotted glass microarrays for their higher density than membrane arrays, also for the relatively low-cost production that is easy to set up in-house.

1.2.2.2 Microarray and gene expression analysis

The use of microarrays in the monitoring of gene expression is based on measuring transcript (mRNA) abundance. This requires extraction of total or mRNA from the sample of interest, which are then reverse transcribed into cDNA. The cDNA is labelled with fluorophors either directly during reverse transcription or by indirect linkage afterwards. For spotted arrays it is most common to use the two-colour labelling method, in which two cDNA samples are labelled as a pair, sample 1 with the red fluorescent Cy5 dye and sample 2 with the green fluorescent Cy3 dye. The pair are then mixed and co-hybridised onto one array. The array is then scanned and the fluorescence intensities measured at around 635 nm (Cy5) and 530 nm (Cy3) of each hybridised probe represent the relative abundance of the particular mRNA sequence in sample 1 and 2, respectively (Figure 1.4). This two-colour system is a very important aspect of the spotted microarray. Microarray experiments are often designed to investigate changes in gene expression over a time series (Ahn et al., 2002; Jenner et al., 2001), under different conditions, e.g. drug treatment vs untreated (Wilson et al., 1999), or to compare expression between two types of cells, e.g. normal versus tumour (Okabe et al., 2001). All of these require the use of more than one microarray and subsequently, comparing data from different arrays. Such comparison is subject to errors as each array hybridisation is unique and variations occur in many processes, such as the amount of DNA probe per spot, labelling efficiency and hybridisation efficiency

(Quackenbush, 2002). Such errors can be reduced using the two-colour system. Because each array probe is hybridised to two samples with different coloured labels, any array-based variations should have the same effect on both Cy5 and Cy3 samples. Therefore comparing the Cy5/Cy3 ratios of each probe across different arrays allows elimination of these errors. This can be achieved by labelling the common reference RNA with one colour (usually the Cy3) and the RNA of interest in the other (Cy5). The same Cy3-labelled reference is used to hybridise all arrays, hence acting as a control so that the Cy5/Cy3 ratios of one gene on different arrays can then be compared.

There are several options of binding chemistry for probe attachment on glass slides, including aminosilane-coating, epis-coating, conjugated coating. Hybridisation depends on the type of array and the binding chemistry (Zammatteo et al., 2002). Spotted glass arrays generally require denaturation and blocking prior to hybridisation to prevent non-specific binding. During the actual hybridisation, optimal temperature, time, and hybridisation buffer containing the correct ingredients are essential to optimise specific binding. The conditions for post-hybridisation washing are also important in order to maximise signal-to-background ratios. Hybridised arrays are scanned under lasers operating at around 635 and 530 nm to excite the fluorophors to produce fluorescent emissions, which are amplified by photomultiplier and recorded as images. These images are then analysed by imaging softwares to produce numerical data (Hegde et al., 2000) for data processing and analysis, discussed in the next section.



RNAs are isolated or amplified from reference and test samples, then reverse-transcribed into Cy3and Cy5-labelled cDNA, respectively. The labelled cDNAs are mixed and hybridised onto microarrays, during which the labelled species bind to the DNA probes of complemetary sequences. The array is then scanned under lasers to produce a combined fluorescent image for the two channels. Spot colour represents the relative abundance of the two labelled cDNA; green spot, higher in Cy3; reds spot, higher in Cy5, yellow spot, similar fluorescence in Cy3 and Cy5.



1.2.3 Analysis of microarray data

After hybridisation and scanning, a series of downstream processes need to be performed before expression patterns can be deduced. First of all, the hybridised spots are identified using spot-finding software, available both open-source reviewed in (Hegde et al., 2000) and commercially (Genepix Pro, Axon Instruments, USA). They operate by overlaying a numerical matrix that defines the position and identity of the array probes onto the scanned image, fitting a circlular 'spot' around each hybridised probe and then calculating the fluorescence signals of every spot. These data are then used in subsequent data processing, including the calculation of Cy5/Cy3 ratios, data filtering, log transformation and data normalisation, which can be performed using user-created spreadsheets, mathematical programs, or specialist softwares (Eisen et al., 1998; Saeed et al., 2003). There are many methods for data analysis, depending on the number of arrays used, the experimental design, and the question being asked. If global expression patterns are of interest, these can be deciphered using clustering methods (section 1.2.3.4). On the other hand, significantly altered genes can be identified by statistical tests (section 1.2.3.3).

1.2.3.1 Data filtering and transformation

Microarray spot-finding softwares store data for every spot, including those with very low hybridisation signals or very high background (i.e. noisy data), which if included in data analysis, could adversely affect the accuracy of interpretations. Therefore microarray data are often filtered to remove questionable or low-quality data, to ensure data integrity and accurate analysis. Normally data below a signal-to-noise threshold or a minimum intensity level are excluded from analysis.

The Cy5/Cy3 ratio is a useful way of representing data but it gives a prejudiced view of up-regulated and down-regulated genes. Genes with double the signal in Cy5 compared to Cy3 have a ratio of 2, whereas genes with twice as much fluorescent in Cy3 than Cy5 have a ratio of 0.5, despite the fold change is the same. When \log_2 transformed these ratios appear as +1 and -1, respectively, hence presenting up- and down-regulated genes

by the same magnitude fold-change. Log_2 transformation also converts the spread of data to normal distribution, thus allowing the use of parametric statistical methods on the data. It is a common pre-analysis adjustment of microarray data (Quackenbush, 2002).

1.2.3.2 Data normalisation

Microarray data need to be normalised before interpretation. The purpose for normalisation is to adjust the data so that the effects of experimental procedures on the data are removed as much as possible. For example, the efficiency of labelling and hybridisation of the two channels can never be exactly identical, resulting in systematic errors in the Cy5/Cy3 ratios. There are several widely used methods to reduce this dyebias effect, based on total intensity, regression, and ratio-statistics. All three approaches assume that the average Cy5/Cy3 expression ratio of all (or a subset of control) genes on the array should be equal to one. Total intensity normalisation assumes the total intensity of all (or a subset of) spots in one channel is equal to that of the other channel. A correcting factor is applied to each spot, so that the total log_2 Cy5_{total}/Cy3_{total} ratio is equal to zero (Quackenbush, 2002). Regression-based normalisation adjusts the intensities so that the best-fit slope of the Cy5/Cy3 ratio to one (Hedenfalk et al., 2001). Ratio statistics-based normalisation is an iterative method that adjusts Cy5/Cy3 ratios to normalise the mean Cy5/Cy3 ratio to one (Chen et al., 2002).

Another reason for data normalisation is the phenomenon of 'intensity-dependent local variation'. It has been observed that the log_2 ratios of low-intensity data generally deviate further from zero than the high-intensity spots. As a result these genes may be falsely classified as up- or down-regulated genes. To understand this phenomenon consider a high-intensity gene spot H and a low-intensity gene spot L. If gene H gave intensity signals of 2200 and 2000 in Cy5 and Cy3 channels respectively, the log_2 Cy5/Cy3 ratio is +0.13 and the expression of gene H would be regarded as relatively unchanged. On the other hand, gene L with signals of 400 and 200 will have a log_2 Cy5/Cy3 ratio of +1, and appear as an up-regulated gene, even though both genes have a small difference of 200 intensity units between the two channels. This intensity-dependent effect can be removed by excluding low intensity spots from analysis, and/or

can be corrected using local weighted scatterplot smoothing (LOWESS) (Yang et al., 2002b). LOWESS also corrects for other systematic spatial variations, such as spotting inconsistencies and uneven hybridisation. There are other methods that apply local corrections, for example, local variance regularisation normalises the variance of the data (Yang et al., 2002a). Expression data are also commonly re-scaled by mean- or median-centring, so that the average log₂ Cy5/Cy3 ratio of each gene across arrays is set to zero. This has the effect of enhancing the relative differences in expression ratios so that subsequent analysis will reveal the patterns rather than the abundance of gene expression (Quackenbush, 2001), as well as eliminating the effects of reference RNA. Specialist softwares are now available for microarray data processing. Some of these even allow users to perform a series of data filtering and normalisation methods in one package (Fielden et al., 2002; Saeed et al., 2003), which have greatly facilitated the processing of microarray data.

1.2.3.3 Identification of differentially expressed genes

One common application of microarrays is to identify differentially expressed genes between two samples, such as drug treated versus untreated cells (Wilson et al., 1999), or normal versus cancer tissue (DeRisi et al., 1996), where a binary system is a simple and efficient method. The two samples may be labelled with different fluorescent dyes and hybridised onto the same array, and the resultant expression ratios represent the changes in gene expression. Genes with log₂ ratios closest to zero have relatively unchanged expression, whereas genes with very highly positive or negative log₂ ratios are more abundant in one sample or the other. Such direct comparison requires dve swap hybridisations to eliminate the dye-bias effect. Alternatively, a common reference approach as mentioned above (section 1.2.2.2) can be used, and the log₂ ratios of one sample compared to another. In either case, replicate arrays should be performed to provide an estimate for the variance in the data (Yang and Speed, 2002), and to enable identification of statistically significant up- or down-regulated genes. Various statistical tests have been used for microarray data, such as analysis of variance (ANOVA) (Churchill, 2004; Kerr et al., 2000), Wilcoxon rank sum test (Chambers et al., 1999). The Significance Analysis of Microarrays (SAM) method, a t-test based approach to assign scores for all genes and to calculate the false discovery rate (Tusher et al., 2001),

have been widely used to identify differentially expressed genes (Jones and Arvin, 2003; Larsson et al., 2005).

1.2.3.4 Expression patterns recognition by clustering analysis

A widely-used approach to study array data is by clustering analysis. The arrangement of genes into clusters according to the similarity of their expression patterns is very useful because genes that are co-regulated (similar expression patterns) are often involved in similar biological processes (Eisen et al., 1998). The power of the technique is maximised by clustering the data from multiple arrays so that samples with similar expression patterns are also grouped together (Tsai et al., 2005). This allows for identification of co-regulated genes across a series of conditions, common or contrasting patterns in different tissue types, or the gradual change of expression at increasing times post-treatment (section 1.2.4.3). Clustering analysis normally consists of 3 steps. The first step is the calculation of the pairwise distance between the expression vectors of the array elements, which are essentially composed of the Cy5/Cy3 ratios of the gene spots. Secondly, clustering algorithms group the elements into clusters by means of their pairwise distance. Thirdly, a visual representation of the clusters is created, usually an expression matrix in which genes are arranged in rows and arrays in columns as a dendrogram, similar to a phylogentic tree, in which the branch lengths represent the distances between the array elements. Various distance metrics exist for the calculation of the distances between array elements. For example, Euclidean distance is a method based on the Pythagorean Theorem. The Manhatten distance is the sum of distances between components of each expression vector. Pearson correlation, a semi-metric distance, may be centred, uncentred or squared (Quackenbush, 2001).

There exist many different clustering algorithms, of which the most popular are hierarchical clustering (Eisen et al., 1998), k-means clustering 1999 (Soukas et al., 2000; Tavazoie et al., 1999), and self-organising maps (Tamayo et al., 1999). Different algorithms have different characteristics, and when used on the same data set will result in slightly different clustering patterns. However, none of these algorithms is more 'correct' than the others; they simply reveal different features of the data set, hence suited for different applications (Quackenbush, 2001). Hierarchical clustering (HCL) is an agglomerative approach, which gradually builds up a cluster by joining elements together based on their similarity. It begins by finding the two array elements with the smallest pairwise distance and joining them together. The distances between this newly joined group and each of the other elements or groups are calculated. The algorithm then repeats the 'search-and-group' process until all array elements are included in the cluster, forming a dendrogram (Quackenbush, 2001). Eisen et al first developed the programs Cluster and Treeview for the application of hierarchical clustering to microarray data analysis (Eisen et al., 1998), which became the standard analysis protocol. HCL has been used in analysing gene expression data of diseases such as psoriasis (Itoh et al., 2005), ischemic heart disease (Satterthwaite et al., 2005), heart failure (Steenman et al., 2005), primary biliary cirrhosis in liver (Honda et al., 2005), Parkinson disease (Hauser et al., 2005); in studying various cancers including HPVassociated cervical cancer (Vazquez-Ortiz et al., 2005), squamous cell carcinoma in lungs (Talbot et al., 2005). It has also been applied to analyse transcriptional profiling data of normal human tissues and organs (Shyamsundar et al., 2005), and microorganisms such as yeast (Eisen et al., 1998) and Streptococcus pneumoniae (Pandya et al., 2005). The advantages of HCL include the presention of results as a dendrogram, and that it does not require any prediction of the resulting clustering pattern. However, because HCL does not offer the option of pre-defined structure to the user, it has been alleged that biological interpretation of the HCL results is a complicated and laborious process (Okada et al., 2005).

In k-means clustering, the array elements are first randomly assigned into k clusters, a user-defined value (Tavazoie et al., 1999). The algorithm then calculates the distances within each cluster and between clusters. Then the array elements are shuffled between clusters in an iterative process, calculating the new distances after each move, until the array element arrives in a cluster where the distance is smallest. The process reiterates until no more movements can be made (Quackenbush, 2001). This algorithm is particularly suited for analyses where a partial structure is desired, as it allows for a preset number of clusters. Examples of k-means clustering applications include gene expression analyses to study human heart failure (Tan et al., 2002), stress-response in rats (Morgan et al., 2005), and the post-feeding processes in the malaria mosquito Anopheles gambiae (Dana et al., 2005).

Self-organising maps (SOMs) also allows the user to dictate the number of sections, in this case a simple geometric structure, such as a 2-dimensional rectangular grid. The nodes of the grid are 'trained' by an iterative process, adjusting their values so that they move closer towards their assigned genes, until the most effective separation is achieved (Quackenbush, 2001). The SOM algorithm has been incorporated into the program GENECLUSTER, (Tamayo et al., 1999). It has been used to analyse expression data in yeast (Tamayo et al., 1999) and *Clostridium acetobutylicum* (Alsaker and Papoutsakis, 2005). It has helped to identify gene expression patterns of the cellular response to HIV (Vahey et al., 2002), and to an oncogene, v-fos (Ordway et al., 2005). It has also been applied to studying the development and molecular signature of many cancer cells, including squamous cell cancer (Hummerich et al., 2005), Hepatitis C virus-associated hepatocellular carcinoma (Iizuka et al., 2005; Tsunedomi et al., 2005), promyelocytic leukaemia (Song et al., 2003), B-cell lymphomas (Wang et al., 2002b), embryonal carcinoma (Harris and Childs, 2002), as well as the molecular classification of tumour type (Covell et al., 2003).

The Institute for Genome Research (TIGR) have developed a suite of programs (TM4) for microarray analysis, which includes an application for data processing and normalization, Microarray Data Analysis System (MIDAS); also a program for the analysis and visualisation of microarray data, Multiexperiment Viewer (MeV). Both applications have incorporated a large number of algorithms, allowing users to process, normalise, cluster and perform statistical analysis for microarray data.

1.2.3.5 Confirmation of microarray results

Like many other scientific experiments, it is important that microarray studies include the use of additional methods to verify the results. Usually a small subset of the genes identified by microarrays are confirmed using follow-up methods, for example reverse transcription PCR (RT-PCR) (Corbeil et al., 2001; Ryo et al., 2000; Tang et al., 2005; van 't Wout et al., 2003), quantitative real-time RT-PCR (Carter et al., 2002; Leong et al., 2005), and northern blotting (Ahn et al., 2002; Chambers et al., 1999; Chang and Laimins, 2000; Paulose-Murphy et al., 2001; Ryo et al., 2000). The results from these methods are gnerally consistent with the microarray results, hence strengthening the reliability of the microarray findings. Methods such as RT-PCR may also differentiate expression patterns of splice variants (Menegazzi et al., 1999; Mirandola et al., 1998). Moreover, since changes in transcript abundance may not necessarily result in changes in protein levels, in some experiments western blotting (Guerra et al., 2004) and immunoassays for proteins (Tang et al., 2005) have been used to correlate transcriptional changes identified by microarrays with protein levels.

1.2.4 Applications of microarrays in host-pathogen interactions

During all stages of infection, viruses subvert the host cellular machineries for their own benefit, altering the expression of many cellular genes. Meanwhile host organisms have developed many ways to control the infection. At any given time, the expression patterns of both viral and cellular genes represent the complex interactions between host and pathogen. The use of microarray techniques in studying these host-pathogen interactions during infection has drawn particular interests (Cummings and Relman, 2000; Diehn and Relman, 2001; Jenner and Young, 2005; Kellam, 2001), and has provided many insights in host-pathogen studies.

1.2.4.1 Microarray studies of microbial gene expression

The availability of microbial genome sequences (section 1.2.1) have prompted the development of whole- (or almost whole) genome microarrays for many microorganisms, and the gene expression of Escherichia coli (Richmond et al., 1999), Mycobacterium tuberculosis (Wilson et al., 1999), Plasmodium falciparum (Hayward, 2000; Rathod et al., 2002), Streptococcus pneumoniae (de Saizieu et al., 2000), Salmonella typhimurium (Eckmann et al., 2000; Rosenberger et al., 2000), and Entamoeba histolytica (MacFarlane et al., 2005b) have been studied. The arrays used to study the physiological state of E. coli during heat shock discovered expression changes in more than 80 previously known heat-shock related-genes and 35 novel genes (Richmond et al., 1999). The temporal transcription programme during the metabolic shift from anaerobic fermentation to aerobic respiration in Saccharomyces cerevisiae has also been investigated using microarrays (DeRisi et al., 1997). Microarrays also revealed coordinated gene expression in *Plasmodium falciparum*, the malarial parasite (Ben Mamoun et al., 2001). Genes involved in protein synthesis were expressed during the early stages of infection, followed by metabolism genes, and genes involved in adhesion and invasion. Microarrays are also useful in studying drug actions. A study in M. tuberculosis discovered that four genes encoding fatty acid biosynthesis enzymes were co-induced with a known target of isoniazid in the presence of the drug. Therefore these genes may be potential targets for new anti-tuberculosis drugs (Wilson et al.,

1999). Oligonucleotide arrays have been used to detect drug-resistant mutants of M. *tuberculosis* (Troesch et al., 1999). Based on the observation that genes specifically induced during infection are often required for virulence (Cotter and Miller, 1998), probable virulence factors can be identified by comparing gene expression profiles of virulent and avirulent strains. For example, microarrays have found that virulent isolates of *Vibrio cholerae* from patient stool had upregulated expression of 44 genes compared to non-pathogenic laboratory strains (Merrell et al., 2002). Many of these genes were involved in cell motility and nutrient acquisition.

1.2.4.2 Microarray studies of viral gene expression

Microarray applications are particularly suited for analysing viruses with large genomes, for the same reasons as other microbes (section 1.2.4.1). Both spotted DNA microarrays and oligonucleotide microarrays have been put to use in studying viral infections. The transcriptomes of many herpesviruses, including herpes simplex virus (HSV) (Stingley et al., 2000), HCMV (Chambers et al., 1999), HHV-6 (Ohyashiki et al., 2005), Kaposi-sarcoma associated herpesvirus (KSHV) (Jenner et al., 2001; Paulose-Murphy et al., 2001) and murine gammaherpesvirus 68 (MHV-68) (Ahn et al., 2002), have been analysed by microarrays. These microarray studies revealed that herpesvirus genes were transcribed in a co-ordinated manner, with viral genes involved in particular functions co-expressed at the corresponding stages of infection. Most of these array studies have confirmed the kinetic classification of herpesvirus genes and suggested putative functions of unclassified viral genes. For example, Paulose-Murphy et al predicted that 4 KSHV ORFs (11, 31, 35, 49) may function as transactivators, one (ORF10) may be involved in viral DNA replication, and 12 ORFs (K7, 18, 23, 24, 27, 28, 30, 33, 42) were possible structural genes (Paulose-Murphy et al., 2001). In another study (Jenner et al., 2001), a region between K10.5 and K11 in KSHV was identified by microarrays, which may represent a novel transcribed ORF (K10.7) with homology to viral interferon regulatory factors (vIRFs). More details on temporal classification of herpesvirus using microarrays are given in section 5.1. Microrrays have also been produced for VZV (Kennedy et al., 2005) and vaccinia virus (Wenli et al., 2004), which are sensitive to viral transcripts, although these have not yet been used in expression profiling.

1.2.4.3 Microarray studies of host gene expression during viral infection

The expression patterns of cellular genes during infection by various viruses have been explored using microarrays (Table 1.4). Amongst these most followed the infection *in vitro*, except a few studies on *in vivo* (Bigger et al., 2001; Domachowske et al., 2002; Taylor et al., 2000). Data showed that viral infection had substantial influence on many cellular processes, not only on defence mechanisms such as DNA repair, cytokine signalling and stress response, but also on normal cellular functions like transcription regulation, mRNA processing, protein synthesis, protein degradation, cell cycle progression and mitochondrial functions (Jenner and Young, 2005). Very often hundreds of genes are significantly upregulated or downregulated during infection (Corbeil et al., 2001; Zhu et al., 1998), demonstrating the complexity of viral infection, also the power of microarray application in host-pathogen interaction analysis.

Microarray technology offers powerful comparative studies. For example, experiments have compared viruses capable of replication with their inactivated counterparts. These have helped to identify host responses that are dependent on viral replication (Browne et al., 2001; Geiss et al., 2001; Jones and Arvin, 2003; Mossman et al., 2001; Wang et al., 2005). In particular, it has been found that replicating HCMV induces fewer interferonresponsive genes and proinflammatory cytokines in human fibroblasts than their nonreplicating counterparts (either by UV-inactivation or by treatment with metabolic inhibitor), suggesting that viral gene products may block cellular antiviral responses (Browne et al., 2001). Alternatively, more changes in cellular gene expression were caused by replicating influenza virus than by inactivated virus (Geiss et al., 2001). Studies have also compared the cellular responses to related viruses that cause different clinical symptoms, offering insights into the pathogensis of the more virulent virus. For example, it was found that the severe acute respiratory syndrome-associated coronavirus (SARS-CoV) induces greater changes in the transcription of cellular genes associated with apoptosis, inflammation, stress response and procoagulation than human coronavirus 229E which causes the common cold (Tang et al., 2005). Differential gene expression has been observed between the oncogenic adenovirus type 12 and nononcogenic adenovirus type 5 (Vertegaal et al., 2000), also between patient samples

obtained from hepatitis B and C lesions (Honda et al., 2001). A study has also compared the gene expression patterns in different cell types to identify cell type-specific responses to VZV (Jones and Arvin, 2003).

Some have studied how host gene expression were affected by knocking out viral genes, such as HSV-1 UL54 (Stingley et al., 2000) and HCMV U28 (Hertel and Mocarski, 2004); and by expressing specific virus proteins such as HIV Tat (de la Fuente et al., 2002), HIV Nef (Shaheduzzaman et al., 2002; Simmons et al., 2001), KSHV LNA-1 (Renne et al., 2001), KSHV vGPCR (Polson et al., 2002), and IE86 of HCMV (Song and Stinski, 2002). Such studies have helped identify host responses to specific viral proteins, for example HCMV glycoprotein B been shown to trigger IFN-stimulated genes (Simmen et al., 2001).

In summary, microarrays are a powerful tool for the large-scale monitoring of changes in gene expression that have been extensively used in many applications, and have provided many useful insights in cell biology, disease, and host-pathogen interactions.

Table 1.4 Microarray studies of host gene expression in viral infections.

			Arrav	Cellular response category ^d								
Family	Virus species *	Cell type ^b	type°	No. probes	1	2	3	4	5	6	7	Reference
dsDNA												
Adenoviridae	AdV type-5 and type-12	HEL, HEK, HER cells	М	18376			+					Vertegaal et al 2000
Herpesviridae	EBV	B-lymphoma cell lines	м	4146	+		+/-		+/-	+		Carter et al 2002
	HCMV	HFF	0	12626	+	+	+/-		-	+		Browne et al 2001
	HCMV	HFF	S	8942	+							Simmen et al 2001
	HCMV	HFF	0	1650x4	+	+		+	-	+	+/-	Zhu et al 1998
	HCMV	HFF	S	29593	+			-		+	mitosis	
	HHV6	SupT1	М	1152		+/-						Mayne et al 2001
	HSV-1	HEL fibroblasts	м	588							arrest G2-M	Hobbs and DeLuca 1999
	HSV-1		М	588				+	-		arrest G2-M	Khodarev et al 1999
	HSV-1	HEL fibroblasts	S	19000	+			+				Mossman et al 2001
	HSV-1		S	57	+						+/-	Stingley et al 2000
	HSV-1		М	18378						÷	+/-	Tsavachidou et al 2001
	VZV	T-cells, fibroblasts, human skin grafts in mice	S	9180	+		+/-	+/-	+/-		+/-	Jones and Arvin 2003
	KSHV	DMVEC	S	4165	+			+			mitosis	Moses et al 2002
	KSHV	DMVEC	М	2350	+			+				Poole et al 2002
	Marek's disease virus	Chicken embryo fibroblasts	М	1126	+							Morgan et al 2001
Poxviridae	Vaccinia virus	HeLa	S	15360	-			-		-		Guerra et al 2003
	Vaccinia virus Ankara	HeLa	S	15360	+	+						Guerra et al 2004
Papovaviridae	HPV 31	NHK	S	7075	-							Chang and Laimins 2000
dsDNA-r												
Hepadnaviridae	Hepatitis B virus	Liver cells from patients	S	1080	+	+	+	+				Honda et al 2001
	Hepatitis B virus	HCCs from patients	S	23040							proliferation	Okabe et al 2001
dsRNA		······································							····			
Reoviridae	Reovirus serotypes 1 & 3	HEK293	0	12626							arrest G2-M	Poggioli et al 2002
	Rhesus rotavirus	Caco3	S	38432	+	+		+	+			Cuadras et al 2002
ssRNA-		<u> </u>										
Orthomyoxoviridae	Influenza virus	HeLa	S	4608		-		+/-	+/-	-		Geiss et al 2001
Paramyoxviridae	Measles virus	PBMCs	м	1176x3	+			+				Bolt et al 2002

			Array		Cellular response category ^d							
Family	Virus species *	Cell type ^b	type ^c	No. probes	1	2	3	4	5	6	7	Reference
	Pneumonia virus of mice	Lung tissues from mice	0	12000	+	+		+				Domachowske et al 2002
Paramyoxviridae	Respiratory syncytial virus	A549	М	268		+		+		+		Zhang et al 2001b
	Respiratory syncytial virus	Hep2	ο	12626	+			+	-			Tian et al 2002
	Respiratory syncytial virus	Hep2, A549	0	12626	+							Zhang et al 2001b
ssRNA+												
Flaviviridae	Hepatitis C virus		ο	7000	+							Bigger et al 2001
	Hepatitis C virus	Liver cells from patients	S	1080	-	-					+/-	Honda et al 2001
	Hepatitis C virus	HCC from patients	S	23040							Anti-cancer	Okabe et al 2001
Picornaviridae	Coxsackievirus b3	Heart tissues from mice	S	7000				+				Taylor et al 2000
	Echovirus 1		М	588				+			arrest G2-M	Pietiainen et al 2000
	Poliovirus	HeLa	S	10000				+				Johannes et al 1999
Coronaviridae	SARS-CoV	Vero E6	S	16000	+/-		+/-	+/-		+/-	+/-	Leong et al 2005
	SARS-CoV	Huh7	0	22000	+	+	+/-	+		+/-	+/-	Tang et al 2005
	Coronavirus 229E	Huh7	0	22000				+/-		-	+/-	Tang et al 2005
SsRNA-RT												
Retroviridae	HIV-1	human T-cell line CEM-CCRF	S	1506			+/-					Geiss et al 2000
	HIV-1		0	6800	+			+	-		arrest G2-M	Corbeil et al 2001
	HIV-1	human T-lymphoblastoid cell lines	S	4608	+/-	+/-	+/-	+/-	-	-	arrest G2-M	van't Wout et al 2003
	Human foamy virus	HEL	м	588							arrest G2-M	Wagner et al 2000
	Human T-cell leukaemia viru	S	М	588				+		+	progression	de La Fuente et al 2000

^a Virus species: Adv, adenovirus; EBV, Epstein-Barr Virus; HCMV, human cytomegalovirus; HHV-6; human herpesvirus type-6; HSV-1, herpes simplex virus 1; VZV, varicella-zoster virus; KSHV, Kaposi's sarcoma-associated herpesvirus; HPV, human papillomavirus; SARS-CoV, severe acute respiratory syndrome-associated coronavirus; HIV-1, human immunodeficiency virus-1

^b Cell type: HEL, Human embryonic lung fibroblasts; HEK, human embryonic kidney fibroblasts; HER, human embryonic retina fibroblasts; HFF, Human foreskin fibroblasts; SupT1, Human T-cell line; DMVEC, Dermal microvascular endothelial cells; NHK, Normal human keratinocytes; HeLa, ; PBMCs, peripheral blood monocytic cells; Huh7, Human hepatoma cell line; CEM-CCRF, Human T-cell line; Hep2, ; A549, Human lower airway epithelial cells; HCC, hepatocellular carcinoma; Caco3, Human intestine cell line, Vero E6, African green monkey kidney cells.

^c Array type: M, membrane array; S, spotted DNA microarray; O, Oligonucleotide chip (synthesised *in situ*).

^d Cellular response category: 1, Interferon and pro-inflammatory response; 2, Cytokines; 3, Signal transduction; 4, Stress response (apoptosis, heat shock, oxidative stress); 5, Protein synthesis; 6, Transcription factors; 7, Cell cycle. +, expression mostly upregulated; -, expression mostly downregulated; +/-, expression altered

1.3 Aim and Scope of thesis

This PhD thesis aims to investigate the complex, intertwined processes during lytic infection of the human T cell by human herpesvirus type-7 (HHV-7), a highly prevalent and relatively non-pathogenic virus. I aim to study the expression of viral and host genes simultaneously, in order to identify the cellular responses to HHV-7 at different stages of lytic infection. I also intend to examine the lytic replication cycle of HHV-7 and compare this to other human betaherpesviruses. To achieve this, the following steps were undertaken:

- A custom-made host-pathogen microarray was created in-house, and its proficiency for use in measuring HHV-7 and host gene transcription was tested (Chapter 3).
- An *in vitro* system for cell-free HHV-7 infection of human T-cells with a high multiplicity of infection was developed to allow the study of the HHV-7 gene expression programme (Chapter 4).
- 3) Finally, the custom made microarrays were applied to the analysis of HHV-7 and human cellular gene expression in parallel. Particularly, the temporal regulation of HHV-7 viral gene expression was examined by expression profiling with and without the use of metabolic inhibitors on *de novo* protein synthesis and viral DNA replication (Chapter 5).

So far no extensive gene expression studies on HHV-7 have been performed, as such studies are limited by the virus's low infectivity. This is the first description of a host-pathogen microarray analysis for HHV-7.

CHAPTER 2

MATERIALS & METHODS

2.1 Buffers and Solutions

Solution	Ingredients
Deoxynucleotide triphosphate mix (dNTPs)	100 mM of deoxyadenosine triphosphate (dATP), deoxythymidine triphosphate (dTTP), deoxyguanosine triphosphate (dGTP), deoxycytidine triphosphate (dCTP)
Luria-Bertani (LB) agar	1% (w/v) bacto tryptone, 0.5% (w/v) bacto yeast, 0.5% (w/v) sodium chloride, pH 7.0 with 1.5% (w/v) bacto-agar
Luria-Bertani (LB) broth	1% (w/v) bacto tryptone, 0.5% (w/v) bacto yeast, 0.5% (w/v) sodium chloride, pH 7.0
10x PCR buffer (20 mM magnesium chloride)	100 mM Tris, pH 8.3, 500 mM potassium chloride, 20 mM magnesium chloride
10x PCR buffer (10 mM magnesium chloride)	100 mM Tris, pH 8.3, 500 mM potassium chloride, 10 mM magnesium chloride
Phosphate-buffered saline (PBS)	137 mM sodium chloride, 2 mM potassium chloride, 10 mM sodium hydrogen phosphate (dibasic), 2 mM potassium hydrogen phosphate (dibasic), pH 7.4
20x Saline sodium citrate (SSC)	3 M sodium chloride, 0.3 M sodium citrate, pH 7.0
Saline-sodium phosphate- EDTA buffer SSPE	2.98 M sodium chloride, 0.02 M EDTA, 0.2 M phosphate buffer, pH 7.4
Tris-acetate-EDTA (TAE)	40 mM Tris pH 7.8, 20 mM sodium acetate, 1 mM ethylenediaminetetraacetic acid (EDTA)
Tris-EDTA (TE)	10 mM Tris pH 7.4, 1 mM EDTA

Table 2.1 Constituents of buffers and solutions

2.2 Mammalian cell culture

2.2.1 Cell lines

Table 2.2 Cell lines

Cell line	Description
CEM-SS	Human T4-lymphblastoid cell line, suspension
HeLa	Cervical epithelium, adherent
Huh7	Hepatoma, adherent
MRC5	Embryonic fibroblasts, adherent
Ramos	B-cell, suspension
SupT1	Non-Hodgkin's T-cell Lymphoma, CD4+, suspension
U937	Monocytes, suspension

2.2.2 Cell growth media

Suspension cell lines were grown in Roswell Park Memorial Institute (RPMI)-1640 medium (Invitrogen, UK) supplemented with 10% foetal bovine serum (FBS, Biowest). When infected with virus, cultures were maintained in RPMI with reduced (5%) FCS to slow down cell growth. Adherent cell lines were cultured in Dulbeccos Modified Eagle

Media (DMEM) medium (Invitrogen, UK) with 5% FCS. All culture media were supplemented with 100 units/ml penicillin and 100 μ g/ml streptomycin (P/S, Invitrogen, UK). All mammalian cells were cultured in tissue culture flasks with filtered caps or multi-well plates (Helena Biosciences, UK) in humidified incubators, 37 °C, 5% CO₂. Centrifugations of cells were carried out at 325g, 5 min, 20 °C, unless otherwise stated. All cells were tested negative for mycoplasma.

2.2.3 Thawing

Cells taken from liquid nitrogen were thawed immediately at 37 °C and added to 2×5 ml pre-warmed medium. Within 24 h cells were washed and medium replaced. Suspension cells were pelleted at 325g, 5 min and washed in 10 ml medium, followed by centrifugation for 5 min and resuspension in pre-warmed medium at 2×10^5 cells/ml. For adherent cells medium was removed and cells were washed with 10 ml medium per flask, and fresh medium was added to 1×10^5 cells/ml.

2.2.4 Passaging cells

Cell counts were obtained using a haemocytometer. Suspension cells were split 1:4 to 1:8, every 3-6 days, depending on cell line and cell density. To split adherent cell lines, cells were first detached from culture surface using 1 ml trypsin/EDTA (Invitrogen, UK) per 25 cm² culture area, and then split 1:2 to 1:6.

2.2.5 Freezing cells

Cells were pelleted by centrifugation and resuspended in corresponding freezing medium (FCS with 10% dimethyl sulphoxide (DMSO, Sigma, UK) at a final concentration of $1-2\times10^6$ cells/ml. Cell stocks were aliquoted into cryovials (Nunc, USA) and frozen gently to -80 °C overnight in an isopropanol-containing cryo-container (Nalgene, USA) before being transferred to liquid nitrogen for storage at -140 °C.

2.3 Culturing HHV-7

2.3.1 Virus strain

Human herpesvirus type 7 (HHV-7) used in this thesis was strain DC, provided by Dr Duncan Clark, Royal Free & University College Medical School, London.

2.3.2 Virus infection

For infection of SupT1 cell line by cell-free human herpesvirus type 7 (HHV-7), 1 ml of virus stock was allowed to adsorb to $1 \times 10^5 - 5 \times 10^6$ cells for 1-2 h at 37 °C. For spinoculation, the virus-cell mixture was centrifuged at 1000 g for 1 h at 20 °C. After adsorbtion (with or without centrifugation), the virus-cell mixture was seeded into tissue culture flasks with pre-warmed medium and incubated at 37 °C, 5% CO₂.

2.3.3 Virus culture and maintenance

Virus culture was maintained by co-culturing as follows. HHV-7 infected cells were monitored for the display of cytopathic effect (CPE), characterised by enlarged, multinucleated cells, and elongated cells attached to bottom. Cultures with greater than 50% CPE were co-cultured with uninfected cells at the ratio of 1:1 to 1:5 (infected: uninfected) by cell count. Medium was added to achieve cell density of $2 \times 10^5 - 1 \times 10^6$ cells/ ml.

2.3.4 Stock preparation

HHV-7 virus stocks were prepared as follows. HHV-7 infected cells were centrifuged and resuspended in 1/6 - 1/100 original volume of FBS. Cells were lysed by freeze-thawing 6 times to release cell-associated virus particles. Cell lysates were centrifuged

again to separate cell debris. Supernatant containing virus particles was retrieved, aliquoted into 1ml stocks, and stored at -140 °C (liquid nitrogen).

2.3.5 Virus quantification

2.3.5.1 Observation of cytopathic effect (CPE)

HHV-7 infected SupT1 cells were monitored for cytopathic effect (CPE) by visual observation under light microscopy ($200 \times$ magnification). Percentage CPE was estimated from the proportion of cells attached to bottom of tissue culture flask, enlarged ($2 \times$ normal diameter or above), elongated or become irregularly shaped (detailed in section 4.2.2).

2.3.5.2 Immunofluorescence assay (IFA)

HHV-7 infected cells were harvested and centrifuged, washed in 2× original volume of Phosphate buffered saline (PBS) and resuspended in PBS to a final concentration of 1×10^{6} cells/ml. 2×10^{4} cells were then spotted onto each well of polytetrafluoroethylene (PTFE)-coated multispot slides (C.A. Hendley Ltd), air-dried for 40 min, and fixed in cold acetone (-20 °C) for 10 min. Fixed cells were then assayed for HHV-7 infection by incubating with mouse-derived anti-HHV-7 monoclonal antibody that target an early HHV-7 antigen (20 µl per well at 2.5 ng/µl in PBS; H2034-18, US Biological, Massachusetts) for 40 min in humidified chamber at 37 °C. The slide was then washed three times in PBS, followed by incubation with FITC-conjugated anti-mouse IgG antibody (20 µl per well at 5 ng/µl in PBS; F2883, Sigma) for 40 min, 37 °C. Slides were washed 3 times in PBS and air-dried. Slides were then counterstained with Vectashield mounting medium with DAPI (Vector Laboratories, USA) and examined by fluorescent microscopy using the Axioplan2 imaging Pol (Zeiss, Germany), to detect the presence of HHV-7 antigen (green fluorescent). The number of green fluorescent (by FITC) cells and the total number of cells (by DAPI counter-stain) were counted in four fields per well. The mean percentage of infected cells was calculated as:

> % Infected cells = <u>HHV-7 antigen-positive cells (FITC)</u> Total cells (DAPI counter stain)

2.3.5.3 Infectious units count (IUC)

SupT1 cells were plated in a 24-well plate at 5×10^5 cells per well and pelleted by centrifugation. 100 µl of serially diluted virus (neat, $\frac{1}{2}$, $\frac{1}{4}$) were added per well in triplicate per dilution and incubated at 37 °C for 2 h to allow virus attachment. 500 µl of medium (RPMI, 5% FBS, P/S) was then added to each well and the cells were incubated at 37 °C for 3-4 days. Cells were then observed for CPE (section 2.3.5.1) or harvested for IFA (section 2.3.5.2). Infectious units count per ml virus (by CPE or by IFA) was calculated as:

% Infected cells ×
$$5 \times 10^5$$
 cells × $\frac{1000 \ \mu l}{100 \ \mu l}$

2.3.5.4 TCID₅₀ determination

SupT1 cells were plated in 24-well plates at $1 \times 10^5 - 5 \times 10^5$ cells per well and pelleted by centrifugation. 100 µl of serially diluted virus (half-log dilution $10^0 - 10^{-5.5}$) were added per well in quadruplicate per dilution and incubated at 37 °C for 1-2 h to allow virus attachment. 100 µl of RPMI-10 medium was then added to each well and the cells were incubated at 37 °C for 3-4 days. After incubation cells were observed for CPE (section 2.3.5.1) or harvested for immunofluorescence detection of infection (section 2.3.5.2). Wells containing unambiguously infected cells were scored as positive and TCID₅₀ was calculated as follows:

 $-\log TCID_{50} = -\log L - \log d(S - 0.5)$

Where: L = highest virus dilution to give 100% positive

d = dilution factor (for half-log dilution, d = 0.5)

S = sum of the ratios (positive wells/total wells) for every dilution from L

2.3.6 Flow cytometry

Cells were prepared for flow cytometry analysis as follows. 5×10^5 cells were washed by adding 2× volume PBS and spinning at 1300 g, 5 min, 4 °C. Cells were fixed on ice for 10 min in 4% paraformaldehyde in PBS. Cells were washed twice in PBS and permeabilised in FACS permeabilisation solution (0.1% Triton X-100, 0.01% sodium azide, 2% FBS in PBS). Permeabilised cells were incubated with anti-HHV-7 monoclonal antibody (2.5 ng/µl; H2034-18, US Biological, Massachusetts) in 50 µl FACS perm + 10% FBS for 1 – 2 h on ice. Unbound antibody was removed by washing three times in 1 ml FACS permeabilisation solution. Cells were incubated with the secondary antibody, FITC-conjugated anti-mouse IgG (5 ng/µl; F2883, Sigma) in 50 µl FACS perm + 10% FBS for 1 – 2 h on ice in the dark. Finally cells were washed three times in 1 ml FACS permeabilisation solution and once in 1 ml FACS wash solution (0.01% sodium azide, 2% FBS in PBS). Ten thousand events of the prepared cells were collected and analysed using a FACScan with Cellquest software (Becton Dickinson, UK).

2.3.7 Time course infection of SupT1 cells by HHV-7

For each time point, 1×10^5 cells of SupT1 (per well in 24-well plate) were inoculated with 1 ml HHV-7 virus stock (MOI = 2.5) or mock infected with uninfected cell lysate for 1 h, with spinoculation (1000 g) at 20 °C. After 1 h the virus inoculum was carefully removed without disturbing the cells. 1 ml of RPMI medium (P/S, 5% FBS) was added to the cells and the plates incubated at 37 °C. At each time point (8, 12, 16, 20, 24, 38, 48, 72 hours post infection) the cells (HHV-7-infected and mock-infected) were centrifuged to remove supernatant. The cells were resuspended in 350 µl RLT lysis buffer (Qiagen, UK) and stored at -80 °C for later RNA extraction (section 2.7.2). In earlier attempts when 5×10^6 cells were used per time point, cells were lysed in 1ml TRIzol reagent (Invitrogen, UK) and stored at -80 °C for later RNA extraction (section 2.7.1).

2.3.8 Metabolic inhibitor assay for temporal gene expression mapping

For metabolic inhibitor assay, SupT1 cells were first treated with 200 μ g/ml cycloheximide (CHX, Sigma, UK), 500 μ g/ml phosphonoacetic acid (PAA, Sigma, UK), or mock (medium) for 1 h at 37 °C prior to infection by HHV-7. Cells were then centrifuged to remove supernatant, and inoculated with HHV-7 stock as in section 2.3.7. Constant concentration of inhibitors was maintained throughout the infection and time course. Cells were harvested at 8 and 16 hpi (CHX-treated and untreated); at 24 and 48 hpi (PAA-treated and untreated) in 350 μ l RLT lysis buffer (Qiagen, UK) and stored at - 80 °C for later RNA extraction (section 2.7.2).
2.4 Producing HHV-7 array probes

2.4.1 Array probe primers

A list of primers used in this thesis is provided in Table 3.1.

2.4.2 HHV-7 DNA preparation

To obtain template DNA for amplification of HHV-7 probes, DNA was isolated from HHV-7 infected SupT1 cells using the Wizard Genomic DNA Purification Kit (Promega, UK). Briefly, 2×10⁶ SupT1 cells (2 ml, infected with HHV-7 with CPE> 50%) were harvested by centrifugation and washed with 1 ml PBS. Cells were then centrifuged at 13,000 g to remove the supernatant, and the cell pellet was resuspended in 200 µl PBS. Cells were lysed with 600µl of nuclei lysis solution. RNA was digested using 3µl RNase solution for 20 min at 37°C, and protein was removed by incubating with 200µl protein precipitation solution on ice for 5 min. The precipitated protein was removed by centrifugation at 16,000 g, 20°C for 4 min. The DNA in the supernatant was recovered into a clean tube which contained 600µl isopropanol (20°C) and was centrifuged at 16,000 g, 20°C for 1 min to precipitate the DNA. Supernatant was removed and the DNA pellet washed with 600µl 70% ethanol (20°C), centrifuged again to remove the ethanol and air-dried for 10 min. The DNA pellet was resuspended in 100µl of DNA rehydration solution (10mM Tris-HCL (pH 7.4), 1mM EDTA (pH8.0)) at 4°C overnight. The concentration of the purified DNA was calculated from the UV absorbance at 260 nm using a UV spectrophotometer (Camlab, UK). An absorbance of 1 cm⁻¹ was taken to be equivalent to 50 µg/ml DNA. DNA was stored at -20°C until use in PCR (section 2.4.3).

2.4.3 Initial amplification of viral DNA

Polymerase chain reaction (PCR) was performed to generate array probes of approximately 300 bp in length for each open reading frame (ORF) of HHV-7 genome.

Probes complimentary to the 5'-end of each ORF (Figure 3.2) were designed using the Primer3 program, so that all probes have similar length and annealing temperature. Each PCR reaction contained 500 pmol of each primer (Table 2.1, Oswel Research Products Ltd., UK), 1.5 μ g of HHV-7 DNA (section 2.4.2), 20 μ mol dNTP (Promega), and 1.25 units Tag DNA polymerase (Promega), in 50 μ l reaction buffer (10 mM Tris (pH8.3), 50 mM KCl, 2 mM MgCl₂ in UV-sterilised water). PCR reactions were carried out in Peltier thermal cycler (MJ Research, USA) with the following programme: 95 °C for 3 min; 30 cycles of 94 °C for 30 sec, 54 °C for 30 sec, 72 °C for 45 s; then a final extension step at 72 °C for 7 min. The presence and size of PCR products were checked by agarose gel electrophoresis (section 2.4.4) and if satisfactory, were used in ligation (section 2.4.5.2) directly.

2.4.4 Agarose electrophoresis of PCR product

PCR amplified probes were analysed by gel electrophoresis using 1.5 % agarose gel Electrophoresis analysis of nuclei acids were performed in 1-2% agarose gel in 1x Trisacetate-ethylenediaminetetraacetic acid (EDTA) (TAE), containing 0.2 μ g/ml ethidium bromide (Sigma, UK), under electrical potential difference of 60-120V (100-200 mA). Separated nucleic acids were visualised under UV illumination.

2.4.5 Cloning array probes

2.4.5.1 Competent cell preparation

Escherichia coli (XL1-Blue) (Stratagene, UK) was grown to saturation at 37 °C in a shaking incubator overnight in 5 ml Luria-Bertani (LB) broth containing tetracyclin (10 μ g/ml). This was used to inoculate LB-broth (1:100) and shaken at 37°C until the culture reached mid-log phase (OD₅₅₀ ~0.5, as determined by spectroscopy). The culture was chilled on ice for 10 min, followed by centrifugation at 3000 g, 5 min, 4 °C. The bacteria were resuspended in half original volume of CaCl₂ (50 mM CaCl2, 10 mM Tris pH8.0, 4 °C) and left on ice 15 min. Bacteria were then centrifuged again and resuspended in 1/15th original volume of CaCl₂. Competent bacteria were either used

immediately or stored frozen in liquid nitrogen in 200 μ l aliquots (100 mM calcium chloride containing 15% (v/v) glycerol).

2.4.5.2 Ligation of DNA into pGEM-T easy vector

The HHV-7 PCR products (2.4.3) were ligated directly into 50 ng of pGEM-T Easy vector (Promega, UK), using 3 units of T4 DNA ligase (Promega, UK), 5 μ l 2×ligase buffer (20mM Tris-HCl (pH 7.8), 10mM MgCl₂, 10mM DTT, 1mM ATP) in 10 μ l at 4°C for 16 h.

2.4.5.3 Transformation

The ligation reaction was incubated with 200 μ l of competent XL-1 Blue *E.coli* (section 2.4.5.1) on ice for 30 min. Uptake of DNA was induced by heat shock at 42 °C, 45 sec, after which cells were returned to ice immediately and cooled for 2 min. Transformed cells were then allowed to equilibrate in 1 ml LB without antibiotics at 37 °C, 30 min. Cells were then pelleted at 3000 g, 10 min, 20 °C, and resuspended cells in residual LB (approx. 200 μ l) before spreading onto LB-agar plates containing 100 μ g/ml ampicillin (Sigma, UK), 0.5 mM isopropylthio- β -D-galactoside (IPTG, Promega, UK), 80 μ g/ml 5-bromo-4-chloro-3-indolyl- β -D-galactoside (X-gal, Promega, UK). The plates were incubated at 37 °C for 16 h to select for plasmid-containing, ampicillin-resistant bacteria, and for the insert-containing white colonies by X-gal/IPTG screening.

2.4.6 Identification of array probes

2.4.6.1 Growth of bacterial clones

To confirm the viral DNA inserts, 5 white colonies (indicate presence of transformed PCR product) were picked from each plate (section 2.4.5.3) and screened for the presence of viral DNA insert. Each colony picked was first dipped into PCR reaction mixture (section 2.4.6.1) and then into 150 μ l LB (100 μ g/ml ampicillin) on 96-well

plate and allowed to grow overnight at 37 °C. Next day the clones that were confirmed positive (contain viral DNA inserts) by PCR were selected for DNA extraction (section 2.4.6.2) and stored as archive in 20% glycerol v/v in 96-well plate at -80 °C. A maximum of two aliquots per clone were stored.

2.4.6.2 Screening for viral insert-containing clones by PCR

Polymerase chain reaction was performed to confirm the presence of viral gene inserts in XL1-Blue transformants using universal pGEM-T forward and reverse primers (Table 2.1). The ingredient of each PCR reaction was the same as in section 2.4.3, except for primer (150 pmol of each pGEM-T primer, Thermal BioSciences GmbH, Germany) and template (1 white colony of transformed XL1-Blue). PCR reaction was carried out as follows: 95 °C for 5 min; 30 cycles of 94 °C for 30 sec, 55 °C for 30 sec, 72 °C for 1 min; then 72 °C for 7 min. Products were checked for the presence of viral inserts by agarose gel electrophoresis (section 2.4.4).

2.4.6.3 Purifying plasmid DNA from positive clones

Plasmid DNA was isolated from the viral insert-containing *E.coli* (XL1-Blue) clones using the QIAprep Spin Miniprep kit (Qiagen, UK), a modified alkaline lysis method (Birnboim and Doly, 1979). 1.5 ml of bacterial culture (section 2.4.6.1) was pelleted at 14,000 g for 10 min and the bacteria resuspended in 250 µl resuspension buffer P1 (50 mM Tris-Cl (pH 8.0), 10 mM EDTA, 100 µl RNase A). Bacteria were lysed with the addition of 250 µl lysis buffer P2 (200 mM NaOH, 1% SDS (w/v)), and the solution neutralised with 350 µl neutralisation buffer N3 (3.0 M potassium acetate (pH 5.5). Genomic DNA was pelleted by centrifugation at 14,000 g for 10 min and the supernatant applied to a QIAprep column. The solution was passed through the column by centrifugation at 14,000 g, 1 min. The column was washed once with 500 µl buffer PB, twice with 750 µl buffer PE, and plasmid DNA eluted with 50 µl distilled water. The concentration of the purified DNA was calculated from the UV absorbance at 260 nm using a UV spectrophotometer (Camlab, UK). An absorbance of 1 cm⁻¹ was taken to be equivalent to 50 µg/ml DNA.

2.4.6.4 Verifying sequence of viral insert

The identity of the clones were verified by DNA sequencing using an automated application of the chain-termination method (Sanger et al., 1977). 1 μ l of plasmid DNA was added to 4 μ l of Beckman QuickStart mix (containing dATP, dCTP, dTTP, dITP, ddUTP, ddGTP, ddCTP, ddATP and thermosequenase DNA polymerase in Tris buffer, pH 8.9) and 5 pmol Sp6 primer (5'-TATTTAGGTGACACTATAG-3'). The solution was placed in a thermal cycler and subjected to the following conditions: 94°C for 3 min; then 30 cycles of 96°C for 20 sec, 50°C for 20 sec, 60°C for 4 min. This incorporates fluorescent di-deoxynucleotides into daughter DNA strands.

DNA was precipitated from the sequencing reaction by adding 2 μ l sodium acetate (1.5 M), 2 μ l EDTA (50 mM, Sigma, UK) and 1 μ l glycogen (20 mg/ml, Beckman Coulter, UK) in a final volume of 20 μ l, vortexing, and then adding 60 μ l 95% (v/v) ethanol (-20°C) and leaving on ice for 10 min. DNA was pelleted by centrifugation at 14,000 g for 15 min at 4°C. The supernatant was removed, the pellet washed with 200 μ l 70% ethanol (-20°C), and the tube left open at room temperature until dry. DNA was resuspended in 40 μ l de-ionised formamide (JT Baker, USA) for 15 min and transferred to a 96-well plate. The sequences were determined using an automated capillary DNA sequencer (Beckman Coulter, UK) as per the manufacturer's instructions.

2.5 Production of human gene probes

2.5.1 PCR optimisation

Unless otherwise stated, all liquid handling processes in the production of human gene probes were performed by Biomek 2000 Liquid Handler (Beckman Coulter) in 96-well and 384-well format plastic wares (Beckman Coulter, UK; Alpha Laboratories, UK; Millipore, UK). 12 DNA polymerases from 8 suppliers (Table 2.4) were tested for their suitability in PCR amplification of array probes. Each 50 μ l PCR reaction consisted of 40 pmol of each HHV-7 U76 primer (Thermal BioSciences GmbH, Germany), 1 μ l template DNA (serial dilution from 10 ng to 1 fg), 1.25 units DNA polymerase enzyme, 40 μ mol dNTP (Promega), in 5 μ l 10× reaction buffer containing 1.5 mM MgCl₂. Two sets of reaction mixes were prepared for all enzymes, one of which proceeded to PCR reaction immediately after preparation and the second set was left at room temperature for 3 h before PCR was performed. PCR was performed in a Primus Multiblock thermal cycler (MWG, UK) as follows: 94 °C for x min (Table 2.4); 40 cycles of 94 °C for 30 sec, 55°C for 30 sec, 72 °C for 30 sec; then 72 °C for 7 min. Products were checked for size and integrity by agarose gel electrophoresis (section 2.4.4).

TADIE 2.3 CONDITIONS IOI DIAA POLYMETASES III DALCH LE	Tab	le	2.3	Conditions	for DNA	polymer	ases ir	n batch te	s 1
--	-----	----	-----	------------	---------	---------	---------	------------	------------

			First heat	ing step
Code	Enzyme Name	Supplier	Temperature °C	Duration (min)
A	Taq	Qiagen	94	3
В	HotStar	Qiagen	94	15
С	Taq	Novagen	n/a	n/a
D	Hot Start	Novagen	95	10
E	Taq	Roche	94	2
F	FastStart	Roche	95	2
G	Taq	Sigma	94	1
н	Taq	Bioline	n/a	n/a
1	Immolase	Bioline	95	7
J	Taq	Promega	94	2
Κ	DyNAzyme	Finnzymes	94	2
L	YieldAce Hotstart	Stratagene	94	2

2.5.2 Preparing lymphochip clones for human array probes

Human gene probes were generated from a lymphochip gene set of ~2100 human clones (MRC gene service, Appendix B). The clone sets were in bacterial glycerol stocks (LB, 8% glycerol, 50 μ g/ μ l ampicillin) arrayed in 96-well formats. 5 μ l of each clone were inoculated into 150 μ l 2×LB (50 mg/ml ampicillin) and allowed to grow for 4-8 h at 37 °C with shaking. 5 μ l of this initial culture was used to inoculate 800 μ l LB (50 mg/ml ampicillin) and grown in the same way for 12-16 h. Bacterial clones were diluted 1:5 with distilled water and heated at 98 °C for 10 min and cooled on ice for 10 min to

break up the cell wall and release plasmid DNA. Cell debris was removed by centrifugation at 1500 g, 3 min to obtain DNA-containing cell-free supernatant for PCR (section 2.6.2).

2.6 Production of custom-made microarrays

2.6.1 Array PCR to amplify viral and human probes

Human array probes for in-house host-pathogen microarrays were produced by PCR amplification using pT7T3 primers (Table 2.3). Each 100 µl PCR reaction consisted of 40 pmol of each primer (Thermal BioSciences GmbH, Germany), 10 µl plasmid DNA-containing supernatant, 40 µmol dNTP (Promega, UK), and 2.5 units Immolase DNA polymerase (Bioline, UK), 10 µl 10× reaction buffer (160 mM (NH₄)₂SO₄, 1 M Tris-HVL (pH 8.3), 0.1% Tween-20). Viral probes were amplified in a similar manner to the human probes, using 40 pmol of each pGEM-T primer, 10 ng purified DNA. PCR was performed in a Primus Multiblock thermal cycler (MWG, UK) as follows: 95 °C for 7 min; 40 cycles of 94 °C for 30 sec, 55°C for 30 sec, 72 °C for 60 sec (30 sec for viral probes); then 72 °C for 7 min. Products were checked for size and integrity by agarose gel electrophoresis (section 2.4.4). Single band products were selected and redistributed onto fresh 96-well plates using Biomek HitPicking Wizard programme (Beckman Coulter, UK).

2.6.2 Human and viral gene probes purification

PCR products for both human and viral genes were transferred into Montage PCR₃₈₄ purification plates (Millipore, UK) for purification. Unincorporated nucleotides were removed by size selective filtration under 10 inches Hg vacuum for 15 min. Purified DNA was resuspended in 20 μ l spotting buffer, containing 3× saline sodium citrate buffer (SSC, Sigma, UK) and 1.5 M Betaine (Sigma, UK) and transferred into 384-well arraying plates (Genetix, UK). The process was automated using Biomek 2000 Liquid Handler (Beckman Coulter, UK).

2.6.3 Microarray spotting and processing

The purified DNA probes in spotting buffer were deposited onto aminopropyl silanecoated Nexterion Slide A (Schott, UK) using the Qarray² arrayer (Genetix, UK) with a 12-pin print head of Genetix aQu 75um split pins (Genetix, UK). The probes were spotted in triplicate, in a format of 36 sub-grids, with 196 spots (14 columns by 14 rows) per grid. Array spotting was performed in collaboration with Edward Tsao, Royal Free & University College Medical School, London. Spotted arrays were airdried for 12+ hours, followed by heating at 80 °C in dry oven for 3 h to immobilise the DNA. Unbound surfaces on the arrays were inactivated by immersing in 60 ml blocking solution: 5× SSC, 0.5% (w/v) bovine serum albumin (BSA, Sigma, UK), 0.5% (w/v) non-fat powdered milk (Tesco, UK) at 50 °C for 45 min, followed by washing 2 times in deionised water. Then the arrays were incubated in deionised water at 95 °C for 2 min to denature the DNA. Denatured arrays were washed and blocked again (with fresh solution) at 50 °C for 30 min. Finally the arrays were washed 2 times in deionised water, immersed in isopropanol (Sigma, UK) for 30 seconds and immediately dried by spinning at 200 g, 2 min. Processed arrays were stored in a dry environment for use within 24 hours.

2.7 Measuring transcript abundance using custommade microarrays

2.7.1 RNA extraction from TRIzol

2.7.1.1 Total RNA extraction

Each ml of frozen TRIzol lysates (section 2.3.7) were thawed at 37° C and centrifuged at 12,000 g, 10 min, 4 °C to remove insoluble material. DNA and proteins were removed from the supernanant by two rounds of chloroform purification, by adding 0.2 ml (first round) and 0.5 ml (second round) of chloroform, and the solution shaken by hand for 15

s. After incubation at room temperature for 2-3 min, the aqueous layer was separated from the organic layer by centrifuging at 12,000 g, 15 min, 4 °C. After the two rounds of chloroform purification, RNA in the aqueous phase was precipitated by incubating with 0.5 ml isopropanol for 10 min at room temperature. Precipitated RNA was pelleted by centrifugation at 12,000 g, 15 min, 4 °C and washed with 1 ml 75% ethanol. Ethanol was removed after centrifugation at 7500 g, 5 min, 4 °C. The RNA pellet was air-dried and resuspended in 50 - 100 μ l RNase-free water (BDH, UK). RNA concentration was estimated by UV absorbance at 260 nm, using the ND-1000 Spectrophotometer (Nanodrop Technologies, USA).

2.7.1.2 DNase treatment of total RNA

To ensure only pure RNA was used in the subsequent analysis, total RNA was treated with 5 units of DNase (Promega, UK) per 50 μ g RNA at 37 °C for 1 h to remove any contaminating DNA. DNase treated RNA was purified two rounds using an equal volume of 25:24:1 phenol:chloroform:isoamylalcohol (Sigma, UK), followed by one round of chloroform extraction (as in section 2.7.1.1), with vigorous shaking and centrifugation at 16,000 g, 10 min at room temperature to separate the RNA from the reaction components. Purified RNA was then precipitated for 1-2 h at -20 °C by adding 1/5th volume ammonium acetate (8M) and 2.5 volume 95 % ethanol (-20 °C), followed by centrifugation at 16,000 g, 30 min, 4 °C. The RNA pellet was washed in 200 μ l ethanol (80 %, -20 °C), centrifuged at 16,000 g, 10 min, 4 °C, and air dried for 5 min before resuspension in 100 μ l Rnase-free water. Concentration and quality of the DNase treated RNA was accessed using Agilent RNA 6000 Nano assay with the Agilent 2100 bioanalyser (Agilent Technologies, UK). If the RNA was not used immediately it was stored at -80 °C.

2.7.1.3 Messenger RNA purification

Messenger RNA (mRNA) was purified from total RNA using Oligotex (Qiagen, UK), which separates nucleic acids containing poly-dA sequences through hybridisation to oligo-dT bound-microscopic polystyrene-latex particles. Before purification, the Oligotex suspension was warmed to 37 °C, vortexed and then returned to room

temperature and buffer OEB heated to 70 °C. Total RNA was made up to 250 μ l and mixed with 250 μ l buffer OBB and 15 μ l Oligotex suspension. This was incubated at 70°C for 3 min to denature the RNA and then annealed at room temperature for 20 min. Subsequently, the Oligotex was pelleted by centrifugation at 14,000 g for 2 min and the supernatant discarded. The pellet was resuspended in 400 μ l buffer OW2, mixed and applied to a Qiagen spin column. This was centrifuged at 14,000 g for 1 min, the Oligotex resuspended in an additional 400 μ l buffer OW2 and centrifuged again. The bound mRNA was eluted by resuspending the Oligotex in 200 μ l of buffer OEB (70 °C) and centrifuging through the column at 14,000 g for 1 min. This was repeated to give a final volume of 400 μ l. The mRNA was concentrated to a volume of 10 – 20 μ l by centrifugation through a Microcon Centrifugal Filter Device YM-30 (Millipore Corporation, UK) at 14,000 g for 8 min. The mRNA was recovered by inverting the column and centrifuging at 1000 g for 3 min. Concentration and quality of the mRNA was accessed using Agilent RNA 6000 Nano assay with the Agilent 2100 bioanalyser (Agilent Technologies, UK). Purified mRNA was stored at -80 °C.

2.7.2 RNA extraction from RLT lysate

2.7.2.1 Total RNA extraction

For time course infection samples of a low cell number (1×10^5) , total RNA was extracted using the RNeasy mini kit (Qiagen, UK). SupT1 cells were harvested by centrifugation, washed in 2 volumes of PBS, and centrifuged again to remove the PBS. 1×10^5 cells were lysed in 350 µl RLT buffer (containing 0.1% (v/v) β-mercaptoethanol) and homogenised by vortexing. 350 µl 70% ethanol was added to the lysate and the mixture transferred onto spin column. The liquid was pulled through by centrifuging at 8000 g, 15 sec, room temperature. The column was washed once with 350 µl RW1 buffer and centrifuged as above, followed by digestion of DNA using 10 µl DNase, 70 µl RDD buffer (Qiagen, UK) at room temperature for 15 min. The column was washed again with 350 µl RW1 buffer and centrifugation, then was washed twice with 500 µl RPE buffer (diluted 1:5 with 100% ethanol) and centrifuged at 8000 g, room temperature (first time for 15 sec and second time for 2 min). RNA was eluted from the column in 30 μ l RNase-free water by centrifugation at 8000 g, 1 min, room temperature. Total RNA extracted was quantified using the Agilent RNA 6000 Nano assay with the Agilent 2100 bioanalyser (Agilent Technologies, UK) and stored at -80 °C.

2.7.2.2 Sense-strand RNA amplification

Messenger RNA was linearly amplified from 250 ng of DNase treated total RNA using the SenseAmp kit (Genisphere, USA). Firstly, total RNA was annealed to 50 ng oligodT24 primer and 500 ng random nanomer primer in 11 µl at 80 °C for 10 min and cooled on ice for 2 min. First strand cDNA was reverse transcribed from the RNA/primer mix using Superscript II reverse transcriptase (Invitrogen, UK), 500 µM of each dNTP, in the presence of 4 µl 5× first strand buffer (250 mM Tris-HCl (pH 8.3), 375 mM KCl, 15 mM MgCl₂), 10 mM DTT, 1 µl Superase-In RNase inhibitor in a total volume of 20 µl. The reaction was performed at 42 °C for 2 h. First strand cDNA was purified using the MinElute PCR purification kit (Qiagen, UK). The cDNA was bound to the MinElute column in the presence of 500 μ l PB buffer. Column was centrifuged at 16,000 g, 1 min; washed in 750 µl PE buffer and centrifuged again. 500 80% ethanol was then passed through the column by centrifuging for 2 min and the column air-dried for 5 min. The purified cDNA was eluted in 10 μ l EB by first equilibrating for 2 min at room temperature and then centrifuging for 2 min. In order to allow RNA amplification directly from the purified first strand cDNA (i.e. no second strand synthesis), a poly-dT tail was added to the 3'-end of the first strand cDNA using 2 µl terminal deoxynucleotidyl transferase (Genisphere, USA), 2 mM dTTP nucleotides, 2 µl 10× reaction buffer, in 20 µl total volume. The reaction took place at 37 °C, 3 min, and was quenched at 80 °C, 10 min. A poly-dA-anchored T7 promoter was then annealed to the poly-dT tail by incubating at 37 °C for 10 min. The 3'-end of the cDNA was then 'filled in' to complete the double stranded T7 promoter using 1 µl Klenow enzyme, 400 µM of each dNTP, 1 µl 10× reaction buffer, by incubating at room temperature for 30 min. In vitro transcription (IVT) was then performed from the T7 promoter-modified cDNA (12.5 µl) using 2 µl T7 enzyme mix, 8 µl nucleotide mix, 2.5 µl 10× T7 reaction buffer. Amplified RNA was synthesised by IVT reaction at 37 °C for 6 h in a Primus Multiblock thermal cycler (MWG, UK). Amplified RNA was stored at 4 °C until clean

up, using the RNeasy kit (Qiagen, UK), in a similar procedure to section 2.7.2.1, except that 250 μ l of 100% ethanol was used instead of 350 μ l 70% ethanol; and that the RW1 buffer washing and DNase treatment steps were eliminated. The concentration of the amplified RNA was measured using the Agilent 2100 bioanalyser (Agilent Technologies, UK).

2.7.3 Reference mix

The reference RNA comprised the following cell lines in the stated proportions: U937 (monocytes) 25%, HeLa (epithelial) 25%, SSCEM (T-cells) 15%, Ramos (B-cells) 15%, Huh7 (liver) 5%, MRC5 (fibroblasts) 5%, HHV7-SupT1 (DC) 5%, HHV6A-SupT1 (U1102) 2.5%, HHV6B-SupT1 (Z29) 2.5%. RNA of the individual cell lines were extracted, purified and quantified separately and mixed before RNA amplification or labelling.

2.7.4 Fluorescent labelling of RNA

Fluorescent dye-labelled cDNA was synthesised from 250 - 1000 ng mRNA or amplified RNA (aRNA) using the CyScribe First-Strand cDNA labelling system-dCTP (Amersham Biosciences, UK). Briefly, mRNA or aRNA were annealed to 1 µl each of oligo-dT and random primers, in a total volume of 11 µl, at 70 °C for 5 min, and cooled to room temperature for 10 min. This RNA/primer mix was added to the labelling reagents: 100 units reverse transcriptase, 5 mM DTT, 50 µM Cy5- or Cy3-dCTP, 1µl dNTPs (except dCTP), 4 µl 5× CyScript buffer in 20 µl total volume. Reversetranscription reaction was performed at 42 °C for 90 min in Primus Multiblock thermal cycler (MWG, UK), to produce fluorescent dye-labelled first strand cDNA.

Labelled cDNA was denatured with 20 μ l sodium hydroxide (0.1 M), 5 μ l EDTA (0.5 M, pH 8.0) at 70 °C for 10 min, and neutralised with 20 μ l hydrochloric acid (0.1M). To suppress hybridisation of repetitive DNA, 3 μ g of Cot-1 DNA (Invitrogen, UK) was added. Unincorporated nucleotides and dye were removed by washing with 450 μ l

Tris-EDTA buffer (TE) through a Microcon YM-30 filter device (Millipore, UK) by centrifugation at 14,000 g for 6 min. This was repeated 2 more times with 300 μ l TE and centrifuging for 8 min. Purified cDNA was collected by inverting the Microcon column and centrifuging at 1000 g for 3 min, to a final volume of 13.5 μ l or less.

2.7.5 Hybridisation onto microarray

For each hybridisation, 250 - 1000 ng of Cy5- and Cy3-labelled cDNA samples were mixed with the hybridisation mix to block non-specific binding and denature the DNA: $12 \ \mu l \ 20 \times$ SSPE (saline sodium phosphate EDTA buffer, Sigma), $1.1 \ \mu l$ of EDTA (0.5 M, Sigma), $2 \ \mu l$ of poly-dA (40-mer, $8 \ \mu g/\mu l$, Thermal Hybaid), $2 \ \mu l$ yeast tRNA (4 $\mu g/\mu l$, Sigma), $1 \ \mu l \ 10\%$ SDS, in TE buffer to a final volume of 45 μl . This mixture was incubated for 2 min at 98 °C to denature the labelled DNA, followed by incubation at 37 °C for 20 min. $1 \ \mu l \ of \ 100 \times Denhardt's$ (Sigma, UK) was added to prevent non-specific binding and the mixture centrifuged at 16,000 g, 15 min to remove any insoluble materials.

Meanwhile the pre-blocked array (section 2.6.3) was placed into a hybridisation chamber and warmed to 65 °C on hot block. A glass coverslip (22×64 mm, thickness No. 0, SLS, UK) was washed in isopropanol (Sigma, UK) for 30 seconds and immediately spun dry in empty centrifuge tubes at 200 g for 2 min. When the labelled cDNA was ready 40 µl was pipetted onto centre of the array, over which the coverslip was gently but quickly placed to allow dispersal of liquid by capillary action. 150 µl of $4 \times$ SSPE (65 °C) was added to the chamber to maintain humidity and the lid immediately closed. Arrays were hybridised at 65 °C by immersion in waterbath for 16 -20 h.

The hybridised array was removed from the chamber and immersed in $2 \times SSPE$, 50 °C until the coverslip was detached. The array was then sequentially washed in three wash solutions of increasing stringency at room temperature. $2 \times SSPE$ for 2 min, $1 \times SSPE$ for 2 min and $0.1 \times SSPE$ for 3 min. After the third wash, the array was immediately

spun dry at 200 g, 2 min. The array was either scanned (section 2.8) or stored in the dark.

2.8 Array scanning and data collection

Each array was scanned using a GenePix 4000 microarray scanner and the GenePix Pro 4.0 software (Axon Instruments, USA), at 10 μ m resolution. Cy3 and Cy5 were simultaneously excited at 532 nm and 635 nm respectively and the fluorescence measured by photomultiplier tubes (PMTs). The voltages across the PMTs were adjusted (generally set around 500V) so that the signals from the array elements were balanced. The software generated a combined image from the measurements, on which a template was overlaid to identify each spot to their corresponding gene. Array spots from which no signal could be detected were flagged as not found by the GenePix software. The image was also examined by eye to flag any gene spots damaged by debris, scratch or smudging. For each spot the software calculated the Cy5 and Cy3 median intensities of all pixels in the spot, as well as the local background signals. The data were exported to the TM4 software package (TIGR, UK), an open-source program developed for microarray data analysis, for processing and analysis (Saeed et al., 2003).

2.9 Array data processing and normalisation

Array data were converted into *mev* file by the TIGR ExpressConvert software (TM4, TIGR, UK), and read into the MIDAS software (TM4, TIGR, UK). The program first removed bad quality data already flagged by the GenePix software or manually. It then removed questionable data based on a 'stringent tolerance policy', so that data with a signal-to-noise ratio of <2.0 in any one channel were removed. This was followed by a low intensity filter, in which data with a integrated intensity below 10,000 were removed. Filtered data were then normalised to correct for various systematic and experimental errors. First of all, total intensity normalisation was applied to balance the values in both channels. This was achieved by applying a normalisation factor (f) to the Cy5 signal of each spot using the Cy3 value as the reference, so that:

Corrected Intensity (Cy5) = Original Intensity (Cy5) /f

The normalisation factor was calculated as follows:

$$f = \frac{\sum Intensity(Cy5)}{\sum Intensity(Cy3)}$$

This was followed by the LOWESS normalisation in the global mode, to correct for intensity-dependent local errors. This was performed by applying LOWESS algorithm (Cleveland 1979) with a smoothing parameter of 0.33, to adjust the Cy5 value using the Cy3 value as a reference.

2.10 Clustering analysis

Normalised data from multiple arrays to be compared were assembled and imported into the Mev software (TM4, TIGR, UK) for multiple array comparison. A low percentage filter was set to include only data points that were present in 80% or 100% of the total number of arrays compared. The data were mean-centred by gene (to set the mean ratio of each gene across all arrays to 0), and by array (to set the mean ratio of all gens within one array to 0) to scale the data so that relative comparisons could be made. Genes and arrays were clustered by average-linkage hierarchical clustering, using the uncentred Pearson correlation as the similarity metrics, based on the method developed by Eisen *et al*, 1998.. Clustered results were visualised as a dendrogram in the MeV software.

2.11 Significance Analysis of Microarrays (SAM)

Significance Analysis of Microarrays (SAM), enables binary comparisons of microarray data, a gene-specific t-test that gives significantly regulated genes, and includes a permutation algorithm that allows estimation of error, termed the false discovery rate (FDR) (Tusher et al, 2001). SAM has been incorporated into the TMev software (TM4, TIGR, UK). Genes that passed the 100%-presence filter (i.e. genes that passed SNR filtering across all arrays analysed) were separated into two groups (inhibitor-treated vs

untreated) and their expression signals compared by SAM. The FDR was determined by a user-defined Delta value, which was set at 3% (i.e. 3 genes in every 100 genes found to be significantly regulated may be falsely identified). Lists of significantly upand down-regulated genes were assembled and used in subsequent hierarchical clustering for visualisation as a dendrogram in TMev.

2.12 Reverse transcription-PCR (RT-PCR)

2.12.1 RT-PCR reaction

Specific HHV-7 transcripts in total RNA, mRNA or amplified RNA were amplified by reverse transcription PCR (RT-PCR) using the OneStep RT-PCR kit (Qiagen, UK). Briefly, 0.05 pg – 50 ng template RNA were reverse transcribed into cDNA by the Omniscript and Sensicript reverse transcriptase enzymes present in the OneStep enzyme mix (Qiagen, UK), in the presence of 0.6μ M HHV-7 ORF specific forward and reverse primers (Table 2.3), 400 μ M of each dNTP, and 10 μ l 5× reaction buffer, in a final volume of 50 μ l. Reverse transcription was allowed to take place at 50 °C for 90 min in Primus Multiblock thermal cycler (MWG, UK). This was followed by a heating step at 95 °C for 15 min, which inactivated the reverse transcriptases and activated the HotStarTaq DNA polymerase which was also present in the OneStep enzyme mix. The newly synthesised cDNA was amplified by PCR with 35 cycles of 94 °C, 30 sec; 55 °C, 1 min; 72 °C, 1 min; followed by a final extension step of 72 °C for 10 min. In control reactions to test for DNA contamination, template RNA was added after the heating step, or the OneStep enzyme mix was replaced by HotStarTaq enzyme (Qiagen, UK). RT-PCR products were analysed by agarose gel electrophoresis as in section 2.4.4.

2.12.2 Gel purification

One volume of the RT-PCR reaction was separated on a 1% w/v agarose gel as in section 2.4.4. The band of the correct size was excised from the gel and DNA purified with the QIAquick gel extraction protocol (Qiagen, UK). 3 volumes of buffer QG (v/w)

were added to the gel slice and incubated at 50° C for 10 min. 1 volume of isopropanol was added, the solution vortexed and then applied to a QIAquick column. The solution was passed through the column by centrifugation (14,000 g, 1 min), washed with 500 µl buffer QG and then with 750 µl buffer PE. DNA was eluted in 30 µl of buffer EB.

2.12.3 Sequence analysis of gel bands

The identity of the DNA extracted from agaorse gels (2.12.2) was confirmed by sequence analysis as detailed in section 2.4.6.3.

CHAPTER 3

PRODUCTION OF HHV-7 MICROARRAYS

3.1 Introduction

3.1.1 Reasons for HHV-7 microarray production

Viruses are intracellular parasites that rely on their continual interaction with the host cellular environment and their exploitation of cellular processes for replication and spread. Understanding these interactions at the molecular level contributes valuable knowledge to virus life cycles, pathogenesis and host defence. Conventional single gene studies provide useful information about the functions of individual genes or proteins, but they do not offer an integrated view on the complex multifactorial infection processes. One approach to study viral-host interaction is by gene expression profiling. Transcription is one of the earliest steps of relaying genomic information to the synthesis of proteins, which are in general the active effectors of many biological processes in both the host and the viral life cycle. Since transcription is a tightly regulated process, changes in transcription during infection usually reflects changes at the protein and functional level (Guerra et al., 2004; Tang et al., 2005). This in turn reveals the biological state and activities in the infected cells (Geiss et al., 2003; Song and Stinski, 2002), or the stages of the viral life cycle (Ahn et al., 2002; Chambers et al., 1999; Jenner et al., 2001; Paulose-Murphy et al., 2001). By applying microarray technology to gene expression profiling, it is possible to monitor the transcriptional changes of not just one gene, but thousands of genes simultaneously (section 1.2). Therefore a global picture of the complicated changes that take place during infection can be obtained.

Microarrays have been used to study global expression changes of cellular genes during infection by viruses, while other studies have monitored changes in viral gene expression (see section 1.2.4). Herpesviruses, in particular, have been the focus of many microarray-based gene expression studies, as the high-throughput technology is ideal for analysing their large DNA genomes. Viral gene expression profiling has been performed on HSV (Stingley et al., 2000), HCMV (Chambers et al., 1999), Kaposi's sarcoma associated herpesvirus (KSHV) (Jenner et al., 2001; Paulose-Murphy et al., 2001), murine herpesvirus type 68 (MHV68) (Ahn et al., 2002) and HHV-6 (Ohyashiki et al., 2005). The microarrays used contain 85 to 180 probes that target the corresponding sequences in the viral genomes. These arrays also contain variable

numbers of human probes, from 4 (Ohyashiki et al., 2005) to 88 (Paulose-Murphy et al.), mainly housekeeping genes for control and normalisation purposes. The probes were produced either by PCR (Ahn et al., 2002; Jenner et al., 2001; Paulose-Murphy et al., 2001) or by oligonucleotide synthesis (Chambers et al., 1999; Ohyashiki et al., 2005; Stingley et al., 2000) before robotic deposition onto membrane (Ahn et al., 2002; Jenner et al., 2001) or glass (Chambers et al., 1999; Ohyashiki et al., 2002; Jenner et al., 2001) or glass (Chambers et al., 1999; Ohyashiki et al., 2005; Paulose-Murphy et al., 2001; Stingley et al., 2000) arrays. Apart from one study (Ohyashiki et al., 2005), all viral microarrays were produced in-house. These viral microarray studies have characterised the transcription profiles of herpesviruses over the course of infection *in vitro* and *in vivo*; some have identified particular genes of interest. More detailed summary of the findings is provided in Chapter 5.

The above mentioned studies focussed on either the host or viral gene expression. However, since viral infection is a multifactorial process involving both parasitic and cellular contributions, analysis of both viral and host genes in parallel during infection would be highly desirable. This could be achieved using a DNA microarray with probes to both viral and host genes to measure transcript abundance of viral and cellular genes together. One study has included probes for HIV as well as cellular genes on the custom-made arrays (van 't Wout et al., 2003). However the study only focussed on cellular expression while the HIV gene probes were used as positive controls rather than for transcription analysis.

Human herpesvirus type 7 (HHV-7), with its high prevalence and apparently nonpathogenic nature, is believed to be a virus very well adapted to humans (section 1.1). It is therefore useful to study the transcription profiles of the virus and the host to understand the life cycle and pathogenesis (or the lack of it) of HHV-7 infection. In order to obtain the global pattern of host and viral gene expression, we have made a DNA microarray containing probes that target all predicted HHV-7 genes and a subset of cellular genes. This is the first description of an integrated host-pathogen microarray for HHV-7.

The aim of work described in this chapter was the production and testing of HHV-7 microarrays, forming the foundation for further analysis in the later sections.

3.2 Results

3.2.1 Production of HHV-7 array probes

To determine global expression patterns of human herpesvirus type 7 (HHV-7) genes, a custom-made DNA microarray was created to measure the abundance of HHV-7 gene transcripts. This involved production of the HHV-7 array probes, which were then spotting onto glass-slides by automated robotics. The production of HHV-7 array probes involved a series of DNA amplification, cloning, sequence confirmation and re-amplification, represented on a diagram (Figure 3.1).

3.2.1.1 Primer design and initial PCR amplification from genomic DNA

Array probes were designed to target the 86 predicted open reading frames (ORFs) in HHV-7, positioned towards the 5'-end of each ORF (Figure 3.2). This positional preference was to distinguish between ORFs with co-terminal 3'-sequences, for example U53 and U53.5. As an attempt to detect any previously unidentified genes, 10 probes (X1-X10) were also designed for some of the larger intergenic 'gap regions' in the HHV-7 genome between known ORFs (Figure 3.2). The production of array probes involved initial PCR amplification of DNA segments approximately 300 bp in length using specific primers (Table 3.1) designed using the Primer3 program (Rozen, 2000), based on the following criteria – optimal melting temperature of 52 °C, primer length of 20 bases, and prevention of secondary structures. Genomic DNA extracted from HHV-7 infected SupT1 cells were used as the template for PCR. The amplified DNA segments were analysed by agarose gel electrophoresis (Figure 3.3), which showed that amplification was successful for 85 ORFs and 4 intergenic probes (X1, X3, X5, X7). One ORF (U77) failed to amplify with the initial primers (U77-1) therefore primers were re-designed (U77-2) to successfully amplify the U77 probe (Table 3.1). No further attempts to amplify the six failed intergenic probes (X2, X4, X6, X8, X9, X10) were performed.

Table 3.1 HHV-7 array primers

				-4	Bevere primer (5' 2')	a ta atb	probe
ORF	Strand"	Acc. No.	Forward primer (5'-3')	start	Keverse primer (5-3)	start	5120
DR1	+ (2)	AAC40718	CTGCTGCAACCACAGAACAT	5		306	302
DR6	+ (2)	AAC40719	AGAAATGCTCCGCGCTGT	9	TGAAGTTCCAGGCTCCAGTT	320	312
U2	•	AAC40720	CCACAAAAGGACAGCAGACA	759	CAGATTCTCCGCTTTTGGAT	1076	318
U3	-	AAC40721	CAAGTTCAAACCCCTCTGGA	863	GCAGATTTGGGGTCTGACTT	1172	310
U4	-	AAC40722	TGAAGAAACGCGAGAAAACC	1319	TGTTTGTAAAGGGCCGCTAA	1609	291
U7	- (2)	AAC40723	GGTCCGGTTCTCGGTAATTT	2409	TGTTCCTGTGTGCAATGATG	2695	287
U8	-	AAC40724	AATTGGCACGTCTTTCCTTC	769	TTGAGACGATGGCAGAACAA	1079	311
U10	+	AAC40725	AGCGGAGCGTTCGAAATAA	16	CTGGTGACGCGACAATAAAA	328	313
U11	-	AAC40726	TGCCCAGTTTAAGCAAATCTT	1852	GGATTTCGGATGAAGCAAAG	2220	369
U12	+ (2)	AAC40727	TACGCCAACAGTACGGAAAG	159	CTGTATGTGCAGGACGCAC	442	284
U13	+	AAC40728	CAACAGCCATCTAATTGCAGA	13	AAAATGGTGACGAACGAACTT	292	280
U14	+	AAC40729	CGAAAACTGCAACTTTGTGC	45	TTCATCGAATCGTCGCATAG	360	316
U15	- (3)	AAC40730	TGTGTGAATTGAGCGACGAT	421	GAAACCTGGAGAAGACAACGA	719	299
U17	- (2)	AAC40731	GCAGCTGCTGTTGTTACGTC	695	TAGGCATGCACTCGCAAAT	1036	342
U18	-	AAC40732	TGTTAGCCCGAGAAGCAAA	531	CAAATTTGAAATGTCACCAGAA	830	300
1110	_	AAC40733	ABGCTGACACCCATTCTTCG	708	ATGTGGCTTAACGACGATGC	967	260
1120	-	AAC40734	TCACCATTAAAATTGTGGGTGAT	848	TGTGGCACATAAGTTGACAATG	1123	276
1124	-	AAC40735	TOLOTRON TONOTON	061	ATCTTCCTCTCTTCCCCTTCCC	1284	324
021	-	AAC40735		407		450	324
023	-	AAC40730	TGCTGCAATTAAAATCACAAGG	127	GIACGCCCACTCTGGAAATG	402	320
U24	-	AAC40737	TTCGTACCCCCATATCGAAC	8	TGCATCAGGAAAATCTATCACC	1/0	109
U24A	-	AAC40738	AGGGATATTTAGAGCACCTGA	3	CAACCATTTTTTCCCAGAGAC	155	153
U25	-	AAC40739	CAAATGCCCTCTGTCACCTT	679	AGCGAATTTGTGCTGTGTCA	952	274
U26	-	AAC40740	CCATTGAATGCGTTGGAAAT	549	GGGCCTTGTTTTAGGGAATG	853	305
U27	-	AAC40741	CTGATGCCGTGACTAAAAACC	696	TCGTAGCAACAGAGAACACCA	1090	395
U28	-	AAC40742	ATGGAATCAAATTCCGAAGC	2120	AAAACATGGGTCTCTTGAGGA	2390	271
U29	-	AAC40743	CTCCAAACAAAGACACCTTTTC	477	TTTGCTAGCTTTTGGTGTTCC	772	296
U30	+	AAC40744	TGCGGCTGAAAGTATCACAG	108	AATAGGCGATGCCAATTCCG	427	320
U31	+	AAC40745	GCAGGAAGTACAAATCAAAACG	13	TTGTTTCGCCAAAATTAGAACT	280	268
U32	-	AAC40746	CCGGTAAAGAGATGATACTTG	847	TGACATCACACTGCGATACG	1136	290
U33	-	AAC40747	CCTTGCATTGCCTGTACAAA	119	ATGGTTCGTACAGTCATTTGC	371	253
U34	-	AAC40748	CGCATCTCGGTTTCGTCTA	361	TCCCGCAGACTTTCGTGT	715	355
U35	-	AAC40749	ATCTCCGAGTGACTGCATTG	95	AACGACAATGCCAAAACAGA	309	215
U36	+	AAC40750	TGGAACTCCGATTCCTTTTC	16	GGGAGTTCTAGGTGCAGCAT	327	312
U37	+	AAC40751	ACACGTAGACTGAGACGAGCA	16	TGCCACCATAGCATCTGAAG	339	324
U38	-	AAC40752	ACCAGATGGAATAACGAAATGT	2704	ATGGATCTGGTGTCCTTCTTT	3042	339
U39	-	AAC40753	TTGGCATTGGCAAAGTTGTA	2124	AGCTGACTTTGTCATGACTGGA	2407	284
U40	-	AAC40754	GCTTGCATTCATTGAGTTGG	1827	TTGCAATCATTGTGTGTGTCTTG	2157	331
U41	-	AAC40755	CAGGCTTTCGATTTCCGTTA	3012	CAATATGTACCGCAGCATGG	3362	351
U42	-	AAC40756	CCGGAATACTTGCGTAAACC	1217	GTATTAGGCGGCAACGATA	1524	308
U43		AAC40757	CACTTCCTTCTGGAACATGAGA	2256	TGTGTTTGCAACGGAATATGA	2575	320
1)44	+	AAC40758	CTACCAGATGCTCTATTCTGAGC	51	TTTTCATCA	2070	202
1145	_	AAC40759	CCGTTTTTCBGGB ATTCCB AG	963		4420	293
1146	<u>.</u>	AAC40760		27		1139	211
1147		AAC40761		21		247	221
1140	-	AAC40701		010	CAATGCTAGTGTTCACGTTGAT	923	308
140		AAC40762	TIGTATGCGAAACAGGTTTGA	1/22	AAATGGATGGAAACACTGGAA	2032	311
049	+	AAC40763	GCCCCCTGTTAGAAGACGTA	18	TCTGTTGCTTGTCGTAGTTGC	368	351
050	+	AAC40764	ATGTACAGACAGAATGCCAGA	206	CCAGCTTCTAAATCGGACAC	619	414
051	+	AAC40765	CAAATTGGAAATTACTGGCAGA	20	TGATACGTTCCAGTGCCAAC	316	297
052	-	AAC40766	GCTGAGCTGCATTTTCTTCA	435	TCGAACAACTCAACCTCCAA	738	304
053	+	AAC40767	GTGGCAGGCTTCCTATGTGT	16	GGATCTGGTGGCAGAGTTTT	308	293
053.5	+	AAC40768	AGCTAGTCTTCCCGCTCCTC	27	TCATCATCCGAATCTGTTGG	323	297
U54	-	AAC40769	TGCGCTCAACTATTTGGACA	1026	TGGAATCCTGTTATGTGGTCTG	1364	339
U55A	-	AAC40770	ACCCTTTTCAAGATAGGCAAA	910	TGCAGCCAGAACTCAATTTT	1220	311
U55B	-	AAC40771	GGAAAAACTCCCATCTGCTC	941	CAAAGAAGAATTCCAGGGTAAA	1240	300
U56	-	AAC40772	GGCAGAATCAGCTCCCAGT	500	AACTCACTAATGCCGTCATCC	818	319
U57	-	AAC40773	GGCGCAGTTACGTCGTTT	3713	CACAGCTGAAATTTTCCCAAA	4024	312
U58	+	AAC40774	TCACTGTGCTAAAAATGTTCCA	54	TATATAGAACCGGGCCTTCG	355	302
U59	+	AAC40775	ACAATGAGACAGCAACTGTGC	53	AAAAACAGTTTGGGCTTTTCA	323	271

USB USB USB USB USB ACC40780 ACCGCABATCCATCTTGRA 4704 TSCAATTCAACCGCGACGTATCA 5085 382 U62 + AAC40776 AATTCGTCAACTGTGRATTC 77 ATTCCACGCGACGACGTAT 176 100 U63 + AAC40776 CATTGTATATACGGAATGGTGTGRATTC 30 AGGGTCTCATATTTGTGGGAGGTGCCCAGA 406 251 U64 + AAC40779 CCCCGAATATCGTTGTGTGG 30 GGCGTAATAGTGCGCCCATT 339 297 U67 + AAC40781 GTCCATGCAGATATCGTCAGTAGAC 22 AGCCATTTTATATAGAGCGCCCCAAA 33 301 U68 + AAC40783 GCAGTTATAGACCCCAAAA 2 GCCATTCCATTGTATAGCGGACAT 303 303 301 U70 + AAC40785 ATGGGCTCAAATAGTCCCCCAAAA 2 CCCATTCTCCATTGTAGGGACATAG 30 302 200 200 200 200 200 200 200 200 200 200 200 200 200 200 200 200 200 <	ORE	Strand*	Acc. No.	Forward primer (5'-3')	start ^b	Reverse primer (5'-3')	start ^b	probe size°
Cut2 + AAC40776 AMATTGCCGAGACTGGAATTC 77 ATTCCACCGCGAGACGATA 176 100 U62 + AAC40777 ATTCCATCCACACTGC 30 ACGGTCCCCATATTGGTCC 299 270 U64 + AAC40778 GCTGTATATACGGAATGGACTGTC 30 ACGGTCATTGCCAATATCGTGTGTGG 33 301 U65 + AAC40778 CCCCGCAATTTATATAAGAACACC 2 ACGGATGTTCACTGCACATT 303 301 U68 + AAC40782 GTCACGGATAATGCACACAA 3 GCCACTGTGTTGAACACCCCCAA 6 CATCCACTGTGTGTGAGAGATTCACTGTGTGTGAGA 301 303 302 U70 + AAC40783 CTGCCACTTAAACACCCCCAA 6 CATCCACTGTGTAGACACCCCAA 303 302 U71 + AAC40786 TTTGTGGAGATCACTCCTG 11 TTGTTCTGTGGTAGACGAT 303 302 U72 - AAC40786 TTTTTGGAGAGATTACCC 74 TGTCTCAGGTTT 303 322 U73 - AAC40786 TTTTTGGAGAGATTTACC 74 TGTGTGGAGATTCCTTTGAGA	1166	-(2)	AAC40780	ACCGCAGATCCATCTTTGAA	4704	TGCAATTCAGAAAGCGTATCA	5085	382
L023 + ACC40777 XTGTCASTCANTCACAGTTGC 30 ACGGTCTCCATATTTSGCTCT 299 270 U64 + ACC40778 CCTTGTATATACGGAARGAC 156 GTANGGAGGTGCCACATT 399 227 U65 + ACC40779 CCACGAATTACGGTTGTGGG 39 GCGCGAATGACTCCCATAT 399 227 U66 + ACC40781 GCGGCTATATATATATATAAAC 2 AGGANTTCACTT 303 302 U67 + ACC40782 GTCACTGGATGAATTATATATATAAAC 2 GCGCATCTGTGTACGAATTCACTCCTCGG 311 306 U70 + ACC40784 TOGGCTAAAGTGCGATACATTACCCACAA 2 CGCATCTGTATACGAACGGAACT 303 302 U71 + ACC40786 TOGGCTAAAGGTGGAACT 303 302 200 U72 - ACC40788 TOGGCTGAAGGTGAACA 1 GCCATCTTACAACCAGTGGAACA 604 213 U73 + ACC40780 TOGTGCGAACTCCATTG 104 GGCATCTTACTAACCAACGGCGAACA 694 213 U75 -	1162	+	AAC40776	AAATTTGTCTGAACTGTGATTTC	77	ATTTCCACCGCGAGCGTAT	176	100
U64 + AAC40778 GCTITITATATACGGANTGGAC 156 GTATGGGAGGGTGCTCCAGA 406 251 U65 + AAC40778 GCTATGTATATACGGANTGGAC 22 AGCGATGTCTCACATGGTCA 343 322 U67 + AAC40781 GCTGGTAATATGTTGTGTGG 83 GCCGTAATAGGTCACATT 303 301 U68 + AAC40782 GCAGCTAAGATACGCCCANA 2 GCGGTAAGGGGTTCACAT 303 302 U70 + AAC40786 TGGCAATAGGTCACACCANA 2 CGGATGTAGGAGATT 303 302 U71 + AAC40786 TCGCCCATCGAAGTTATATATACAC 1 GCGGTGTAGGAGATT 303 302 U73 + AAC40786 TCCCCCATCCAAGTTATATATACC 674 TGGCATGAGGGTACAGATT 322 289 U74 + AAC40787 TTTTTGTGGGAGACATA 31 GACGTTAACGAGACATT 322 289 U75 - AAC40789 CTCAGCGGAATCTCTTCAGC 422 ACTAMACATCGTCCAATT 567 3055 U77-2 +<	1163	+	AAC40777	ATGTCAGTCAATCACAGTTGC	30	ACGGTCTCCATATTTGGCTCT	299	270
U65 + AAC40779 CCACQUATATCOTTOTOTOG 93 GGCGTAATAGGTCGCACALT 389 297 U67 + AAC40781 GCTGCANTTERLANGANACGAC 22 AGCGATGTCACACALT 343 322 U68 + AAC40782 GTCGCATTAGATTAATANAAC 3 GTCGGTAGGATTCACTT 303 301 U69 + AAC40783 GCGGTAAGATTAACATTAATANAAC 3 GTCGGTAGGATTCACTTT 303 302 U70 + AAC40784 TGGCATAGGTCCAAAT 1 TTCTTCTGGTAGGACACAT 303 302 U71 + AAC40785 TTGTGGATGGCAAAAT 1 GCGATGTACAAACGACAATAGATCCATTGT 328 288 U73 + AAC40786 TTTTTGGAAGCCAAATCCCTTGT 104 GGCATGTACAAAACGACAATAGACGACGATT 328 288 U75 - AAC40798 CTCGACGCAAATCTCCATGC 217 TCGCGCGAAACA 642 213 U76 - AAC40796 TGCTGACGCAGATCAA 386 AGGACAATCTATACACAGAT 563 GGACGTGTACACATTG 7703 <td< td=""><td>U64</td><td>+</td><td>AAC40778</td><td>GCTTGTATATACGGAATGGAC</td><td>156</td><td>GTATGCGAGGTGCTCCAGA</td><td>406</td><td>251</td></td<>	U64	+	AAC40778	GCTTGTATATACGGAATGGAC	156	GTATGCGAGGTGCTCCAGA	406	251
U67 + AAC40781 GCTGCARTTATANARGANCGAC 22 AGCGATGTTCTCACTGGTCA 343 322 U68 + AAC40782 GTCACTGCATGATTATANARC 3 GTCGGTGACGGATTCACTT 303 301 U69 + AAC40783 GGCATGACTGCATGATTATCCACCCANA 6 CATGGACTCATCCATCGG 311 306 U70 + AAC40786 TGGCGATTGAGCACAA 2 CGCATCTTGTGAGAGAACG 200 200 U71 + AAC40786 TGCCCCATGCACAA 1 TTCTTGGTGTAGAGAGAGAGAACG 200 200 U72 - AAC40786 TGCCCCATGCAGATTATATCC 674 TGGCATGAGAGAGAGAACG 228 288 U74 + AAC40786 TGCCCCAGGATCTCTTGGC 104 GGCATGTACAAACGAGCGAACA 644 213 U75 - AAC40791 GATATGGAGAGACAATTAA 283 CATTTGGATGACAACA 647 322 321 U75 - AAC40791 GATATGCAACACTCCAA 283 CATTTGGATGACAACA 647 305 U77-1	065	+	AAC40779	CCACGAATATCGTTGTGTGG	93	GGCGTAATAGGTCGCACATT	389	297
U68 + AAC40782 GTCACTGCATGAATTAATAAAAC 3 GTCGGTGACGGATTTCACTT 303 301 U69 + AAC40783 GCCACTTAAGACACCCCAAA 6 CATGCACTCATCACGTTTG 303 302 U70 + AAC40784 TGGCATCAAGACACCCCAAA 2 CGCATCTATGACGACAT 303 302 U71 + AAC40786 TGGCCTCAAGGGCTTAAG 2 CGCATCTTGATACGAACGAC 200 U72 - AAC40786 TCGCCCCACCCAAGTTATATCC 674 TGGCATGAAGAAGAAG 2289 U74 + AAC40788 TGCGTGAGCATCTTCTTG 104 GGCATGTACAAAACCGCCAAT 322 289 U75 - AAC40780 TGCTGAGGATCACCGTAA 1365 AGGGACTATAAACAACCGCCGAT 328 248 U77-1 + AAC40790 TGCTGAGGATGCACGTAC 287 CACTTGGCTGCAACATTG 703 339 U77-2 + AAC40791 GAATTGCAACCGTTGCC 217 TCTGCGGATGCCCAATTG 7103 329 U77-4 + AAC40793	U67	+	AAC40781	GCTGCAATTTATAAAGAAACGAC	22	AGCGATGTTCTCACTGGTCA	343	322
Construction Construction<	1168	+	AAC40782	GTCACTGCATGAATTAATAAAAAC	3	GTCGGTGACGGATTTCACTT	303	301
0000 + AAC40784 TGGCAATAGATTAGGCACAAA 2 CGCATCTTGTAAGCGAACAT 303 302 U71 + AAC40785 ATGGCAATAGATTAGCCACAAA 1 TTCTTCTGGTATAGCGAACAT 303 302 U72 - AAC40785 ATGGCATCAAGTGCTATAATC 674 TGGCATTGAAGCAGAGTAGATGG 1040 367 U73 + AAC40786 ATGCTGAAGTCACTCCTTG 104 GGCATGTACAAAACGGCGAACA 694 213 U74 + AAC40789 CTCAGCGAAATCCATTCTTTGAGC 482 ACTAAACAAGCGCGAACA 694 213 U75 - AAC40790 TGCTAAGGGTACACGTTGC 217 TCTGGCGATGACGTTCAACAGTG 1703 339 U77-1 + AAC40791 GAATTGCCAGGGTAA 283 CACTTGGCATTTGATGCAGCATGA 322 221 U81 - AAC40791 GAATTGCCAGAGGATAGAGAT 2 CCTTTGCCGTTTTTCTGA 322 221 U82 - AAC40795 TGGTGAAAGGATTCCATA 283 CACGTTGGCAATTGATTACA 749 297 U84 - AAC40796 TCGCTGAATCGTGGAA 331 GGCGGCCGGAGGGACAAGG <td>1169</td> <td>+</td> <td>AAC40783</td> <td>GCAGCTTAAGACACCCCAAA</td> <td>6</td> <td>CATGCACTCAATCCATCTCG</td> <td>311</td> <td>306</td>	1169	+	AAC40783	GCAGCTTAAGACACCCCAAA	6	CATGCACTCAATCCATCTCG	311	306
U71 + AAC40785 ATGGGCTCDAAGTGCTGTAA 1 TTCTTCTGGGTTAACGAAACG 200 201 U72 - AAC40786 TCCCCCCATCCAAGTTATATCC 674 TGGGCTCTAAAGGAAAGGT 310 GACGCTCTTAATATTCGCTGT 328 288 U73 + AAC40788 TTTTTGGATGGGTTGGACA 31 GACGCTCTTAATTCGCAGT 322 289 U74 + AAC40788 TTTTTGGATGGGTTGGACA 31 GACGCTGTAAAACGGCAACA 694 213 U75 - AAC40788 CTCAGCGAAATCCTCTCTAGC 482 ACTAAACAACGGCAACA 694 213 U76 - AAC40791 GCAAAACAAGCGGTGCC 217 TCTGGGATGACAGTCAATTA 541 325 U77-2 + AAC40791 GCAAAACAAGCAGCATCAA 283 CACTTGGCTTTCCAATTA 543 322 321 U81 - AAC40795 TGTTGAAAGTATCCGTGA 453 GGATGCTGAAAACTTCA 749 297 U84 - AAC40795 TGTTGCAAGTTTGCTGTG 33 AACTTTGTTAAGGCAAAGG 913 281 U86 - AAC40798 CAGGCATGTTAAGGAAA	U70	+	AAC40784	TGGCAATAGATTACGCACAAA	2	CGCATCTTGTAAGCGGACAT	303	302
U72 - AAC40786 TCCCCCATCCAAGTTATATCC 674 TGGCATTGAACAGAGTAGATG 1040 367 U73 + AAC40787 TTTTTGGAATGGGTTGGACA 31 GACGCTCTTTATATCCCCTGT 328 288 U74 + AAC40788 TCTGTGAGACTCATCCTTG 104 GGCATGTACAAAACGGCCAACG 644 213 U75 - AAC40789 CTCACGCGGAATCTTCTTCTAGC 482 ACTAAACAAACGGCGAACG 1703 339 U76 - AAC40790 TGCTAAGGGTGCACCTCTACAC 282 CATAACAACAGCGACAT 541 325 U77-1 + AAC40791 GAAATCGAACGAGCAGCATCAA 283 CACTTGGATGACATTCCA 761 322 321 177-2 + AAC40792 TGTTGGAAGCAGCATCCAA 283 CACTTGGCATTCCATTG 567 305 177 177 176 322 321 181 - AAC40793 TGCTGAAAGCTATCCGTGA 283 CACTTTGGATGACATTCCATTGA 749 297 U82 - AAC40795 TTGTGGAAAGTTTGCGTTG 397 ATTCTTAGTAGGCAAGTGAGGATTCA 749 297 U84 - AAC40796	U71	+	AAC40785	ATGGGCTCAAAGTGCTGTAA	1	TTCTTCTGGGTTAACGAAACG	200	200
U73 + AAC40787 TITTIGGANGGTTIGGACA 31 GACGCTCTTTAATTICGCTOT 328 298 U74 + AAC40788 ATGCTGAGGCTAACCTCCTTG 104 GGCATGTACAAAACCGCCGAAC 664 213 U75 - AAC40789 CTCAGCCGAACTCTTCTCTAGC 482 ACTAAACAAACGGCCGAAC 664 213 U76 - AAC40790 TGCTAAGGGTGCAGCGTAA 1365 AAGGACAATCTAACACGGTGC 1703 339 U77.1 + AAC40791 GAAAACAGGGCATCAA 283 CACTTGCGCTGATTCCCATTG 587 305 U79 + (3) AAC40792 TGATGGAAGATGAATCGTGA 453 GGATGCTGGAAAACTTCA 749 292 321 U81 - AAC40793 TGCTGTAATCCGTGA 453 GGATGCTGGAAATTCA 749 297 U82 - AAC40795 TGTTGCAAATCGTGATG 597 312 CCTTTCCTAGGTAACCATTCA 3403 AACGTTGAATTCA 442 331 U84 - AAC40796 CCAGCATCGTTATATTGTG 512 TGCCTTTCTTAAGGTGACTGGTT 3142 313 U85 - AAC407979	U72	-	AAC40786	TCCCCCATCCAAGTTATATCC	674	TGGCATTGAGCAGAGTAGATG	1040	367
U74 + AAC40788 ATGCTGAGACTCACTCCTTG 104 GGCATGTACAAAACCGCGAAT 392 289 U75 - AAC40789 CTCAGCGGAATCTCTTCTAGC 482 ACTAAACAAACGGCGAACA 694 213 U76 - AAC40790 TGCTAAGGGTGACGGTAA 1385 AGGGACATGTATATCACACGGTCC 1703 339 U77.1 + AAC40791 GCAAAACAAGCAGCGTTCCA 283 ACCTTTGGCTTCCCAATAT 541 325 U77.2 + AAC40791 GCAAAACAAGCAGCATTCA 283 GCATTTGCCAAACTTTG 322 321 U81 - AAC40793 TGCTGAAAGCTAATCCGTGA 453 GGATGTGGAAAACATTCA 749 297 U82 - AAC40795 TTGTGCAAAGTTGTTGTTG 397 ATTCTTATCCCATTCTAAACC 716 320 U84 - AAC40796 CCAGGACTGTTATTGTTGG 512 TGCCTTCTTGTAAGGCAAATT 442 331 U85 - AAC40796 CGAGGAAGCTCTTCTTCAA 3405 TGGACAACCATCGAAATT 342 U91 + (2) AAC40798 CGAGGAAGGTGTTGTTC 1 ATTCCTTGGTGTGTGTTGC 365	U73	+	AAC40787	TTTTTGGAATGGTTTGGACA	31	GACGCTCTTTAATTTCGCTGT	328	298
U75 - AAC40789 CTCASCCGANTCTCTTAGC 482 ACTAAACAAACACGCGGAACA 694 213 U76 - AAC40790 TGCTAAGGGTGTAGCGTAA 1365 AGGGACAATCTATACACAGTG 1703 339 U77.1 + AAC40791 GAATTTGCCAACGGTGCC 217 TCTGCGAGACGTCCAATAT 541 325 U77.2 + AAC40791 GCAAAACAAGCAGCATTCAA 283 CACTTTGGCTTCCCATTG 587 305 U79 +(3) AAC40792 TGGTGAAAGCATCAA 283 CACTTTGGCTTGGAAAGATTCCA 749 297 U81 - AAC40794 CCAGTTGTGCCCTGATTCT 397 ATTCTTATCTCCATTGAAAGC 716 320 U84 - AAC40795 TTGTTGCAAAGTTGTGCTG 633 AACTGTTTGTAAGCCAAAGG 913 281 U86 - AAC40796 CCAGGTGTAATTGTCTGCTG 321 GGCCTTACAGTGGAATT 824 331 U90 -(3) AAC40796 CTGGGGTGTATCTGAGGAT 3405 TGGCGTTGTCTGCTGA 345 324 U91 +(2) AAC40800 ATGGAGAGGGGATAGTTC 1 ATGCGGGGGTGCCATGT <t< td=""><td>1174</td><td>+</td><td>AAC40788</td><td>ATGCTGAGACTCACTCCTTG</td><td>104</td><td>GGCATGTACAAAACCGCATT</td><td>392</td><td>289</td></t<>	1174	+	AAC40788	ATGCTGAGACTCACTCCTTG	104	GGCATGTACAAAACCGCATT	392	289
U76 - AAC40790 TGCTAAGGTOTCAGCGTAA 1365 AGGGACAATCTATACACAGTG 1703 339 U77-1 + AAC40791 GAAATTGCTCAACCGTTGCC 217 TCTGCGATGACGTTCCAATAT 541 325 U77-2 + AAC40791 GCAAAACAAGCAGCATTCAA 283 CACTTGGCTTCCCAATAT 567 305 U79 + (3) AAC40792 TGATGGGAAGATACAAGAAT 2 CCTTTCCCTGGTTTTCCTGA 322 321 U81 - AAC40795 TGGTGAAAGCTAATCCGTGA 453 GGATGCTGGAAAACATTCA 749 297 U82 - AAC40795 TGGTGAAAGCTAATCGTGA 353 AACTGTTTGTACAACTTCAA 749 297 U84 - AAC40796 CCAGCATCGTTATTGTGTG 633 AACTGTTTGAGGAAAACCT 716 320 U84 - AAC40797 TTCTCCACTGCTGAAGGAT 3186 GCACTACGATGGTAAACCAT 3414 297 U90 -(3) AAC40798 CGAGGAAGCTCCTCTCAA 3405 TGGGACAACCATCGATGAT 324 U91 + (2) AAC40800 TGGAGAAGCTCCTCAAA 22 CACTCGGGTTGTCTCTC 345	U75	-	AAC40789	CTCAGCCGAATCTCTTCTAGC	482	ACTAAACAAACACGCCGAACA	694	213
U77-1 + AAC40791 GAATTTGCTCAACCGTTGCC 217 TCTGCGATGACGTTCCAATAT 541 325 U77-2 + AAC40791 GCANAACAAGCAGCATTCAA 283 CACTTTGCCTTTCCCATTG 587 305 U79 +(3) AAC40792 TGATTGCGGAGGATAGAGAAT 2 CCTTTCCCTCGTTTTCCTGA 322 321 U81 - AAC40793 TGCTGAAAGCTAATCCGTGA 453 GGATGCTGAAACATTCA 749 297 U82 - AAC40796 CCAGTTGTGCCCTGATTCT 397 ATTTCTTATCTCAATCCATTCAAAGG 913 281 U84 - AAC40796 CCAGTAGTGCCTGATTCG 512 TGCCTTTCTAAGGCCAAAGG 913 281 U85 - AAC40797 TTTCTCCACTTGCTTGAGGAT 318 GCACTACAGCTGGGTT 3614 297 U90 -(3) AAC40799 AACGTTTGTCATCAA 3405 TGGCGTGAACCATCTGACT 3723 319 U91 +(2) AAC40800 ATGGAGAGGGATAGTTTC 1 ATGTTGCTGGTTACTTGC 345 324 U95 + AAC40800 ATGGAGAGGGATAGTTTC 1 ATGTGGGCACTGAGCACATGCCTTGC	U76	-	AAC40790	TGCTAAGGGTGTCAGCGTAA	1365	AGGACAATCTATACACAGTG	1703	339
U17-2 + AAC40791 GCAAAACAAGGAGCATTCAA 283 CACTTTGGCTTCCCATTG 587 305 U79 + (3) AAC40792 TGATTGGGAGGAATAGAGAAT 2 CCTTTCCCTTCGTTTTCTGA 322 321 U81 - AAC40793 TGCTGAAAGCAGCATTCT 397 ATTCTTCATCTCCATTCTAAACC 716 320 U82 - AAC40794 CCAGTTGTGCCTGATTCT 397 ATTCTTATCTCACATTCTAAACC 716 320 U84 - AAC40795 TTGTTGCAAAGTTGTCGTG 633 AACTGTTGTAAGGCAAAAGG 913 281 U85 - AAC40796 CCAGCATCGTTATGTGGGTG 512 TGCCTTCTTGAAGGCAAAGG 913 281 U86 - AAC40797 TTCTCCCCTTGCTTGAGAT 3405 TGGCAACCATCGTAT 3723 319 U90 -(3) AAC40797 TTGGTGACAGGCAAAA 22 CACTCGGGATGTTTCCTGC 289 289 U900 -(10) AAC40800 ATGGAGATGGGATAGTTTC 1 ATGTTCCTGGTTGCTTGC 306 302 DR6' +(2) AAC40802 CTGCTGCAACACAGACAT 5 GCGAGGTGTGCTGCTG <td< td=""><td>U77-1</td><td>+</td><td>AAC40791</td><td>GAATTTGCTCAACCGTTGCC</td><td>217</td><td>TCTGCGATGACGTTCCAATAT</td><td>541</td><td>325</td></td<>	U77-1	+	AAC40791	GAATTTGCTCAACCGTTGCC	217	TCTGCGATGACGTTCCAATAT	541	325
U79 + (3) AAC40792 TRATTGCGAGGATAGAGAAT 2 CCTTTCCTGGA 322 321 U81 - AAC40792 TGCTGAAAGCTAATCCGTGA 453 GGATGCTGGAAAACATTCA 749 297 U82 - AAC40794 CCAJTTGTGCCAAAGTTGCGTG 633 AACTGTTTGTAAGGCCAAAGG 913 281 U84 - AAC40795 TTGTTGCAAAGTTGTCGT 397 ATTTCTTATCTCAAAGG 913 281 U85 - AAC40796 CCAGCATCGTTAATTGTCG 512 TGCCTTTCTAAGGTGCAATTT 842 331 U86 - AAC40797 TTTCTCCACTGCTTGAGGAT 3405 TGGACTACAGCTGAACCAT 3723 319 U91 + (2) AAC40799 AACGGAGATACAACTTCTAA 3405 TGGACTACAGCTGGACACT 3245 324 DR1' + (2) AAC40800 ATGGAGAGTGGGAATGAT 5 GCGAGGCTCGAGCT 306 302 302 DR6' + (2) AAC40802 CTGCTGCAACCACAGAAACAT 5 GCGAGCTCGAGCACACA 271 272 X2 n/a n/a TAGGAGAGTGTGAACCACACAGGGAG 23 TGAAGTTCCAGGCCCA	1177-2	+	AAC40791	GCAAAACAAGCAGCATTCAA	283	CACTTTGGCTTTCCCATTTG	587	305
UB1 AAC40793 TGCTGAAAGCTAATCCGTGA 453 GGATGCTGGAAAACATTTCA 749 297 UB2 AAC40794 CCAGTTGTGCCCTGATTCT 397 ATTCCTTATCTCCATTCTAAACC 716 320 UB4 AAC40795 TTGTTGCAAAGTTTGTCGTTG 633 AACTGTTTGTAAGGCCAAAGG 913 281 UB5 AAC40796 CCAGGCATCGTTAATTGTCG 512 TGCCTTTCTTAGGGCCAACGAGGTGGTT 842 331 U86 AAC40797 TTTCTCCACTTGCTTGAGGGAT 3318 GCACTACAGCGACGATGGTT 3614 297 U90 -(3) AAC40799 CGAGGAAGCCCTTCTCAA 3405 TGGACAACCATCGAACGAT 3723 319 U91 +(2) AAC40799 AACGTGTATCAAGGCCAAAA 22 CACCGGGTGTTACTGCT 3645 324 U95 + AAC40800 TGGAAAGCACACAAGAACAT 5 GCGAGGCTCTGTCCT 306 302 DR1' +(2) AAC40800 TGGAACCACAGAACAT 5 GCGAGGCTCCAGTT 320 298 X11 n/a n/a CGAGGACCACAGAACAT 5 GCGAGGCCCAGGT 302 204 X32 n/a	U79	+ (3)	AAC40792	TGATTGCGGAGGATAGAGAAT	2	CCTTTCCCTCGTTTTTCTGA	322	321
UB2 AAC40794 CCAGTTGTGCCCTGATTCT 397 ATTTCTTATCTCATTCTAAACC 716 220 U84 - AAC40795 TTGTTGCAAAGTTTGTCGTTG 633 AACTGTTTGTAAGGCCATAAGG 913 221 U85 - AAC40796 CCAGCTGGTTAATTGTCG 512 TGCCTTCTTAGGGCCAAGGGGT 3614 297 U90 -(3) AAC40799 TTTCTCCACTGCTTGAGGAT 3318 GCACTACAGCGGGGGT 3614 297 U90 -(3) AAC40799 CGAGGGAAGCTCCTTCTTCAA 3405 TGGACAACCATCTGAAAGCCAT 3723 319 U91 + (2) AAC40799 AAACGTGTATCAAGGCCAAAA 22 CATCCGGGTTGTTTCCTGCT 345 324 U95 + AAC40800 ATGGAGGTGTGTGTTCC 2322 CCAATACGGTCCCATTGA 2645 324 U96' + (2) AAC40802 CTGTGCACCAGGAAG 23 TGAAGTTGTCCCAGGTT 320 298 X1 n/a n/a CGAGGTGTACCACAGAACCAT 5 GCGAGCTGTGGCTGCAGGT 320 294 X2 A	U81	-	AAC40793	TGCTGAAAGCTAATCCGTGA	453	GGATGCTGGAAAACATTTCA	749	297
U84-AAC40795TTGTTGCAAAGTTGTGTGG33AACTGTTGTAAGGCCAAAGG913281U85-AAC40795CCAGCATCGTTAATTGTCG512TGCCTTTCTTAAGGCCAAAGG913281U86-AAC40797TTTCTCCACTTGCTGAGGAT3318GCACTACAGCTGACGTGACTT842331U86-AAC40798CGAGGAAGCTCCTTCTCAA3405TGGACAACCATCGAAACCAT3723319U90-(3)AAC40799AAACGTGTATCAAGGCCAAAA22CATCCGGGTTGTTTACTGCT345324U95+AAC40800ATGGAGAGGGGATAGTTC1ATGTTCCTGGTTCCTGC289289U100-(10)AAC40801TGGAACGCACACAGAAAAA22CCAATACGGTCCACTTGA2645324DR1'+(2)AAC40802CTGTGCAACCACAGAAACAT5GCGAGCTTGTGCTTGTCC306302DR6'+(2)AAC40803CTGTTCAGCTCCAGGCCAAG23TGAAGTCCAAGGCCATCA271272X2n/an/aCGAGGTCATCATACAACGG1CAATGGTCAACGACAGGG203204X3n/an/aGCTGCGTACTCATGACAG1TATCTCCGGGAAGTGGG203204X4n/an/aGCTGGGTACTCATGCACG1TATCTCCGGGAAGCAC264265X5n/an/aGACGGTCATCATACAACGG1CAATGGTCAACGAAGCATCA301302X4n/an/aGTCTGCGTAACTACAACGG1TATCTCTCAGGAGAGAACAC264265X5n/an/aAGGCAACCTAAAACGGCG1TATCTCTCAGGAGAGA	1182	_	AAC40794	CCAGTTGTGCCCTGATTCT	397	ATTICTTATCTCCATTCTAAACC	716	320
U85 - AAC40796 CCAGATCGTTAATTGTTCG 512 TGCCTTTCTTAGGTGCAATTT 642 331 U86 - AAC40797 TTTCTCCACTTGCTTGAGGAT 3318 GCACTACAGCTGAATT 642 331 U80 -(3) AAC40797 TTTCTCCACTTGCTTGAGGAT 3318 GCACTACAGCTGAAACAT 3723 319 U91 +(2) AAC40799 AAACGTGTATCAAGGCCAAAA 22 CATCCGGGTGTTTACTGCT 345 324 U95 + AAC40800 ATGGAGAGTGGGGAAAGTTC 1 ATGTTCCTTGGCT 366 302 U100 -(10) AAC40801 TGGAGAACCACAGAAACAT 5 GCGAGCTTGTGCTGTGCTC 306 302 DR6' +(2) AAC40803 CTGTGCGAACCACAGAACAT 5 GCGAGCTTGAGCACCA 271 272 X2 n/a n/a CGAGATCGTAACGAACTTG 1 CAATGGTCACGAGCATCA 271 272 X2 n/a n/a CGAGGTCGTCATCATCATCAGGG 1 GCATGGCACACACACAGGG 203 204 X3 n/a n/a GTGCGCGCAAGGGA 1 GCTTCCTGAGCACACAGGAAGGG 229	1184	_	AAC40795	TTGTTGCAAAGTTTGTCGTTG	633	AACTGTTTGTAAGGCCAAAGG	913	281
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	1185	_	AAC40796	CCACCATCGTTAATTGTTCG	512	TACCTTTCTTAGGTGCAATTT	842	331
U90 -(3) AAC40798 CGAGGAAGCTCCTTCTTCTAA 3405 TGGACAACCATCTGAAACCAT 3723 319 U91 +(2) AAC40798 CGAGGAAGCTCCTTCTTCAA 3405 TGGACAACCATCTGAAACCAT 3723 319 U95 + AAC40799 AAACGTGTATCAAGGCCAAAA 22 CATCCGGGTTGTTACTGCT 345 324 U95 + AAC40800 ATGGAGAGTGCGGGATAGTTC 1 ATGTTCCTGGTTTCCTGC 289 289 U100 -(10) AAC40801 TGGAATGCAAGAGTGTTGTTTC 2322 CCAATACGGTCCACCTGA 2645 324 DR1' +(2) AAC40802 CTGCTGCAACCACAGAACAT 5 GCGAGCTTGTGTCTC 306 302 DR6' +(2) AAC40803 CTGTTCAGCTCCAGCCAAG 23 TGAAGTTCCAGGCATCAA 271 272 X2 n/a n/a TTAGCGTCATCTAACAACGG 1 CTATCTTGCGGAAGGCCATCA 271 272 X2 n/a n/a TTAGCGTCATCTATCAACAACGG 1 CTATCTTGCGGAAAGGCATCA 301 302 X4 n/a n/a GAGGAACCTGATAGAAGGTGCGG 1 TATCTCTGGGAAAGAGGACACC <td>U86</td> <td>_</td> <td>AAC40797</td> <td>TTTCTCCACTTGCTTGAGGAT</td> <td>3318</td> <td>GCACTACAGCTGACGTGGTT</td> <td>3614</td> <td>297</td>	U86	_	AAC40797	TTTCTCCACTTGCTTGAGGAT	3318	GCACTACAGCTGACGTGGTT	3614	297
U91 +(2) AAC40799 AAACGTGTATCAAGGCCAAAA 22 CATCCGGGTTGTTACTGCT 345 324 U95 + AAC40800 ATGGAGGTGGGGATAGTTC 1 ATGTTCCTTGGTTTCCTTGC 289 289 U100 -(10) AAC40801 TGGAATGCAGGTGTTCTC 2322 CCAATACGGGTCCACTTGA 2645 324 DR1' +(2) AAC40802 CTGCTGCAACACAGAAGTGTTGTTC 2322 CCAATACGGGCCCACGTGA 2645 302 DR6' +(2) AAC40803 CTGTTCAGCTCCAGCAAG 23 TGAAGTCCAGGCTCACTGA 271 272 X1 n/a n/a CGAGATCGTAACGAACTTG 1 CAATGGTCACTGAGCCACA 203 204 X3 n/a n/a CGAGATCGTAACGAACTGA 1 TATCTCCGTGACAAGCAAC 264 265 X5 n/a n/a GCTGCGTTACTTCTGTCAC 1 TATCTCCGTGACAAGCAAC 264 265 X5 n/a n/a CAACTGATAAGAAGTAGGGG 1 TTTGGGAGTAAGCAAC 264 265 X5 n/a n/a AGCACTTAATAGAAGTAGGGG 1 CTACCTCAGTAGGAACAC 264	1190	- (3)	AAC40798	CGAGGAAGCTCCTTCTTCAA	3405	TGGACAACCATCTGABACCAT	3723	310
U95+AAC40800ATGGAAGTGGGGATAGTTC1ATGTTCCTTGGTTTCCTTGC289289U100-(10)AAC40801TGGAATGCAGAGTGTTGTTC2322CCAATACGGTCCCACTTGA2645324DR1'+(2)AAC40802CTGCTGCAACCACAGAACAT5GCGAGCTTGTGTCTC306302DR6'+(2)AAC40803CTGTTCAGCTCCAGCCAAG23TGAAGTTCCAGGCTCCAGTT320298X1n/an/aCGAGATCGTAACGAACTTG1CAATGGTCACCAGGAGCACCA271272X2n/an/aTTAGCGTCATCATACAACGG1GCTTACTTTGCGGAAGCAACAC301302X3n/an/aGTCTGCGTTACTTCTGTCAC1TATCTCCGTGACAAAGCAACAC264265X4n/an/aGTCTGCGTTACTTCTGTCAC1TATCTCCGTGACAAAGCAACAC264265X5n/an/aGTCTGCGTTACTTCTGTCAC1TATCTCCGTGACAAAGCAACAC264265X5n/an/aCAACCTGATAGAGGACTGCAG1CAACCTTCAGTAGTGTGGG229230X6n/an/aAGGCAACCTAAAACTGGCTG1CAGCGGAAGGAGAATTG339340X7n/an/aAGGCAACCTAAAACTGGCTG1ATGCAGTAGCAAAACC359360X8n/an/aTGTGTGCAGTTTGGAAACATG1TTTCTTATATTGCTAGCCTCC400401X9n/an/aAGGAACTGCGGGTGCCTTC1GGGGGGCAGAAGCCAATAACACAAAACC359360X10n/an/aAGGAACTGCGGGGCGGGGGAATTGGGGGCCGCGGAATTCACTAG<	1191	+ (2)	AAC40799	ABACGTGTATCABGGCCABAA	22	CATCCGGGTTGTTTACTGCT	345	324
U100- (10)AAC40801TGGAATGCAGAGTGTTGTTC2322CCAATACGGTCCACTTGA2645324DR1'+ (2)AAC40802CTGCTGCAACCACAGAACAT5GCGAGCTTGTGCTTGTCTC306302DR6'+ (2)AAC40803CTGTTCAGCTCCAGCCAAG23TGAAGTTCCAGGCTCCAGTT320298X1n/an/aCGAGATCGTAACGAACTTG1CAATGGTCACGAGGCTCCAGTT320298X2n/an/aTTAGCGTCATCATACAACGG1CAATGGTCACTGAGCCATCA271272X2n/an/aGTCTGCGTTACTCTGTCAC1TATCTCCGTGACAAAGCAAC301302X4n/an/aGTCTGCGTTGATCTCTGTCAC1TATCTCCGTGACAAAGCAAC264265X5n/an/aGACCTGATAGAAGGAGG1TTTGGGAGTCAAGCAAGCAC264265X5n/an/aCAGTTGAATGAGGACTGCAG1TATCTCCGTGACAAGCAAC264265X5n/an/aCAGTTGAATGAGGACTGCAG1TATCTCCGTGACAAGCAAC264265X6n/an/aCAGTTGAATGAGGACTGCAG1CAACCTTCAGTAGGAGCAACC264265X6n/an/aAGGCAACCTAAAAACTGGCTG1CAACCTTCAGTAGCAAAGCAATG339340X7n/an/aAGGCAACCTAAAACTGGCTG1CTGCAGAGAAGAATACC344345X8n/an/aGTGTGTGCAGTTTGAGAACATG1TTTCTTATATTGCTAGCACTAAAAACC359360X10n/an/aAGGAACTGCGGGCGGGGGAATTGGCGGCCGCGAATTCACTAG <td>U95</td> <td>+</td> <td>AAC40800</td> <td>ATGGAGAGTGGGGATAGTTTC</td> <td>1</td> <td>ATGTTCCTTGGTTTCCTTGC</td> <td>280</td> <td>280</td>	U95	+	AAC40800	ATGGAGAGTGGGGATAGTTTC	1	ATGTTCCTTGGTTTCCTTGC	280	280
DR0(10)AAC40802CTGCTGCACCACAGAACAT5GCGAGCTTGTGCTGCTC306302DR6'+ (2)AAC40803CTGTTCAGCTCCAGCCAAG23TGAAGTTCCAGGCTCCAGCT271272X1n/an/aCGAGATCGTAACGAACTTG1CAATGGTCACTGAGCCATCA271272X2n/an/aTTAGCGTCATCATACAACGG1GCTTACTTGCGGAACGAGG203204X3n/an/aGTCTGCGTTACTTCTGTCAC1TATCTCCGTGACAAAGCATAC301302X4n/an/aCAACCTGGATAGAAGTAGGGG1TTTGGGAGGTGAGGACAC264265X5n/an/aCAGTTGAATGAAGGACTGCAG1CAACCTTCAGTAGTGTGTGG229230X6n/an/aAGGCAACCTAAAACTGGCTG1CAGCGAAAGAGGTGAAGAATTG339340X7n/an/aAGGCAACCTAAAACTGGCTG1ATGCAGTAGCAATACCAAATCC344345X8n/an/aTGTGTGGCAGTTGAGAACCT1GGGGGCACCAAAAACC359360X10n/an/aAGGAACTGCGGTTGCCTTC1GGGGGCCGCGAAATCTACTAG301301pGEM-TGCATGGCGGCCGCGGGGAATTGGCGGCCGCGGAAATCACTAG301301301301pGEM-TFTGTAAAACGACGGCCGCGGGGAATTGGCGGCCGCGGAAATCACTAG300301301pGEM-TFTGTAAAACGACGACGGCCAGTGGGGAATTTGGCCCTCGAAGGTCCAAACCTACGAGG300301pGEM-TFTGTAAAACGACGACGGCCAGTGGGCGGCCGCGGAAATCACTAGSP6gGGAATTGGCCTCGAAGTGGCGGCCCCGAAGG </td <td>11100</td> <td>- (10)</td> <td>AAC40801</td> <td>TGGALATGCAGAGTGTTGTTGTTC</td> <td>2322</td> <td>Сартасаатсосастаа</td> <td>2645</td> <td>324</td>	11100	- (10)	AAC40801	TGGALATGCAGAGTGTTGTTGTTC	2322	Сартасаатсосастаа	2645	324
DR6'+ (2)AAC40803CTGTTCAGGTCCAGGCCAAG23TGAAGTTCCAGGCTCCAGGT320298X1n/an/aCGAGATCGTAACGACTTG1CAATGGTCACTGAGCCACGA271272X2n/an/aTTAGCGTCATCATACAACGG1GCTTACTTGCGGAAAGTGGG203204X3n/an/aGTCTGCGTTACTTCTGTCAC1TATCTCCGTGACAAAGCATAC301302X4n/an/aCAACCTGATAAAAGTAGGGG1TTTGGGAGGTGAGCAAAGCAC264265X5n/an/aCAGTGAATGAGGACTGCAG1CAACCTTCAGTAGTGTGTGG229230X6n/an/aAGGCAACCTAAAACTGGGCTG1CAGCGGAAAGGGTGAAGATTG339340X7n/an/aAGGCAACCTGAAACC1ATGCAGTAGCAATACCAAATCC344345X8n/an/aTGTGTGCAGTTTGAGAACATG1TTTCTTATATTGCTAGCCCTCC400401X9n/an/aAGGAACTGCGGTTGCCTTC1GGGGGCGCGAAAACCTAAAAACC359360X10n/an/aCAGTTAGTCGCTGCGGGTGCCTTC1GGCGGCCGCGGAAATGAACC301301pGEM-TGCATGGCGGCCGCGGGGAATTGGCGGCCGCGGAAATTGACTAG1TCCAAACCTACGAAGTATACC359360X10n/an/aCAGTGGCGCCGCGGGGAATTGGCGGCCGCGGAAATTGACTAG1TCCAAACCTACGAGG1pGEM-TGCATGGCGGCCGCGGGGAATTGGCGGCCGCGGGAATTGACTAG301301301pGEM-TTTGTAAAACGACGACGCCAGTGGGGAATTGGCCCTCGAAGG301301 <td>DR1'</td> <td>+ (2)</td> <td>AAC40802</td> <td>CTGCTGCAACCACAGAACAT</td> <td>5</td> <td>GCGAGCTTGTGCTTGTCTC</td> <td>306</td> <td>302</td>	DR1'	+ (2)	AAC40802	CTGCTGCAACCACAGAACAT	5	GCGAGCTTGTGCTTGTCTC	306	302
X1n/an/aCGAGATCGTAACGAACTTG1CAATGGTCACTGAGCCATCA271272X2n/an/aTTAGCGTCATCATACAACGG1GCTTACTTTGCGGAAGTGGG203204X3n/an/aGTCTGCGTTACTTCTGTCAC1TATCTCCGTGACAAAGCATAC301302X4n/an/aGTCTGCGTTACTTCTGTCAC1TATCTCCGTGACAAAGCATAC301302X5n/an/aCAACCTGATAAGAAGTAGGGG1TTTGGGAGGTCAAGCAAGCAC264265X5n/an/aCAGTGAATGAGGACTGCAG1CAACCTTCAGTAGTGTGTGG229230X6n/an/aAGGCAACCTAAAACTGGGCTG1CAGCGGAAAGAGGTGAAGATTG339340X7n/an/aAGGCAACCTGAAAACTGGGCTG1ATGCAGTAGCAATACCAAATCC344345X8n/an/aTGTGTGCAGTTTGAGAACATG1TTTCTTATATTGCTAGCACCTCC400401X9n/an/aAGGAACTGCGGTTGCCTTC1GGGTGAGAACCCAAAAAACC359360X10n/an/aCAGTTAGCTGCTGGAAAGC1TCCAAACCTACGCAATGTATC300301pGEM-TGCATGGCGGCCGCGGGGAATTGGCGGCCGCGGAAATTGACTAGGGCGGCCCCGGAGGGTTTGTAAAACGACGGCCAGTGGGGAATTGGCCCTCGAGGTTT3TTGTAAAACGACGACGGCCAGTGGGGGACCTCGAAGGSP6 promoter primerTTCTATAGTGTCACCTAAAT	DR6'	+ (2)	AAC40803	CTGTTCAGCTCCAGCCAAG	23	TGABGTTCCBGGCTCCBGTT	320	202
X1I/dI/dCONCRETED/INFORMATIONICARTAGATCATACAAGCA211212X2n/an/aTTAGCGTCATCATACAACGGIGCTTACTTTGCGGAAGTGGG203204X3n/an/aGTCTGCGTTACTTCTGTCACITATCTCCGGGAAAGCATAC301302X4n/an/aCAACCTGGATAGAAGTAGGGGITTTGGGAGTCAAGCAAAGCATAC264265X5n/an/aCAGTGAATGAGGACTGCAGICAACCTTCAGTAGTGTGTGG229230X6n/an/aAGGCAACCTAAAACTGGCTGICAGCGGAAAGAGGTGAAGATTG339340X7n/an/aAGGCAACCTAAAACTGGCTGICAGCGGAAAGAGGTGAAGATTG339340X8n/an/aATTCCATGGGCCCTGGAAACCIATGCAGGTAGCATACTACAATCC344345X8n/an/aTGTGTGCAGTTTGAGAACATGITTTCTATATTGCTAGCCCTCC400401X9n/an/aAGGAACTGCGGTTGCCTTCIGGGGGGCAAACCAAAACC359360X10n/an/aCAGTTAGTCGCTGGAAAGCITCCAAACCTACGCAATGATACC300301pGEM-TGCATGGCGGCCGCGGGGAATTGGCGGCCGCGGAAATTCACTAGGGCGGCCCCCGAGGGSP6 promoter primerTTCTATAGTGTCACCTAAAT	X1	· (2)	n/a	CARATCATACCAACTTC	1		271	200
X2IV2IV2INTOCONSTRUCTIONINTOCONSTRUCTION203204X3n/an/aGTCTGCGTTACTTCTGTCAC1TATCTCCGTGACAAAGCATAC301302X4n/an/aCAACCTGATAGAAGTAGGGG1TTTGGGAGAAAGCATAC264265X5n/an/aCAGTTGAATGAAGGACTGCAG1CAACCTTCAGTAGTGTGGG229230X6n/an/aAGGCAACCTAAAACTGGCTG1CAGCGGAAAGAGGTGAAGATTG339340X7n/an/aAGGCAACCTAAAACTGGCTG1CAGCGGAAAGAGGTGAAGATTG339340X7n/an/aATTCATGGGCCCTGAAACC1ATGCAGTAGCAATACTACAAATCC344345X8n/an/aTGTGTGCAGTTTGAGAACATG1TTTCTATATTGCTAGCCCTCC400401X9n/an/aAGGAACTGCGGTTGCCTTC1GGGTGAGAACCCAATAAAACC359360X10n/an/aCAGTTAGTCGCTGGAGAAGC1TCCAAACCTACGCAATGTATC300301pGEM-TGCATGGCGGCCGCGGGGAATTGGCGGCCGCGGGAATTCACTAGGGCGGCCCCCGAGGGTTTTGTAAAACGACGACGGCCAGTGSP6 promoter primerTTCTATAGTGTCACCTAAAT	x2	n/a	n/a	TTACCGTCATCATACAACGG	. 1	CTTACTTCCCCA & CTCCC	203	204
X4 n/a n/a CAACCTGATAGAAGTAGGGG 1 TTTGGGAGTCAAGCAAGCAAC 264 265 X5 n/a n/a CAGTTGAATGAGGAGCAGG 1 CAACCTTCAGTAGTGTGG 229 230 X6 n/a n/a AGGCAACCTAAAACTGGCTG 1 CAGCGGAAAGAGTGG 339 340 X7 n/a n/a AGGCAACCTGAAACC 1 ATGCAGTAGCAAGCAACC 344 345 X8 n/a n/a TGTGTGCAGTTTGAGAACATG 1 TTTCTTATATTGCTAGCCCTCC 400 401 X9 n/a n/a AGGAACTGCGGTTGCCTTC 1 GGGTGAGAACCCATAAAACC 359 360 X10 n/a n/a CAGTTGGCGGCGCGGGGAATT GGCGGCCGCGGAATTCACTAG 1 TCCAAACCTACGCAATTAAACC 300 301 pGEM-T GCATGGCGGCCGCGGGGAATT GGCGGCCCCTCGAAGG 1 TCCAAACCTACGCAATTAGCATGG 300 301 pGEM-T TTGTAAAACGACGACCGGCCAGTG GGGGAATTTGGCCCTCGAGGG GGGGAATTGGCCCTCGAGGG 501 300 301	X3	n/a	n/a	GTCTGCGTTACTTCTGTCAC		TATCTCCCTCACAAAACCATAC	205	204
X1 INC IN	X4	n/a	n/a	CAACCTGATAGAAGTAGGGG	1	TTTGGGAGTCAAGCAAGCACAC	264	302 265
X6 n/a n/a AGGCAACCTAAAACTGGCTG 1 CAGCGAAAGAGGGGAAGATTG 339 340 X7 n/a n/a ATTTCATGGGCCCTGAAACC 1 ATGCAGTAGCATACTACAATCC 344 345 X8 n/a n/a TGTGTGCAGTTTGAGAACATG 1 TTTCTTATATTGCTAGCCCTCC 400 401 X9 n/a n/a AGGAACTGCGGTTGCCTTC 1 GGGTGAGAACCTAAAACC 359 360 X10 n/a n/a CAGTTAGTCTGCTGGAAAGC 1 TCCAAACCTACGCAATGTATC 300 301 pGEM-T GCATGGCGGCGCGGGGGAATT GGCGGCCGCGGAATTCACTAG 1 TCCAAACCTACGCAATTAGG 1 T7T3 TTGTAAAACGACGGCCAGTG GGGAATTTGGCCCTCGAGGG SP6 promoter primer TTCTATAGTGTCACCTAAAAT	X5	n/a	n/a	CAGTTGAATGAGGACTGCAG	1	CAACCTTCAGTAGTGTGTGTG	204	200
X7 n/a n/a ATTTCATGGGCCCTGAAACC 1 ATGCAGTAGCATACTACAATCC 344 345 X8 n/a n/a TGTGTGCAGTTTGAGAACATG 1 TTTCTTATATTGCTAGCCCTCC 400 401 X9 n/a n/a AGGAACTGCGGTTGCCTTC 1 GGGTGAGAACCCATAAAACC 359 360 X10 n/a n/a CAGTTAGTCGCGGTGCCTTC 1 GGGTGAGAACCATGATATTC 300 301 pGEM-T GCATGGCGGCGCGCGGGGAATT GGCGGCCGCGGAATTCACTAG 1 TTCTATAGTGGCCCTCGAGG 300 301 T7T3 TTGTAAAACGACGGCCAGTG GGGAATTTGGCCCTCGAGGG SP6 promoter primer TTCTATAGTGTCACCTAAAAT 559 540	X6	n/a	n/a	AGCALCTAALCTGGCTG	1	CACCORAGAGAGAGAGAGAGATAG	220	230
X8 n/a n/a TGTGTGCAGTTTGAGAACATG 1 TTTCTTATATTGCTAGCCCTCC 400 401 X9 n/a n/a AGGAACTGCGGTTGCCTTC 1 GGGTGAGAACCCATAAAACC 359 360 X10 n/a n/a CAGTTAGTCGCGGTGCCTCC 1 GGGTGAGAACCCATAAAACC 300 301 pGEM-T GCATGGCGGCCGCGGGGAATT GGCGGCCGCGGAAATTCACTAG 1 TTCTATAGTGGCCCTCGAGG 300 301 T7T3 TTGTAAAACGACGGCCAGTG GGGAATTTGGCCCTCGAGG GGGAATTGGCCCTCGAGG 5P6 promoter primer TTCTATAGTGTCACCTAAAT	X7	n/a	n/a	ATTTCATGGGCCCTGAAACC	1	ATGCAGTAGCATACTACAATCC	344	245
X9 n/a n/a AGGAACTGCGGTTGCCTTC 1 GGGTGAGAACCCATAAAACC 359 360 X10 n/a n/a CAGTTAGTCTGCTGGAAAGC 1 TCCAAACCTACGCAATGTATC 300 301 pGEM-T GCATGGCGGCCGCGGGGAATT GGCGGCCGCGGGGAATT GGCGGCCCCCGGGGAATTCACTAG T7T3 TTGTAAAACGACGGCCAGTG GGGGAATTGGCCCTCGAGGG SP6 promoter primer TTCTATAGTGTCACCTAAAT	X8	n/a	n/a	TGTGTGCAGTTTGAGAACATG	. 1	TTTCTTATATATACTACCCCTCC	400	401
X10 n/a n/a CAGTTAGTCTGCTGGAAAGC 1 TCCAAACCTACGCAATGTATC 300 301 pGEM-T GCATGGCGGCGCGGGGAATT GGCGGCCGCGGGAATTCACTAG T7T3 TTGTAAAACGACGGCCAGTG GGGAATTTGGCCCTCGAGG SP6 promoter primer TTCTATAGTGTCACCTAAAT	X9	n/a	n/a	AGGAACTGCGGTTGCCTTC	1	CCC CATABAACC	250	401
pGEM-T GCATGGCGGCGCGGGGAATT GGCGGCCGCGGGAATTCACTAG T7T3 TTGTAAAACGACGGCCAGTG GGGAATTTGGCCCTCGAGG SP6 promoter primer TTCTATAGTGTCACCTAAAT	X10	n/a	n/a	CAGTTAGTCTGCTGGAAAGC	, 1	TCCAACCTACCCAATATACC	309	201
T7T3 TTGTAAAACGACGGCCAGTG GGGAATTTGGCCCTCGAGG SP6 promoter primer TTCTATAGTGTCACCTAAAT	DGFM-T			GCATGGCGGCCGCGCGAAATT	··	GGCGGCCGCGA & TTCA CTAC	300	
SP6 promoter primer TTCTATAGTGTCACCTAAAT	T7T3			TTGTAAAACGACGGCCAGTG		GGGAATTTGGCCCTCGAGG		
	SP6 pro	moter prime	r	TTCTATAGTGTCACCTAAAT				
T7 Promoter primer TAATACGACTCACTATAGGG	T7 Prom	oter primer	•	TAATACGACTCACTATAGGG				

Strand +, ORF in same direction as genome; -, ORF on anti-strand; number in brackets indicate number of predicted exons if >1
Position of the first base of primer, relative to first base of the corresponding HHV-7 ORF (in the direction of genome); in the case of ORFs on the complementary strand, the forward primer is closer to the 3'-end of the ORF; for integenic probes, the start position is always 1

° Size of array probe in bp

.







Figure 3.2 DNA fragments of approximately 300 bp (small black rectangles) were PCR amplified from the 5'-end of each predicted ORF in the HHV-7 genome (block arrows); also for ten intergenic regions (small grey rectangles). The PCR products generated were processed for use as microarray probes to detect their corresponding sequences in order to measure the expression level of each predicted gene. The intergenic probes were included to detect any previously unidentified genes.



Figure 3.3 PCR amplification of HHV-7 fragments using specific primers (Table 3.1) and DNA extracted from HHV-7–infected SupT1 cells.

3.2.1.2 Cloning of array probes

Each amplified DNA segment was cloned into the bacterial vector pGEM-T by ligation. The pGEM-T plasmid was transformed into competent bacteria *Escherichia coli* (XL1-Blue) by heat shock. Bacteria were plated onto selective agar plates containing ampicillin, X-gal (5-Bromo-4-chloro-3-indolyl b-D-galactoside) and IPTG (Isopropyl b-D-1-thiogalactopyranoside) to identify clones with the viral DNA inserts (white colonies).

From each transformation, 5 random (or all if less than 5) white colonies were picked from the plate for PCR screening for the DNA insert. Universal primers (Table 3.1) that target the pGEM-T plasmid sequences flanking the insertion site were used to amplify the viral DNA insert. PCR products were analysed on agarose gels to check for the presence and approximate size of the inserts (Figure 3.4). The first round of cloning produced positive colonies (containing viral inserts) in 71 of 90 clones. Transformation was repeated for the remaining 19 clones, which produced satisfactory results, each giving at least 2 positive colonies. Finally, out of the 90 clones screened, all had at least two positive colonies apart from 8 clones (A12, B10, B11, C3, C5, D3, E11, F2) that had only one positive colony. 64 clones had the insert in all colonies tested. Therefore cloning was successful for all 90 probes (86 ORFs and 4 intergenic) (Table 3.2).





3.2.1.3 Sequence verification of clones

PCR screening of clones (section 3.2.1.2) confirmed the presence of DNA insert in all 90 clones but the use of universal primers was not able to confirm the sequences of the inserts. To ensure the clones contained the correct DNA probe, sequence verification was necessary, which was performed on plasmid DNA extracted from the expanded cultures for each clone. All 90 insert sequences were analysed by BLASTN 2.2.2 (Altschul et al., 2001) and were shown to match their corresponding ORFs (or intergenic region) in the HHV-7 RK genome (Megaw et al., 1998), as well as the JI genome (Nicholas, 1996), with adequate percentage identity scores (Table 3.2). All sequences matched with significant p-values, ranging from 1^{-15} to 1^{-128} . A few clones (C12, D4, E8 and G3) initially matched their respective ORFs (U39, U43, U57 and U79) with lower p-values of 1^{-2} to 1^{-7} , probably due to the short sequences (92 - 134 bases) generated and the quality of sequencing signal, resulting in a number of unidentified bases. A second attempt to sequence these clones produced better signals, higher p-values and identities (Table 3.2), confirming that the poorer blast match in the first sequencing attempt was due to sequencing errors, and the insert DNA in the clones were indeed of the correct sequences. Most sequences produced significant matches only to their corresponding region in HHV-7 sequences, with the exceptions A2 and H1, (HHV-7 DR6' and HHV-7 DR1, respectively) which generated many matches to sequences in other organism and cloning vectors. This was perhaps not surprising as these two clones represented the direct repeats at the end of the HHV-7 genome. This was taken into account in the microarray analyses of the four probes DR1, DR6, DR1' and DR6'. The sequencing results are shown in Appendix A.

		No.	with	%		Genome (RK)	%	
Clone	ORF	tested*	insert ^a	+veª	Blast match	location	identity ^b	p-value⁵
A1	HHV7DR1	5.0	4.0	80.0	HHV7DR1	4568-4475	93.8	1.0E-29
					HHV7DR1'	147614-147721	93.8	1.0E-29
A2*	HHV7DR6	2.0	2.0	100.0	HHV7DR6	6517-6813	94.8	8.1E-53
					HHV7DR6'	149563-149859	94.8	8.1E-53
A 3	HHV7U2	3.0	3.0	100.0	HHV7U2	11501-11634	93.6	1.0E-18
A4	HHV7U3	2.0	2.0	100.0	HHV7U3	12639-12950	95.9	3.4E-57
A5	HHV7U4	5.0	3.0	60.0	HHV7U4	14328-14578	88.7	7.8E-36
A6	HHV7U7	3.0	3.0	100.0	HHV7U7	17036-17323	99.3	2.0E-56
A7	HHV7U8	3.0	3.0	100.0	HHV7U8	18198-18397	78.6	5.4E-17
A8	HHV7U10	3.0	2.0	66.7	HHV7U10	18889-19151	84.4	1.9E-33
A9		3.0	2.0	66.7	HHV7U11	22076-22443	97.8	1.6E-72
A10		30	3.0	100.0	HHV7U12	22723-23008	100.0	7.7E-57
A11	HHV71113	3.0	3.0	100.0	HHV7U13	23754-24033	99.6	4.5E-55
A12		3.0	1.0	33.3	HHV7U14	24152-24456	86.8	5.3E-45
R12			2.0	100.0		26484-26737	95.8	8 0E-45
 		<u> </u>	<u> 2.0</u>	100.0		20404-20131	100.0	1 75 60
B2		5.0	5.0	100.0		2/00/-20020	100.0	0.45.00
B3	HHV7U18	5.0	5.0	100.0		29404-29/04	99.7	0.1E-00
84	HHV7U19	3.0	3.0	100.0	HHV/019	30872-31043	100.0	8.8E-30
B 5	HHV7U20	2.0	2.0	100.0	HHV7U20	32102-32378	97.9	1.7E-52
B6	HHV7U21	5.0	5.0	100.0	HHV7U21	33382-33706	99.7	3.4E-65
B7	HHV7U23	3.0	3.0	100.0	HHV7U23	34249-34574	100.0	7.2E-66
B8	HHV7U24	5.0	4.0	80.0	HHV7U24	34751-34920	100.0	1.2E-30
B9	HHV7U24A	3.0	3.0	100.0	HHV7U24A	34997-35129	93.4	2.4E-18
B10	HHV7U25	1.0	1.0	100.0	HHV7U25	35880-36107	88.9	7.5E-33
B11	HHV7U26	1.0	1.0	100.0	HHV7U26	36757-37062	98.4	2.1E-59
B12	HHV7U27	6.0	3.0	50.0	HHV7U27	37915-38152	92.8	1.4E-38
C1	HHV7U28	5.0	5.0	100.0	HHV7U28	40404-40677	99.6	3.0E-53
C2	HHV7U29	5.0	5.0	100.0	HHV7U29	41184-41479	100.0	4.3E-59
C3	HHV7U30	2.0	1.0	50.0	HHV7U30	41689-41990	98.7	2.3E-58
C4	HHV7U31	5.0	5.0	100.0	HHV7U31	44412-44679	99.3	5.3E-52
C5	HHV7U32	1.0	1.0	100.0	HHV7U32	51536-51711	98.9	2.9E-31
C6	HHV7U33	3.0	3.0	100.0	HHV7U33	51950-52194	95.1	9.3E-41
C7	HHV7U34	3.0	3.0	100.0	HHV7U34	52573-52927	99.7	5.7E-72
C8	HHV7U35	30	20	66.7	HHV7U35	53115-53335	95.1	0.05.39
C9	HHV71136	5.0	5.0	100.0	HHV71136	53354 53665	90.0	1 25 64
C10	HHV7U37	2.0	2.0	100.0	HHV71127	54912 55126	99.0	1.35-01
C11		2.0	2.0	100.0		54012-00130	100.0	1.25-00
C12**	HLN/71120	4.0	4.0	100.0	FIFTV7 036	00200-00020	99.7	1.0E-68
D1		4.0	4.0	100.0	HHV7039	60748-61031	100.0	5.3E-56
D2		4.0	4.0	100.0	HHV7U40	62895-63205	95.5	5.0E-56
D2		4.0	3.0	/5.0	HHV/U41	66235-66585	100.0	1.6E-71
D3	HHV/042	1.0	1.0	100.0	HHV7U42	68232-68520	100.0	1.6E-57
U4	HHV7U43	2.0	2.0	100.0	HHV7U43	71003-71231	99.6	4.4E-43
D5	HHV7U44	5.0	5.0	100.0	HHV7U44	71417-71709	100.0	2.0E-58
D6	HHV7U45	5.0	3.0	60.0	HHV7U45	72845-73120	98.9	3.2E-53
D7	HHV7U46	5.0	4.0	80.0	HHV7U46	73180-73400	99.5	1.1E-41
D8	HHV7U47	5.0	5.0	100.0	HHV7U47	74477-74785	93.2	8.0E-52
D9	HHV7U48	5.0	5.0	100.0	HHV7U48	76761-77073	97.2	1.9E-59
D10	HHV7U49	5.0	5.0	100.0	HHV7U49	77252-77598	97.7	4.8E-67
D11	HHV7U50	5.0	5.0	100.0	HHV7U50	77966-78398	99.8	1.2E-89
D12	HHV7U51	5.0	3.0	60.0	HHV7U51	79591-79828	72.4	7.3E-17
E1	HHV7U52	5.0	4.0	80.0	HHV7U52	80842-81148	92.9	1.1E-51
E2	HHV7U53	5.0	4.0	80.0	HHV7U53	81195-81487	99.7	5.2E-58
E3	HHV7U53.5	5.0	5.0	100.0	HHV7U53.5	82110-82344	95.5	1 0F_40
E4	HHV7U54	5.0	5.0	100.0	HHV7U54	83762-84099	 	7 25 50
E5	HHV7U55A	5.0	50	100.0	HHV71155A	85057_25362	00.3	6 75 64
		<u> </u>				00001-00000	50.7	0./ =-01

Table 3.2 Cloning and sequence verification of HHV-7 array probes

Cione	ORF	No. tested*	No. with insert ^a	% +ve*	Blast match	Genome (RK) location ^ь	% Identity ^b	p-value ^b
E6	HHV7U55B	5.0	5.0	100.0	HHV7U55B	86455-86754	100.0	5.3E-60
E7	HHV7U56	5.0	3.0	60.0	HHV7U56	87358-87676	98.1	1.3E-61
E8	HHV7U57	5.0	5.0	100.0	HHV7U57	91585-91696	98.2	3.0E-16
E9	HHV7U58	5.0	5.0	100.0	HHV7U58	91873-92109	75.9	2.3E-17
E10	HHV7U59	5.0	5.0	100.0	HHV7U59	94114-94391	95.0	1.1E-49
E11	HHV7U66	3.0	1.0	33.3	HHV7U66	99899-100167	74.4	3.1E-23
E12	HHV7U62	5.0	5.0	100.0	HHV7U62	96323-96424	99.0	7.1E-15
F1	HHV7U63	2.0	2.0	100.0	HHV7U63	96475-96748	91.3	8.9E-45
F2	HHV7U64	2.0	1.0	50.0	HHV7U64	97214-97465	99.6	1.0E-48
F3	HHV7U65	5.0	5.0	100.0	HHV7U65	98461-98727	86.8	2.2E-36
F4	HHV7U67	2.0	2.0	100.0	HHV7U67	100238-100507	80.8	3.8E-29
F5	HHV7U68	2.0	2.0	100.0	HHV7U68	101255-101557	100.0	1.1E-60
F6	HHV7U69	5.0	5.0	100.0	HHV7U69	101606-101911	97.7	1.8E-58
F7	HHV7U70	2.0	2.0	100.0	HHV7U70	103244-103546	97.4	1.6E-57
F8	HHV7U71	5.0	4.0	80.0	HHV7U71	104621-104822	100.0	7.3E-38
F9	HHV7U72	2.0	2.0	100.0	HHV7U72	105539-105905	100.0	3.9E-75
F10	HHV7U73	2.0	2.0	100.0	HHV7U73	105953-106251	100.0	9.0E-60
F11	HHV7U74	2.0	2.0	100.0	HHV7U74	108338-108584	93.1	7.6E-41
F12	HHV7U75	2.0	2.0	100.0	HHV7U75	110843-110684	97.5	1.9E-26
G1	HHV7U76	5.0	5.0	100.0	HHV7U76	112268-112599	96.4	1.7E-61
G2	HHV7U77	5.0	5.0	100.0	HHV7U77	112950-113253	99.0	7.3E-60
G3	HHV7U79	5.0	5.0	100.0	HHV7U79	117734-118055	100.0	1.7E-64
G4	HHV7U81	5.0	3.0	60.0	HHV7U81	119867-120164	100.0	1.5E-59
G5	HHV7U82	5.0	5.0	100.0	HHV7U82	120746-120968	90.2	2.9E-33
G6	HHV7U84	5.0	5.0	100.0	HHV7U84	121970-122251	96.2	1.7E-51
G7	HHV7U85	5.0	3.0	60.0	HHV7U85	122810-123140	98.8	3.1E-65
G8	HHV7U86	5.0	4.0	80.0	HHV7U86	126633-126930	100.0	4.5E-59
G9	HHV7U90	5.0	5.0	100.0	HHV7U90	133050-133371	97.5	1.0E-61
G10	HHV7U91	5.0	2.0	40.0	HHV7U91	133560-133833	99.1	2.5E-64
G11	HHV7U95	7.0	7.0	100.0	HHV7U95	137406-137672	70.4	1.1E-17
G12	HHV7U100	5.0	2.0	40.0	HHV7U100	142646-142948	79.6	8.7E-36
H1*	HHV7DR1'	5.0	3.0	60.0	HHV7DR1	4503-4790	96.2	1.0E-128
					HHV7DR1'	147549-147836	96.2	1.0E-128
H2	HHV7DR6'	5.0	3.0	60.0	HHV7DR6	6351-6828	99.7	3.8E-59
				·····	HHV7DR6'	149577-149874	99.7	3.8E-59
H3	X1 (U17 < U18)	5.0	5.0	100.0	U17< U18	28411-28683	100.0	6.7E-54
H4	X2 (U41 < U42)	n/a	n/a	n/a	n/a	n/a	n/a	n/a
H5	X3 (U46 < U47)	5.0	5.0	100.0	U46 < U47	73522-73822	100.0	1.9E-60
H6	X4 (U70 < U71)	n/a	n/a	n/a	n/a	n/a	n/a	n/a
H7	X5 (U77 < U79)	5.0	4.0	80.0	U77 < U79	115444-115682	100.0	1.0E-43
H8	X6 (U77 < U79)	n/a	n/a	n/a	n/a	n/a	n/a	n/a
H9	X7 (U86 > U90)	5.0	5.0	100.0	U86 > U90	127115-127459	99.4	2.1E-69
H10	X8 (U86 > U90)	n/a	n/a	n/a	n/a	n/a	n/a	n/a
H11	X9 (U91 < U95)	n/a	n/a	n/a	n/a	n/a	n/a	n/a
H12	X10 (U91 < U95)	n/a	n/a	n/a	n/a	n/a	n/a	n/a

a No. tested, number of colonies used in PCR screening for viral insert; No. with insert, number of colonies containing viral insert; % positive, percentage of colonies containing viral insert out of total number tested

b BLAST results: genome location, location of array probe sequence in the HHV-7 (RK) genome (Megaw et al, 1998); % identity and pvalue; calculated by BLAST

3.2.1.4 Amplification and purification of HHV-7 array probes

Using optimised procedures (section 3.2.2), the 90 HHV-7 probes for array spotting were PCR amplified from the sequence verified clones using universal pGEM-T primers (Table 3.1). Therefore the 96-well PCR plate contained 90 probes and 6 blank wells. The products were purified by filtration and resuspended in spotting buffer. All probes were successfully amplified and purified, as shown by agarose gel electrophoresis (Figure 3.5a). It can be seen that despite extensive PCR, purification and robotic liquid handling, the negative controls showed there was no apparent cross contamination. Probes U10 and U86 displayed a doublet, which could be due to cross contamination by another clone, as a common primer was used for all clones. PCR was therefore repeated for these two probes (Figure 3.5b)



Figure 3.5 a) HHV-7 array probes were re-amplified by PCR from sequence verified clones and purified by filtration. - , negative controls (reaction without DNA template) were included. b) Re-amplification was repeated for probes U10 and U86.



3.2.2 Optimisation of array probe amplification

To produce the in-house host-pathogen microarrays, it was necessary to amplify array probes for 90 HHV-7 genes and approximately 1000 human genes. Before this large-scale PCR amplification was carried out, the procedures for scaling up PCR reactions and the use of extensive robotic liquid handling were first optimised.

3.2.2.1 Optimisation of PCR reaction

12 different DNA polymerases from 8 suppliers (Table 3.3) were assessed for their suitability for amplifying array probes. The criteria for assessment were 1) robust amplification across a range of target concentration; 2) absence of additional PCR product bands; 3) ability to maintain activity after 3 hours at room temperature. Each enzyme was used to amplify a fragment of approximately 300 bp from template DNA of a dilution series from 10 ng to 1 fg, using the same cycling conditions except where specified by the manufacturer's instructions (section 2.5.1). Robotic handling was desired for the preparation of a large number of PCR reactions, but this could only take place at room temperature, hence the enzymes were tested for their ability to withstand room temperature for a prolonged period of time. To do this, two sets of reaction mixes were prepared for all enzymes. One set proceeded to PCR reaction immediately after preparation and the second set was left at room temperature for 3 hours before PCR was performed.

Results (Figure 3.6) showed that when used with our primers, templates, and experimental conditions, enzymes A, E and L produced inconsistent yield, enzymes C and K produced low yield and enzyme G gave no product at all. Enzymes B and C had the lowest sensitivity whereas enzymes F and J had the highest sensitivity and yield amongst those tested. However, these two high-yield enzymes also gave rise to smaller and fainter bands in addition to the product; and enzyme J produced inconsistent product size. In general enzymes H and I gave good yield and clean products. The 3-hour-wait at room temperature reduced the sensitivity of enzyme H, whereas enzyme I was not affected. Therefore enzyme I (Bioline Immolase) was chosen for the consistency, high yield, sensitivity and clean product. The fidelity of the enzymes was

not tested, as the sequences of the human clones were unknown. Any human genes found to show interesting expression patterns will be sequence verified subsequently.

Enzyme Label	Supplier	Name of enzyme
A	Qiagen	Taq DNA polymerase
В	Qiagen	HotStarTaq DNA polymerase
С	Novagen	NovaTaq DNA polymerase
D	Novagen	NovaTaq Hot Start DNA polymerase
E	Roche	Taq DNA polymerase
F	Roche	FastStart Taq DNA polymerase
G	Sigma	Taq DNA polymerase
н	Bioline	BIOTAQ DNA polymerase
I	Bioline	IMMOLASE DNA polymerase
J	Promega	Taq DNA polymerase
К	Finnzymes	DyNAzyme EXT DNA polymerase
L	Stratagene	YieldAce Hotstart DNA polymerase

Table 3.3 DNA polymerase batch test

3.2.2.2 Robotic liquid handling

A Biomeck 2000 liquid handler (Beckman Coulter, UK) was used to quickly and reproducibly prepare PCR reaction plates from master mixes of reagents; to aspirate small volume of PCR products for agarose gel electrophoresis; to select good PCR products, re-organise them onto clean plates and to purify these probes (section 3.2.3.1) for microarray printing.




Figure 3.6 DNA polymerase samples from different suppliers were tested for their suitability for use in array probe amplification PCR. Two sets of reaction mixes were prepared for each enzyme, according to manufacturer's guidance, using the same primers and diluted DNA template for all enzymes. One set was run immediately after preparation (left panel) and the second reaction mix left at room temperature for 3 hours before PCR was performed (right panel). a) Qiagen Taq DNA polymerase; b) Qiagen HotStarTaq DNA polymerase; c) Novagen NovaTaq DNA polymerase; d) Novagen NovaTaq Hot Start DNA polymerase; e) Roche Taq DNA polymerase; f) Roche FastStart Taq DNA polymerase; g) Sigma Taq DNA polymerase; h) Bioline BIOTAQ DNA polymerase; i) Bioline IMMOLASE DNA polymerase; j) Promega Taq DNA polymerase; k) Finnzymes DyNAzyme EXT DNA polymerase; I) Stratagene YieldAce Hotstart DNA polymerase. Template DNA dilution in ng; +, positive control; -, negative control (no DNA)

3.2.3 Production of HHV-7-human microarrays

3.2.3.1 Production of human gene probes

We created in-house host-pathogen arrays using the HHV-7 array probes described above (section 3.2.1) as well as human gene array probes. The human gene array probes were produced by PCR amplification from The I.M.A.G.E. Consortium Human Lymphochip subset [IRAY, IRBA, IRAZ, IRBB] (MRC gene service, UK), a set of lymphoid selected for their association with tissue rearrayed clones (http://www.geneservice.co.uk/products/image/image_subsets.jsp). This clone subset consisted of several libraries cloned in various vectors, of which the vector pT7T3-Pacl was used for the majority of clones. 1452 clones from the plates IRAY7-IRAY17, plates IRBA7-IRBA10, and plate IRBA12 were amplified using T7T3 Forward and Reverse primers (Table 3.1). PCR was successful for 907 clones (63%), giving good yield (approximately 10-20 µg per 100 µl reaction) and clean single bands on gel electrophoresis analysis of the PCR products. The remainder either failed to produce a product or resulted in multiple bands (Figure 3.7a). The 907 good products were selected for purification by filtration (Figure 3.7b), using a hit-picking protocol developed on the robotic system. Together this means that the PCR amplification, selection of good products and purification were carried out by optimised robotic liquid handling procedures (section 3.2.2). The identities of all 907 human gene array products are provided in Appendix B.



Figure 3.7a An example (plate IRBA7) of PCR amplified products from the Human Lymphochip subset. 68 out of 90 clones (75%, underlined) produced good, clean, single bands and were selected for purification and use as array probes.



Figure 3.7b Random sample of purified PCR products from the Human Lymphochip subset. Good PCR products were selected, purified by filtration, and analysed by agarose gel electrophoresis. For data tracking purposes, each purified probe were catalogued by their positions on the arraying plate as well as on the original clone plate.

3.2.3.2 Microarray printing

96 custom-designed HHV-7 probes (including 6 blanks) and 907 human probes (section 3.2.3.1) were printed in triplicate (287 spots of HHV-7 and 2721 spots of human probes) onto aminosilane-coated microarray slides by a 12-pin print head (3×4) using automated robotic deposition. Probes for HHV-6 and KSHV genes were also included on the array for the use of other projects. The arrays were laid out in 36 sub-grids (4×9), each of which was a square of 14 columns by 14 rows. In total the array consisted of 7056 spots, some of which were empty (Figure 3.8). The microarray printing was performed in collaboration with Edward Tsao, Royal Free & University College Medical School, London.



Figure 3.8 Format of HHV-7 microarray. Each array consisted of 36 sub-grids, with 196 spots (14 columns and 14 rows) per sub-grid. Each probe is spotted in triplicate, so that a probe in sub-grid 1, column 1, row 1 was also present in sub-grids 13 and 25, in the corresponding column and row.

3.2.4 Characteristics of HHV-7-human microarrays

The custom-made host-pathogen microarrays were hybridised in various combinations (Table 3.4) to test the characteristics of the HHV-7 probes, as well as for the performance of the whole array. 1) Two arrays (S14 and S20) were hybridised in reverse-dye format to assess dye bias and the effects of different batches of RNA preparation. 2) Three arrays (S02, S29 and S30) were hybridised to mock-infected samples to determine the cut-off point of data filtering. 3) Array S17 was hybridised to RNA from HHV-7–infected cells to check the sensitivity of the HHV-7 probes. 4) The specificity of the HHV-7 probes were evaluated by comparing arrays hybridised to virus-infected (S17) or mock-infected (S02, S29, S30) samples. 5) Reproducibility of arrays was assessed using technical and biological repeats (section 3.2.4.5). 6) The microarrays were also tested for their ability to produce distinguishing expression patterns for different samples by cluster analysis (section 3.2.4.6).

Array Slide	Cy5-labelled (red)	Cy3-labelled (green) Reference mix batch 2	
S14	Reference mix batch 1		
S20	Reference mix batch 2	Reference mix batch 1	
S02	RNA from mock-infected SupT1 cells	Reference mix batch 2	
S29	RNA from mock-infected SupT1 cells	Reference mix batch 2	
S30	RNA from mock-infected SupT1 cells	Reference mix batch 2	
S17	RNA from HHV-7 infected (72 hpi) SupT1 cells	Reference mix batch 1	

Table 3.4 Test hybridisations for the host-pathogen microarrays

The microarray experiment uses a comparative hybridisation approach. In this study we used the two-colour mode, measuring fluorescent intensities of samples labelled with two fluorescent dyes (red Cy5-dye and green Cy3-dye). PolyA-tailed mRNA samples to be analysed on microarrays were extracted from cells, amplified and reverse transcribed with red (Cy5) or green (Cy3) fluorescent-labelled dCTP into labelled-cDNA. The two species of labelled cDNA (Cy5 and Cy3) were co-hybridised onto each array. After washing to removal of unbound cDNA, arrays were scanned to measure the fluorescence intensities at 635 nm (Cy5) and 532 nm (Cy3) to generate an array image,

which was then analysed by image analysis software GenePix Pro 4.0. A matrix (the .gal file) defining the position and identity of all the array spots was overlaid onto the image, allowing the calculation of the signal and background intensities (both Cy5 and Cy3 channels) of each spot. These numerical data were used to generate Cy5/Cy3 Throughout this thesis arrays were hybridised to a Cy3-labelled common ratios. reference, so that the Cy3 signals acted as a constant control and the Cy5/Cy3 ratios were thus comparable between arrays. The common reference was a mixture of RNA extracted from 9 populations of cells, composed in the following proportion: U937 (25%), HeLa (25%), CEM-SS (15%), Ramos (15%), Huh7 (5%), MRC5 (5%), HHV7-SupT1 (5%), HHV6A(U1102)-SupT1, (2.5%) and HHV6B(Z29)-SupT1 (2.5%) cells. The purpose of using a mixture of RNA from different cell lines was to include many different RNA transcripts, so that more array probes will be bound and produce fluorescent signals, which in turn enables accurate Cy5/Cy3 ratios to be deduced for as many spots as possible. The inclusion of HHV-7 and HHV-6 RNA in the reference was to produce a signal for viral probes in the Cy3 channel, thus allowing the calculation of Cy5/Cy3 ratios.

3.2.4.1 Effects of Dye-bias and RNA preparation

The reference was made by first mixing the total RNAs in their corresponding proportions, followed by RNA amplification to give enough amplified RNA (aRNA) for many arrays. During the course of the thesis the first batch of reference exhausted hence a second batch was made using the same total RNA sources. It was therefore necessary to ensure there was no significance difference between the two batches of references. To assess the differences between the two batches of RNA reference mix, Ref-1 and Ref-2, they were each labelled in Cy5 and Cy3. Array S14 were hybridised to Cy5-labelled Ref-1 and Cy3-labelled Ref-2; the labelling was reversed for array S20 (Table 3.4).

If the two reference batches gave identical signals on the array, we would expect the median Cy5/Cy3 ratios of all spots on each array to be equal to 1 (or $\log_2 ratios = 0$) and the Pearson correlation values between the two channels should also be 1. Results show that good correlations were indeed obtained. For S14 (Cy5/Cy3 = Ref-1/Ref-2), median

 $log_2(Cy5/Cy3) = -0.417$; Pearson = 0.979 and for S20 (Cy5/Cy3 = Ref-2/Ref-1), median $log_2(Cy5/Cy3) = -0.068$; Pearson = 0.984 (Figure 3.9). Hence the median $log_2(Cy5/Cy3)$ value was close to zero in S20 and not so close in S14; whereas the Pearson correlation values were both close to and less than 1. To show that the high Pearson correlation values were indeed indicative of the high degree of similarity between Ref-1 and Ref-2 RNA batches, data from array S17 (different species of RNA in the two channels, see Table 3.4) gave a lower correlation of 0.82, indicative of more variation between the RNAs, as expected.

From this observation some conclusions can be made: 1) the Cy5- and Cy3-labelled species (Ref-1 and Ref-2) were highly correlated, as shown by the high Pearson correlation, hence there was negligible variation between the two batches of RNA references; 2) the deviation of the median $log_2(Cy5/Cy3)$ values from zero in S14 may be due to a bias towards Cy3, as a result of variations in labelling efficiency, hybridisation efficiency, or fluorescence detection, a phenomenon well documented for Cy3; 3) S20 has a median $log_2(Cy5/Cy3)$ value closer to zero hence is less affected by dye-bias than S14; 4) normalisation should be applied to remove dye-bias effect when arrays were used for HHV-7 expression profiling. In addition, the distribution of log_2 Cy5/Cy3 ratios around zero on the M/A plots (Figure 3.9) for S14 and S20 was an important observation for subsequent analyses. This was because many subsequent normalisation methods were based on this assumption.



Figure 3.9 Two batches of reference RNA mixture (Ref-1 and Ref-2) were each labelled in Cy5 and Cy3 and hybridised to each other in reverse-dye format. Figure shows m/a plots of two arrays, a) Slide S14 hybridised with Cy5-labelled Ref-1 and Cy3-labelled Ref-2; b) Slide S20 hybridised with Cy5-labelled Ref-2 and Cy3-labelled Ref-1. The log₂ (Cy5/Cy3) ratios of each spot is plotted against the geometric mean intensity of that spot. Pearson correlation values between the two labelled-species in each array are shown.

3.2.4.2 Data filtering and Signal-to-Noise Ratio threshold

Depending on factors such as transcript abundance and probe sequence, hybridised spots on an array gave signals of varied intensities. Some spots with very low intensities that were close to background level would result in inaccurate calculations and data interpretations. Therefore, a signal-to-noise ratio (SNR) filtering method was adopted to eliminate these 'noisy' data, as previously described (Jenner et al., 2001). Briefly, the SNR of each hybridised spot was determined by:

$$SNR = \frac{(Intensity_{total} - Intensity_{background})}{Intensity_{background}}$$

Spots with an SNR below a threshold value were considered not significantly different from background and were eliminated from analysis. This was a conservative approach and may remove some low intensity but authentic spots. SNR thresholds of 2.0 for Cy5 and 1.5 for Cy3 have been previously suggested to eliminate noisy data while retaining majority of 'good' data (Jenner et al., 2001). The determination of the SNR threshold value for data filtering in HHV-7–human microarrays was performed as follows.

Three microarrays (S02, S29, S30) were hybridized to Cy5-labelled RNA from mockinfected cells and Cy3-labelled reference RNA. Since HHV-7 RNA was absent in the mock-infected cells, any Cy5 fluorescence intensities detected by the HHV-7 probes were due to noise or unspecific hybridisation. The mean Cy5 intensity (of nine repeats – three slides with three spots per slide) of the 96 HHV-7 spots ranged between 60 and 250 fluorescence units (Figure 3.10a). In order to select the SNR threshold which achieves the maximum removal of noisy data while retaining as much 'good' data as possible, the number of probes that passed the SNR filtering in both channels were plotted against increasing values of SNR threshold (Figure 3.10b). The number of false positives in the Cy5 channel decreased rapidly with increasing SNR threshold, while the number of 'good' data points in the Cy3 channel showed a much slower drop. At the SNR of 1.5, all 6 negative control blank spots were removed in both Cy5 and Cy3 channels, hence was regarded as the lowest acceptable value for SNR filtering purposes. However, the number of noisy data points in the Cy5 channel was still high (44 of 90 non-blank spots, or 49.9%). The SNR threshold of 2.0 was the highest value at which the pass-rate in the Cy3 channel remained above 90% (85 out of 90, or 94%), while only 8 of 90 spots (8.9%) in the Cy5 channel were higher than the threshold (DR6, DR6', U7, U19, U36, U42, X5 and U79) (see specificity section 3.2.4.4). Therefore by raising the threshold from 1.5 to 2.0, the number of noisy data points in Cy5 was reduced by more than 5-fold, while the number of data points in Cy3 was only reduced by 4%, causing 3 additional spots (U49, U71, one blank) to fail the filtering.

The software used for data filtering (section 2.9) did not allow different threshold values to be set for the two channels, hence it was not possible to filter Cy5 and Cy3 data separately with respective SNR thresholds of 2.0 and 1.5 as described previously (Jenner et al., 2001). When SNR threshold of 2.0 was used instead of 1.5, the pass-rate in the Cy3 channel was reduced by 4%, but an extra 36 false positives in the Cy5 channel were removed. Therefore a SNR threshold of 2.0 was selected for filtering the HHV-7 microarray data, to obtain a lower false positive rate.



Figure 3.10 Three microarrays (S02, S29 and S30) were hybridised to Cy5-labelled RNA from mockinfected cells and Cy3-labelled reference RNA. a) Most of the Cy5 fluorescence intensities (average of 9 replicates, error bars represent standard deviation) detected by the HHV-7 probes (black dots) were below the SNR threshold of 2.0 (solid line); some of these were above the SNR threshold of 1.5 (dotted line). b) The number of probes passing SNR filtering decreased with increasing SNR threshold in both Cy5 (red) and Cy3 (green) channels. An SNR threshold of 1.5 or above (dotted line) was required to filter out all 6 blank spots. The SNR threshold of 2.0 (solid line) was selected to filter out maximum number of probes in Cy5 while retaining maximum number of probes in Cy3.

3.2.4.3 Sensitivity of HHV-7 probes

To assess the sensitivity of the HHV-7 array probes, RNA from an HHV-7-infected culture of SupT1 was Cy5-labelled and hybridised to slide S17, together with Cy3-labelled reference RNA. Cy5 fluorescence signals were detected by all HHV-7 probes (Figure 3.11). To determine the sensitivity of the HHV-7 probes, the SNRs of probes were determined and compared to the thresholds of 1.5 and 2.0 (section 3.2.4.2).

All HHV-7 probes passed the SNR threshold of 1.5, with the exception of two probes, U74 and X3, which had a SNR of 1.22 and 0.99, respectively. When the SNR threshold was raised to 2.0, one more probe (X7, SNR=1.71) also failed the SNR filtering. Out of the 90 probes, 88 probes (97.7 %) had SNR above 1.5; 87 probes (96.6 %) above SNR of 2; 62 probes (68.9 %) above SNR of 5; and 37 probes (41.1 %) above SNR of 10. Hence the majority of HHV-7 probes had high SNRs and the use of SNR threshold at 2.0 resulted in a high pass rate of 96.6%. Lowering the SNR threshold to 1.5 increased the pass rate by only 1.1 %. This finding further confirmed the use of 2.0 as the SNR threshold. Moreover, we have shown that majority of HHV-7 array probes were sensitive to HHV-7 RNA, giving good signal intensities when hybridised to RNA from infected cells of a late time point.



Figure 3.11 Microarray S34 was hybridised to Cy5-labelled RNA from HHV-7 infected cells (72 hpi) and Cy3-labelled reference RNA. Average fluorescence signals were calculated of triplicate spots (black dots). All except two probes (U74, X3) were above the signal-to-noise threshold of 1.5 (dotted line). Raising the SNR threshold to 2.0 (solid line) resulted in probe X7 also failing SNR filtering. Gene probes on graph in the same order as the corresponding ORFs in genome. 90 array probes were successfully amplified (black rectangles) and 6 did not amplify (grey rectangles). Guide lines (orange) are included to aid tracking the data points to their corresponding ORFs.

3.2.4.4 Specificity of microarrays

The specificities of the HHV-7 probes were evaluated by comparing arrays hybridised to Cy5-labelled virus-infected (S17) or mock-infected (S02, S29, S30) samples. The use of a common reference RNA in the Cy3 channel allowed comparison of the log_2 Cy5/Cy3 ratios between the different arrays. Figure 3.12 showed that all six blank spots (spotting buffer without DNA) had negative ratios in both mock- and HHV-7–infected slides. They also had low SNRs (average Cy5 SNR = 1.40 Cy3 = 1.39) hence were removed by the filtering criteria mentioned in section 3.2.4.2. This observation suggested two things: i) since all the blank spots failed the SNR filtering, the fluorescence in HHV-7 spots that have passed the SNR filtering must be due to real hybridisation signals rather than to auto-fluorescence in the spotting buffer; ii) the negative ratios of blank spots indicated a higher background fluorescence in the Cy3 channel than in the Cy5 channel, supporting the finding of a slight dye-bias towards Cy3 (section 3.2.4.1).

All HHV-7 probes had negative ratios on mock-infected slides; while all except two probes (X3 and X7) gave positive ratios on the HHV-7–infected slide (Figure 3.12). This difference clearly illustrated the ability of the HHV-7 probes to distinguish between mock- and HHV-7–infected samples. Moreover, majority of the HHV-7 spots passed the SNR filtering when hybridised to HHV-7–infected samples, but failed when hybridised to mock-infected samples, indicating these probes were specific to HHV-7 sequences. However, the specificity of 8 HHV-7 probes (DR6, U7, U19, U36, U42, X5, U79, DR6') were questioned, as they produced above-background signals when hybridised to mock-infected samples (striped green bars). Nonetheless, the SNRs of these 8 probes were only just above the SNR threshold of 2.0, which were very low compared to the true hybridisation signals against the HHV-7–infected samples (Table 3.5). However, two probes (U7 and U19) gave low SNRs for both mock- and HHV-7–infected samples. It was therefore more difficult to differentiate signal from noise for these two probes.

The sequences of these 8 probes gave no significant matches to human genes by BLAST searches. In fact, the probes only produced significant matches to their target positions in HHV-7 genome (Table 3.2), except U36 and U42, which also blasted to the corresponding conserved sequences in HHV6A and HHV6B, with lower similarities (62% and 66% similarity respectively) and lower E-values $(10^{-9} \text{ and } 10^{-13} \text{ respectively})$. Therefore the reason for these 8 probes passing the SNR filtering was more likely to be due to a non-perfect filtering method (i.e. false positives) rather than cross-hybridisation with cellular transcripts in the uninfected cells. These data showed the HHV-7 array had a high degree of specificity to HHV-7 RNA. Although all probes were analysed in future experiments, 11 of the probes (DR6, DR6', U7, U19, U36, U42, U74, U79, X3, X5 and X7) were considered with caution due to either high SNR in mock infected cells (section 3.2.4.2) or low SNR in sensitivity test (section 3.2.4.3).

Probe	SNR (mock-infected)	SNR (HHV-7-infected)	
DR6	3.49	67.65	
U7	2.18	2.81	
U19	3.70	2.61	
U36	2.21	40.74	
U42	2.95	44.38	
X5	2.85	212.64	
U79	2.25	23.89	
DR6'	3.14	35.66	

Table 3.5 Comparison of SNR for a subset of HHV-7 probes



Figure 3.12 Microarrays were hybridised to Cy5-labelled RNA from HHV-7 infected cells (red bars) or mock infected cells (green bars) and Cy3-labelled reference RNA. The average log2 Cy5/Cy3 ratios were calculated for data before (solid bars) and after (striped bars) signal-to-noise ratio filtering.

123

3.2.4.5 Reproducibility of microarrays

To access the reproducibility of the arrays, technical replicates were performed. Three slides were hybridised with the same sample (mock-infected SupT1 cells). S29 and S30 were identical repeats hybridised to the same labelled cDNA (product of one labelling reaction was split in half); whereas S02 was hybridised to cDNA from a separate labelling reaction that used the same RNA template as S29 and S30. Pearson correlation was calculated for the medians of (Cy5/Cy3) ratios of all spots (including viral and host). The Pearson correlation for unfiltered data of S29 and S30 were slightly higher at 0.90, compared to 0.88 and 0.85 for S02 vs S29 and S02 vs S30, respectively. After SNR filtering, the Pearson correlations were approximately 0.93 for all three comparisons (Figure 3.13a), showing the noise introduced by low intensity spots adversely affected inter-array correlation. These high correlations between technical replicates indicated the HHV-7–human arrays were very reproducible.

To further measure the reproducibility of the custom-made microarrays, the Pearson correlation was also determined for technical and biological replicate arrays hybridised to RNA from HHV-7 infected cells (Table 3.6). SNR filtering of data improved the correlation between replicates by 0.8 - 8.8 %, except in S32 vs S27, where a large improvement of 102.2 % was observed, probably due to large amount of noisy data, as seen in the exceptionally low correlation coefficient of 0.44 in unfiltered data. In S33 vs S13 the correlation coefficient in filtered data was actually lower than that in unfiltered data. To show that the correlation coefficients measured were indeed indicative of reproducibility of microarrays, correlation was also calculated for the arrays S34 (72 hpi) and S29 (mock-infected), where no correlation was observed, as expected. The average Pearson correlation coefficients of the technical repeats, whether mock or infected, were higher than that of the biological repeats in both unfiltered and filtered data (Figure 3.13b). The stronger correlation in technical repeats was expected, as a result of lower variation, since biological replicates would have been subjected to both biological and technical variations.

		Pearson correlation coefficient		
		Unfiltered	Filtered	Improve by
Mock technical		Ave 0.88	Ave 0.93	
S02 Mock	S29 Mock	0.88	0.93	5.6%
S02 Mock	S30 Mock	0.85	0.93	8.8%
S29 Mock	S30 Mock	0.90	0.93	3.5%
Infected technical		Ave 0.92	Ave 0.95	
S33 48hpi	S19 48hpi	0.96	0.97	0.8%
S34 72hpi	S17 72hpi	0.74	0.73	-1.6%
Infected biological		Ave 0.71	Ave 0.83	
S33 48hpi	S13 48hpi	0.87	0.94	7.2%
S34 72hpi	S28 72hpi	0.90	0.91	1.3%
S15 48hpi PAA S11 48hpi PAA		0.74	0.7 9	7.2%
S32 24hpi	S27 24hpi	0.44	0.89	102.2%
Infected vs I	Mock			
S34 72hpi	S29 Mock	-0.05	0.12	N/A

Table 3.6 Reproducibility of microarrays – Pearson correlation coefficients

In summary, the HHV-7-human microarrays produced highly correlated data amongst technical replicates (average correlation after filtering = 0.93 - 0.95) and biological replicates (average correlation post-filter = 0.83). These values were comparable with previously published data for both technical replicates: 0.96 (DeRisi et al., 1997), 0.91 (Ahn et al., 2002), 0.87 (Cohen 2000), 0.84 - 0.86 (Mayne et al., 2001), and biological replicates 0.97 (Scherf et al., 2000), 0.90 (Zhang et al., 2001b), 0.83 - 0.92 (Ross et al., 2000). Therefore the reproducibility of the HHV-7-human microarrays were satisfactory.



Figure 3.13 Pearson correlation coefficients of median (Cy5/Cy3) ratios between replicate slides before (white bars) and after (shaded bars) SNR filtering. a) Microarrays S29 and S30 were identical technical replicates hybridised to the same labelled cDNA in both Cy5 (mock-infected) and Cy3 (reference RNA); S02 was hybridised to cDNA of a different labelling reaction using the same RNA as S29 and S30. b) Pearson correlation coefficients of technical repeats of mock-infected (S02/S29/S30); technical repeats of HHV-7 infected (S33/S19, S34/S17); biologcial repeats of HHV-7 infected (S33/S13, S34/S28, S15/S11, S32/S27); and infected vs mock. Also see Table 3.6

3.2.4.6 Use of HHV-7-human microarray in gene expression profiling

The usability of the HHV-7-human microarrays in gene expression profiling of HHV-7-infection was tested. A selection of microarray data were analysed by clustering (section 2.10) to test whether the arrays were able to produce distinguishing patterns for different samples. Arrays were arranged into nodes in a tree according to the similarity of their patterns (Figure 3.14). Several findings were observed. Firstly, replicates were clustered together, indicating that similar expression patterns were detected by the arrays. Furthermore, technical replicates were more similar than biological replicates (S33 and S19; S07 and S13), consistent with results obtained in section 3.2.4.5. The arrays were even able to recognise identical repeats (S29 and S30) as more similar than non-identical ones (S02). Perhaps the most important observation was that the arrays were able to produce distinguishing patterns not only between mock- and HHV-7infected samples, but also between samples at different time point post-infection (8 hpi and 48 hpi). It can be seen that the variation between different time points was higher than the variation between biological replicates of a particular time point.



Figure 3.14 Tree structure of clustering analysis of selected microarrays.

Arrays with most similar patterns are clustered together. Samples with the same Roman numbers are technical replicates; different Roman numbers denote biological replicates

3.2.5 Data Normalisation

In order to ensure accurate measurement of biological-based differential gene expression in HHV-7 infected cells, microarray data were normalised (after filtering) to remove experimentally introduced variations, including different efficiencies in labelling, hybridisation, and effects introduced during scanning. Microarray data were subjected, in sequential order, to total intensity normalisation and LOWESS normalization using the Microarray Data Analysis System (MIDAS). MIDAS is part of a suite of programs known as TM4, developed at The Institute for Genomic Research (TIGR) (Saeed et al., 2003).

In this section, data from arrays S14 and S20 were shown as examples. Since these arrays were hybridised to reference RNA in both Cy5 and Cy3 channels, the signal in both channels should be equal, meaning the effectiveness of each normalisation step could be assessed by comparing real data to an idealised view of the data.

3.2.5.1 Effects of Total Intensity Normalisation

Due to differences in labelling efficiencies, most arrays in this study were systematically biased towards the Cy3 channel (section 3.2.4.1). Array S14 hybridised to reference RNA in both Cy5 and Cy3 channels showed a high Pearson correlation coefficient between the two dye channels (section 3.2.4.1) but also had a Cy3-bias, as illustrated by the negative $log_2(Cy5/Cy3)$ ratios (Figure 3.15a); and this bias was not limited to a specific area of the slide, but a common phenomenon across the whole slide (Figure 3.15b). In order to correct for this dye-bias, data were adjusted by a normalisation factor (section 2.9) so that the total intensities of all spots were equal in both channels. This resulted in an upward shift, where the $log_2(Cy5/Cy3)$ ratios of all the spots were increased proportionally to their original values (Figure 3.15c, 3.15d). Data from S20 were also subjected to total intensity normalisation, but the effect was not as noticeable because the two channels were already fairly balanced in raw data (Figure 3.16a-d).

3.2.5.2 Effects of Lowess Normalisation

Total intensity normalisation assumed that the variation in the $log_2(Cy5/Cy3)$ ratios of the data were independent of signal intensity. In reality, lower intensity spots tend to have $log_2(Cy5/Cy3)$ ratios that deviate more from zero (section 1.2.3.2), as shown in the raw data of array S20 (Figure 3.16a). This phenomenon could not be corrected by the use of a common normalisation factor in total intensity normalisation (Figure 3.16c).

Local weighted scatterplot smoothing (Lowess) normalisation was used to correct for intensity-dependent variation by applying a local weighted linear regression as a function of the intensity, with the Cy3 as reference point, as previously suggested (Quackenbush, 2002). The Lowess-adjusted data had a more balanced distribution of median $\log_2(Cy5/Cy3)$ ratios vs intensity (Figure 3.16e). The number of spots with $\log_2(Cy5/Cy3)$ ratios below -1.0 was 9 before normalisation and 0 after normalisation; For above +1.0, these numbers were 2 and 7, respectively. Therefore intensity-dependent variation was reduced by Lowess normalisation, preparing the data for more accurate analyses downstream.

Although one may argue that Lowess normalisation is an artificial adjustment of data, and may result in loss of definition in low abundance transcripts, it allows the correction of data points with low signal and falsely high ratios which may otherwise appear as highly differentially expressed genes. A previous study has shown that although the spread of data was reduced by Lowess normalisation, the ability to detect genes known to be differentially expressed was not affected (Yang et al., 2002b).



Figure 3.15 Microarray data of S14, hybridised to reference RNA labelled in Cy5 and Cy3, shown in M/A plots (a,c,e) and box plots (b,d,f). Raw data (a-b) were adjusted by total intensity normalisation (c-d); followed by Lowess normalisation (e-f).



Figure 3.16 Microarray data of S20, hybridised to reference RNA labelled in Cy5 and Cy3, shown in M/A plots (a,c,e) and box plots (b,d,f). Raw data (a-b) were adjusted by total intensity normalisation (c-d); followed by Lowess normalisation (e-f).

3.3 Summary

We have produced custom-made DNA microarrays containing probes to all 86 predicted ORFs and 4 intergenic regions of the HHV-7 genome, plus 907 probes targeting human genes. Validation tests showed that 1) the viral probes were sensitive and specific to HHV-7; 2) the high reproducibility of the arrays results were comparable to published data; 3) the arrays were able to distinguish between mock infected and HHV-7 infected samples, as well as samples infected for different periods of time; 4) the arrays were generally balanced in both channels but individual arrays may be prone to a bias towards Cy3 to varying extents.

This chapter also investigated the data adjustment steps required before useful interpretations could be made. It was found that the microarray data need to be filtered using a signal-to-noise threshold of 2.0 in both channels to remove low intensity spots; the filtered data need to be normalised by total intensity normalisation and Lowess normalisation to correct for dye-bias and intensity-dependent variation.

In summary, this chapter describes the production and validation of a host-pathogen microarray for HHV-7; and the optimisation of data filtering and normalisation procedures required for accurate analysis of the microarray data. The result is a microarray system that produces reliable data for the gene expression profiling of HHV-7 infection.

CHAPTER 4

INFECTIVITY OF HUMAN HERPESVIRUS TYPE-7 (HHV-7) IN SUPT1 CELLS

.

4.1 Introduction

In order to study the expression patterns of viral and host genes during human herpesvirus type-7 (HHV-7) infection, a time course infection of SupT1 cells by cell-free HHV-7 virus was required, and the transcription profiles of the infected cells analysed using the custom-made HHV-7 microarrays. To generate sufficient mRNA (200-500 ng) for each microarray analysis, 2×10^6 cells per infection are needed. Ideally a multiplicity of infection (MOI) of 1.0 or above is desirable to ensure majority (67 %) of the cells are infected, so that the expression patterns obtained are representative of HHV-7 infected cells rather than uninfected ones. Therefore this chapter is concerned with developing and optimising procedures for growing the HHV-7 virus in cell culture, producing high titre virus stocks, accurately quantifying the virus stocks, infecting SupT1 cells at a MOI of 1.0 or above, and using RNA amplification to facilitate microarray analysis.

4.1.1 Infectivity of HHV-7

HHV-7, like other betaherpesviruses (Shiraki et al., 1991), has long been known to be a low titre virus (Pellett, 2001), and have low infectivity-to-viral particle ratios (less than 1 infectious unit per 10⁴ particles) (Ablashi et al., 1998), which has hindered gene expression studies of the virus (Pellett, 2001). Such low ratios have been observed in human cytomegalovirus (HCMV), and are attributed to the production of noninfectious enveloped particles and dense bodies (Stinski et al., 1979), but it is unknown whether HHV-7 produce such particles. Direct pelleting by ultra-centrifugation resulted in higher viral particles/ml but not higher infectivity titre (Ablashi et al., 1998), probably because the virus was damaged during ultra-centrifugation. HHV-7 is a cell-associated virus, and infected cell lysates have been shown to be about 10 times more infectious than virus supernatant (Ablashi et al., 1998). However, in order to accurately monitor the changes in gene expression over time, a homogenous infection is required in this study. Therefore it is necessary to use cell-free virus and to remove the virus after the inoculation period. This constraint has limited the infectious titre used in this study. In fact, the low infectivity of HHV-7 has limited past attempts on gene expression studies (Pellett, 2001), therefore highlighting the importance of producing high titre stocks in this project.

4.1.2 Spinoculation and Polybrene

Different methods have been developed to improve the infectivity of viruses with low infectivity-to-particle ratios, including the use of spinoculation (also known as centrifugal infection) and the addition of cationic polymers such as polybrene. Spinoculation has been applied to successfully increase the infectivity of many viruses, for example, HIV-1 (Ho et al., 1993; O'Doherty et al., 2000; Pietroboni et al., 1989), herpes simplex virus (Tenser and Dunstan, 1980), human herpesvirus type-8 (HHV-8) (Inoue et al., 2003), cytomegaloviruses (Ho et al., 1993; Hodgkin et al., 1988; Hudson, 1988; Hudson et al., 1976), HHV-6 (Pietroboni et al., 1989), and human adenovirus (Nyberg-Hoffman et al., 1997). The enhancement effect of spinoculation on viral infectivity has been attributed to several factors that are believed to be increased by centrifugation, including cellular receptivity (Hudson, 1988; Hudson et al., 1976), viral fusion (Tenser and Dunstan, 1980), and viral deposition (Hodgkin et al., 1988; O'Doherty et al., 2000; Osborn and Walker, 1968), bringing viruses to closer contact with the cells. A study on influenza A virus had shown that reducing the volume of virus inoculum has the same effect as centrifugation (Reading et al., 2001) therefore favouring the close-contact hypothesis. The presence of polybrene (2-8 µg/ml) has been used to enhance the infectivity of HHV-8 (Inoue et al., 2003) and HIV (Pan et al., 1993), but had no effect on HHV-6 infectivity (Black et al., 1989). The addition of polybrene, together with an anionic polymer, has also been shown to enhance retrovirus infection of cells in a concentration-dependent manner (Landazuri and Le Doux, 2004), due to formation of virus-polymer complexes which aided sedimentation and delivered the viruses to the cells more rapidly than by simple diffusion.

4.1.3 RNA amplification

The application of the RNA amplification technique in gene expression analysis has been well documented (McClintick et al., 2003; Petalidis et al., 2003; Rajeevan et al.,

2003; Schindler et al., 2005), especially for detecting low copy number gene transcripts (Van Gelder et al., 1990), and for RNA extracted from limiting amounts of starting materials obtained by biopsy, laser capture microdissection (Luo et al., 1999) and flow cytometry-sorting. A number of different methods for RNA amplification have been published, which can be roughly divided into polymerase chain reaction (PCR)-based (Frohman et al., 1988) and in vitro transcription (IVT)-based approaches. Both methods essentially involve cDNA synthesis from the RNA by reverse transcription, followed by amplification of the cDNA. PCR based methods amplify the cDNAs using Taq polymerase. Due to the low fidelity of the enzyme and the exponential nature of the reaction, PCR-based methods have been largely replaced by the IVT-based approach, also known as the Eberwine method (Van Gelder et al., 1990). This approach involves the incorporation of a T7 promoter sequence in the synthesis of first strand cDNA using a specific cDNA synthesis primer, followed by second strand synthesis by DNA polymerase I. Antisense RNAs are then generated by IVT using T7 RNA polymerase. This method has the advantage over PCR-based approaches in that RNA is amplified in a linear rather than exponential manner. However, the aRNAs generated by the Eberwine method are anti-sense to the mRNA and 3'-biased, due to the use of oligo-dT primer in first strand synthesis (Nam et al., 2002). Therefore recent methods have made modifications to correct for this bias, such as the use of random primers (Goff et al., 2004), template switching (Petalidis et al., 2003; Puskas et al., 2002; Seth et al., 2003), and terminal continuation (Che and Ginsberg, 2004). Methods have also been developed to amplify sense strand RNA that are in the same orientation as the mRNA transcript (Goff et al., 2004; Marko et al., 2005; Rajeevan et al., 2003), which are suitable for use with oligonucleotide arrays and for array probes that require RNAs without a 3'-bias. RNA amplification has been shown to generate results that were reproducible within each method, although minor, method-specific systematic biases have been detected (Schindler et al., 2005; Wilson et al., 2004) when compared with unamplified RNA. Nonetheless, as long as the bias is consistent in all samples, such effects may be cancelled out in two-colour spotted microarrays, which measure relative expression ratios between two samples (Goff et al., 2004). Moreover, it has been shown that the correlation between aRNAs amplified by the T7-based method was good and the overall bias was low (Zhao et al., 2002), and the template switching method was able to preserve the known relative abundance of plant mRNA spiked into human mRNA samples at proportions between 0.1% to 0.00001% (Rajeevan et al., 2003).

RNA amplified using the SenseAmp kit (Genisphere, USA) has been shown to be reproducible, as well as having a high Pearson correlation coefficient of 0.90 when compared to unamplified RNA using quantitative real-time PCR (Goff et al., 2004).

4.1.4 Scope of chapter 4

The aim of this chapter was to establish a high MOI in SupT1 cells with HHV-7 cellfree virus, and to produce sufficient RNA for subsequent gene expression analysis. For this purpose the suitability of spinoculation, polybrene, and RNA amplification for HHV-7 infection of SupT1 cells was explored.

4.2 Results

4.2.1 Maintaining HHV-7 infection in SupT1 cells

HHV-7 (strain DC) infection of SupT1 cells *in vitro* was established in the laboratory. Infection was maintained by co-culture at a ratio of 1:1 to 1:2 (infected-to-uninfected cells). This culture ratio had to be carefully followed, because over-dilution with uninfected cells resulted in the loss of active infection. This was probably because HHV-7 has a slow infection profile hence the proliferating uninfected cells quickly outgrew the spread of infection. This finding agreed with the observation that HHV-7 grew best in media supplemented with a lower concentration (5%) of FBS (normally 10% is used for SupT1), which retarded the proliferation of uninfected cells (Ablashi et al., 1998). However, our results were contrary to the 1:5 co-culture ratio previously suggested by Secchiero *et al* (Secchiero *et al.*, 1997a), who used a different HHV-7 isolate (AL), which may have different growth kinetics. Some isolates have indeed been shown to be more cytopathic than others in SupT1 cells (Ablashi et al., 1998). The infection kinetics of the DC strain had not been determined in this study.

4.2.2 Morphology of infected cells

Infected cultures displayed cytopathic effect (CPE), characterised by the formation of giant multi-nucleated syncytia and attached cells - large, elongated, flattened cells that were loosely attached to the base of the culture container (Figure 4.1), consistent with previous studies (Ablashi et al., 1998). Flow cytometry analysis (FACS) of infected cultures revealed a population of larger and more granular cells, indicated by higher forward scatter (FSC) and side scatter (SSC) values, respectively (Figure 4.2e, h), in contrast to the small uninfected cells (Figure 4.2b). When comparing a cell-free virusinfected culture 4 days post inoculation to an ongoing infection maintained by coculture, the degree of CPE (proportion of large and granular cells) did not correlate with the proportion of cells bearing HHV-7 antigen as detected by FACS and immunofluorescent microscopy (Figure 4.2). The percentage of granular cells exceeded those bearing HHV-7 antigen. It may be possible that these cells displaying CPE were not infected, but experienced morphological changes as bystander effect. A similar cellular phenotype had been observed previously, where apparently uninfected cells in an infected culture underwent apoptosis (Secchiero et al., 1997a). As a result, visual judgement of CPE may be falsely high in the presence of cell debris in an ongoing culture, whereas a culture infected by cell-free virus may contain a portion of infected, HHV-7 antigen bearing cells that have no visible CPE. In summary, these finding demonstrated that although CPE may be crudely indicative of the degree of infection, it might not give accurate results as a direct measure of virus titre. This will be discussed further in the following sections.



Figure 4.1 Confocal microscopy photograph of unstained Sup-T1 cells. a) uninfected cells; b) cells treated with uninfected cell lysate; ; c-d) HHV-7 infected cells showing cytopathic effect: giant multinucleated syncytia (i); flattened, elongated cells loosely attached to bottom of cell culture flask (ii); cells that are both multinucleated and attached (iii), and cells with plasma membrane 'bubbles' (iv).



Figure 4.2 Flow cytometry and immunofluorescence of uninfected SupT1 (a-c), SupT1 infected with cell-free HHV-7 virus (d-f), and HHV-7 infected SupT1 cells maintained by co-culture (g-i). Percentage of large and granular cells (upper right quadrant) in dot plots (e,h) do not correlate with percentage of HHV-7 antigen-bearing cells (FL1-FITC greater than 60, dotted line in panels d,g). FSC, forward scatter; SSC, side scatter; red histogram, stained with anti-HHV-7 antibody; white histogram, unstained
4.2.3 Quantification of HHV-7 virus

In order to achieve HHV-7 infection with a high MOI, an accurate way of measuring infectious titre of the virus was required. Cell free stocks of HHV-7 were quantified using two approaches: determination of tissue culture infectious dose₅₀ (TCID₅₀) and infectious units count (IUC). For each of these approaches, two detection methods were used and compared. Detection by CPE or by indirect immunofluorescence assay (IFA) was used for the TCID₅₀ approach; while IFA or FACS was used for the IUC approach.

4.2.3.1 Quantification by Tissue Culture Infectious Dose₅₀ (TCID₅₀)

Briefly, TCID₅₀ is the viral dilution that results in a 50% chance of infection in target cultures (Peden, 1995) therefore is an end-point estimation of the infectivity. To determine TCID₅₀, 1×10⁵ SupT1 cells were incubated with serial dilutions of two HHV-7 stocks in a 24-well plate. Infection was detected by CPE observation or IFA. In CPE observation, a well containing one or more attached cell (hence HHV-7 infected) was counted as positive; whereas in IFA any well containing cells stained green by monoclonal antibody specific to an HHV-7 early antigen was scored positive. Interestingly, results of the TCID₅₀-CPE method were approximately one log higher than the TCID₅₀-IFA method for the same virus stock (Table 4.1), suggesting that either the CPE observation was a more sensitive method than IFA, or that it was affected by false positives. Unlike direct estimation (4.2.3.2), when CPE was used as end-point detection method, it was unlikely for the results to be affected by false positives, because attached cells were only present in HHV-7 infected culture and had previously been shown to contain virion particles capable of causing infection (Ablashi et al., 1998). Also, no attached cells were seen in SupT1 cultures mock-infected with freezethawed lysate of uninfected cells (Figure 4.1b). Therefore we believe that the $TCID_{50}$ -CPE method was a robust measure of virus infectious titre. Another possible reason for the different results might be because the CPE observation was made at 6 dpi, but cells were harvested for IFA at 4 dpi. Sensitivity of the TCID₅₀-IFA method may be improved by growing the cells for a longer period before staining to allow extra time for antigen production. However, it was unknown whether this would account for the difference between $TCID_{50}$ -CPE and $TCID_{50}$ -IFA results.

4.2.3.2 Quantification by Infectious units count (IUC)

In the second approach, IUC was calculated from the percentage of infected cells bearing HHV-7 antigen, which was detected either by IFA (Figure 4.3) or FACS (Figure 4.4) using the anti-HHV-7 antibody and the results translated into infectious units per ml (Table 4.1). The anti-HHV-7 antibody specifically bound to HHV-7 infected cells, but not to uninfected cells or cells infected by HHV-6, a close relative of HHV-7 (Figure 4.3). Therefore this IUC–IFA method had a low false positive rate and was considered a highly specific method for detecting HHV-7 infected cells. Results obtained were approximately 3 times higher than $TCID_{50}$ –CPE results (Table 4.1 & Figure 4.6).

IUC-FACS was the only method to show that stock 020723 had a higher titre than stock 040520 and the difference was about 20 times (Table 4.1). Note that only the major population of small cells (within gated region) were taken into account (Figure 4.4), whereas any HHV-7 infected cells in the large and granular population (outside gate) were excluded. The gating was necessary to remove false positives, because large and/or granular cells in HHV-7 infected cultures displayed high background fluorescence even in the absence of anti-HHV-7 antibodies (Figure 4.5). The drawback for this filtering was that the proportion of fluorescent cells in the gated region may be different from that in the whole cell population, which may account for the low estimate for stock 040520. Perhaps most of the cells infected by stock 040520 became larger and more granular, hence excluded by FACS gating, whereas the cells infected by stock 020723 remained small. Unfortunately, it was not possible to confirm this by FACS, due to the high background fluorescence of the large, granular cells (Figure 4.5). In contrast, the human eye was more accurate in differentiating the antigen-bearing cells, by their characteristic punctate staining pattern, from the background fluorescence (Figure 4.3). Of the two methods for infectious unit count, IUC-IFA was therefore considered more accurate than IUC-FACS.

HHV-7	TCID ₅₀	/ ml	Infectious units / ml		
stock	by CPE ^a	by IFA ^b	by FACS ^c	by IFA ^c	
020723	3.44E+04	1.81E+03	4.36E+05	8.44E+04	
040520	4.87E+04	7.24E+03	1.90E+04	1.33E+05	

Table 4.1 Results using different quantification methods of HHV-7 stocks

* TCID₅₀ value determined by scoring positive wells by the presence of attached cells

^b TCID₅₀ value determined by scoring positive wells by the presence of fluorescent cells

° percentage of HHV-7 antigen-bearing cells × number of cells/ volume of virus



Figure 4.3 IFA for HHV-7 infection. SupT1 cells uninfected (a-b); HHV-6 infected (c-d); HHV-7 stock 020723 infected (e); HHV-7 stock 040520 infected (f) were fixed and stained with anti-HHV-7 monoclonal antibody, followed by FITC to stain HHV-7 antigen-bearing cells (b,d,f,h); and DAPI counter-stain to visualise all cells (a,c,e,g)



Figure 4.4 Flow cytometry analysis of SupT1 uninfected (a,d,g); infected with HHV-7 stock 020723 (b,e,h); or stock 040520 (c,f,i). Results are representative of triplicate measurements. Infection was estimated using the percentage of HHV-7 antigen-bearing cells (FL1-FITC greater than 10^2 , dotted line) in the gated region (g-i). See also the legend for Figure 4.5



Figure 4.5 Flow cytometry analysis of uninfected (a,c,e) and HHV-7 infected SuptT1 cells (b,d,f). Infected cells (b) contained more larger (FSC) and granular (SSC) cells than uninfected cells (a). Without gating, infected cells displayed a 'tail' portion (d) of higher fluorescence (FL1 FITC) when stained with anti-HHV-7 antibody (red shaded), secondary isotype control (black curve), or unstained (green curve). Gating the population of smaller cells (black polygon) completely removed these tails in unstained and isotype controls, and reduced the tail in anti-HHV-7 stained cells (f). Gating has no effect on the fluorescence of uninfected cells (c,e)



Figure 4.6 Summary of Quantification methods. Results of different quantification methods on two HHV-7 stocks 020723 (white bars) and 040520 (shaded bars) are compared.

4.2.3.3 Summary of Quantification

In summary, several methods were used for HHV-7 quantification. IFA was highly specific for antigen positive cells but could not detect low level infection hence was not suitable for end point determination. It was nevertheless more specific than FACS when used for direct counting of infected cells. CPE was a sensitive method for endpoint determination using the cell attachment phenotype, but percentage CPE could not be directly translated into percentage of infected cells, due to bystander effect. (uninfected cells undergoing morphological changes) FACS offered a quick determination of infected population, but results were inconsistent with the other methods. Out of the four methods, the TCID₅₀-CPE end-point method and the IUC-IFA cell counting method gave most similar results. The results from these two methods had a reproducible difference of 2.5-fold (Figure 4.6 and Table 4.1) in both virus stocks tested. This systematic difference was probably because one method calculates TCID₅₀ value and the other measures infectious units. This also suggested that both methods produced reproducible results, therefore were believed to be the most accurate methods. Subsequent virus quantification experiments were therefore based on This thesis did not apply other methods of assessing viral these two approaches. infection, such as the detection of viral DNA by southern blotting, PCR or real-time PCR; viral proteins by ELISA; or viral particles by electron microscopy.

4.2.4 Optimisation of HHV-7 culturing to achieve a high multiplicity of infection

In order to achieve high multiplicity of infection for the gene expression studies, it was desirable to produce virus stocks of high infectious titre and to optimise the efficiency of infection. Having established accurate methods for HHV-7 titration, we were able to examine the effect of cell culture conditions on the titre of HHV-7 virus stocks.

4.2.4.1 Production of high tire virus

The first attempt to grow high titre HHV-7 virus was to expand the virus culture (in SupT1 cells) into multiple large volume (100 ml) flasks as described in section 2.3.4. When 15-30 flasks (1500 - 3000 ml) of HHV-7 infected cells (5 - 25% CPE) were harvested, the virus was released by freeze-thawing in $1/100^{\text{th}}$ volume as described in section 2.3.4. This meant $7.5 \times 10^7 - 7.5 \times 10^8$ CPE-displaying cells were concentrated to a volume of 15 - 30 ml, resulting in $5.0 \times 10^6 - 2.5 \times 10^7$ CPE-displaying cells per ml lysate (Table 4.2). This method has been used previously (Menegazzi et al., 1999), but the titres of virus stocks we obtained using this method were rather low (average titre = 1.9×10^3 infectious units/ml) (Table 4.2). In light of this, HHV-7–infected cultures were harvested in medium (300 ml) and small (6 - 10 ml) volumes, and their titres determined and compared (Table 4.2). Results showed that cells (40-75% CPE) grown in small volumes (6-9 ml) for freeze-thawing in 1-2 ml gave an average titre of 1.2×10^5 infectious units/ml (Table 4.2). This was almost 100 times higher than that obtained using the large volume method.

HHV-7 Culture volume (ml)	Cells/ml	% CPE	No. CPE- displaying cells ^a	Lysate volume (ml)	CPE/ml lysate ^b	Mean titre (infectious units/ml) ^c
1500-3000	1×10 ⁶	5-25%	7.5×10 ⁷ - 7.5×10 ⁸	15-30	5.0×10 ⁶ - 2.5×10 ⁷	1.9×10 ³
300	1×10 ⁶	25- 50%	7.5×10 ⁷ - 1.5×10 ⁸	30	2.5×10 ⁶ - 5.0×10 ⁶	2.0×10 ⁴
6-10	1×10 ⁶	40- 75%	2.4×10 ⁶ - 7.5×10 ⁶	1-2	2.4×10 ⁶ - 3.8×10 ⁶	1.2×10 ⁵

Table 4.2 Production of high titre virus stocks

* No. of CPE-displaying cells = culture volume × no. cells/ml × % CPE

^b CPE/ml lysate ≈ no. of CPE-displaying cells / lysate volume

^e For large volume (1500-3000ml), mean titre of 4 different stocks; for medium volume (300ml), mean titre of 2 different stocks; for small volume (6-10ml), mean titre of 7 different stocks; all calculated using the IUC-IFA quantification method

The reason for this was probably because it was more efficient for a culture in a small volume to reach high CPE percentage (40-75% at 6-8 days post co-culture); whereas cultures in large volumes often had lower CPE (5-25%), perhaps because a larger culture volume made cell-to-cell spread of viral infection more difficult. However, if the CPE estimation was correct, each ml of virus stock from a 1500 ml culture at 5% CPE would still contain more CPE-displaying cells than that from a 10 ml culture at 75% CPE (Table 4.2). This finding suggested that CPE of the culture was not directly related to number of infectious particles produced. HHV-7 was subsequently cultured in multiple (10-20) flasks of small volumes (6-10 ml) and harvested as 1-2ml stocks when the cells reached a CPE of 40-75%, as we have found that this provided stocks with the highest titre. The stocks were then pooled and quantified (section 2.3.5) before use.

There could be another possible reason for the unproportionally low degree of CPE in the large volume cultures. As the virus was passaged in SupT1 cells by co-culturing, the virus might have established latent infection in some cells, resulting in a loss of CPE and reduced production of infectious virus as passage number increased. Subsequent lysis of the cells and virus stock production therefore yielded less infectious virus per ml. In contrast, in the small volume culturing method, viruses were harvested as soon as the small culture reached an adequate CPE, thus avoiding the loss of productive infection with passaging.

4.2.4.2 Effect of Spinoculation & Polybrene on Inoculation efficiency

After optimisation of virus stock preparation, infectious titres of approximately 1×10^5 infectious units/ml were achieved (section 4.2.4.1). The next step was to improve the efficiency of HHV-7 virus inoculation, thereby increasing the effective MOI. Firstly, 1×10^5 SupT1 cells were inoculated with 100 µl of the virus stock 020723 for 1 hour at 37 °C. After the inoculation period, the virus supernatant was removed and used to inoculate fresh 1×10^5 SupT1 cells, in order to determine the amount of any unadsorbed virus remaining and if these were able to cause further infection. Fresh medium was added to the originally inoculated cells, and all cells were cultured at 37°C for 3 days, after which they were harvested, washed, and fixed onto slides for infectious unit count by IFA.

Results (Table 4.3) indicated that 1 ml of stock 020723 virus (estimated infectious titre $3-8\times10^4$ infectious units/ml. Table 4.1) was able to infect 6.8×10^4 cells after a 1-hour incubation period at 37°C, and the remaining virus supernatant was able to infect a further 1.4×10^5 cells, hence a total of 2.1×10^5 cells. This suggested the virus only achieved 33% of total possible infection after 1h incubation. This value coincided with the titre obtained for stock 020723, which was also determined after inoculation. However, this also showed that much infectious virus remained after 1 hour inoculation. Therefore methods to increase infection efficiency were investigated. Infection by spinoculation at 1000 g, 20 °C for 1 hour increased the number of infected cells by about 3.5-fold to 2.3×10^5 cells. Interestingly, the remaining virus only infected 1.6×10^4 more cells, bringing the total number of infected cells to 2.5×10^5 cells/ ml virus, similar to the total infection obtained without spinoculation. Therefore spinoculation for 1 hour achieved about 94% of total possible infection. If 2.5×10^5 cells/ ml virus was the maximum achievable infection of stock 020723, then the inoculation efficiency of 1hour incubation and 1-hour spinoculation was 27% and 94%, respectively (Table 4.3 & Figure 4.7a). These findings suggested that spinoculation was a more efficient inoculation method for HHV-7 infection of SupT1 cells.

The effect of polybrene on HHV-7 infection of SupT1 cells was also investigated. Results showed that the presence of polybrene (8 μ g/ml) during spinoculation led to a

decreased number of infected cells to 4.3×10^4 cells/ml virus, $1/5^{\text{th}}$ of the number infected when spinoculation was used alone (Table 4.3 & Figure 4.7a). When polybrene was used during incubation at 37 °C without spinoculation, no infected cells were detected at all. Therefore the presence of polybrene (8 µg/ml) had an inhibitory effect on HHV-7 infection of SupT1 cells with or without spinoculation.

	Number o	f infected cells			
	Original inoculum	Remaining supernatant	Total ^a	% Total ^b	% Maximum ^c
Incubation	6.8×10 ⁴	1.4×10 ⁵	2.1×10 ⁵	33	27
Spinoculation	2.3×10⁵	1.6×10⁴	2.5×10 ⁵	94	94
Incubation + polybrene	0	0	0	n/a	n/a
Spinoculation + polybrene	4.3×10 ⁴	1.4×10⁴	5.7×10 ⁴	75	17

Table 4.3 Effects of spinoculation and polybrene on HHV-7 stock 020723

* Total = infected cells/ml virus in original inoculum + infected cells/ml remaining virus in supernatant

^b % Total = infected cells/ml virus in original inoculum / total infected cells/ml virus achieved by the particular inoculation method × 100

6 % Maximum = infected cells/ml virus in original inoculum / total infected cells/ ml virus achieved by spinoculation × 100

In order to consolidate these findings, the experiment was repeated with several other virus stocks (Table 4.4). Again, spinoculation resulted in higher numbers of infected cells compared to incubation, with fold increases of 2.1, 2.5, 1.1, 3.4, and 3.8 for stocks 050118A, B, D, F and G, respectively, (Figure 4.7b). Except for stock 050118G, spinoculation also resulted in a higher percentage (91-100%) of total infection, compared to incubation alone (30-58%), as observed before in virus stock 020723 (Table 4.4). Moreover, for stocks 050118A-D, there were no detectable viruses in the remaining supernatant after spinoculation. Although the different stocks had varying infectious titres, spinoculation at 1000 g increased the infectious titres of all stocks (except stock 050118D) by 2.1- to 3.8-fold, clearly demonstrating that spinoculation was the more effective method. The enhancement effect of spinoculation on HHV-7 infectivity was consistent with previous studies in HIV (Ho et al., 1993; O'Doherty et al.,

2000; Pietroboni et al., 1989), HSV (Tenser and Dunstan, 1980), HHV-8 (Inoue et al., 2003), CMV (Ho et al., 1993; Hudson, 1988; Hudson et al., 1976) and HHV-6 (Pietroboni et al., 1989). However, our results showed that HHV-7 infectivity was only enhanced slightly, compared to 10-fold in HIV and 100-fold in HHV-6 (Pietroboni et al., 1989).

On the other hand, the presence of polybrene (8 μ g/ml) had a negative effect on HHV-7 infectivity with or without spinoculation (Table 4.3, 4.4), consistent with stock 020723 and published data on HHV-6 (Black et al., 1989), but not with previous studies on HHV-8 (Inoue et al., 2003) and HIV (Pan et al., 1993). Perhaps the presence of polybrene had adverse effects on SupT1 cells proliferation, therefore inhibiting secondary infection by cell-to-cell spread.

Therefore spinoculation in the absence of polybrene will be the method of choice for inoculating SupT1 cells with HHV-7 virus in subsequent experiments. From these data we have shown that, with the HHV-7 virus stock production method available (section 4.2.4.1), we can therefore achieve an MOI of 2.5 on 1×10^5 SupT1 cells, giving a theoretically synchronous infection.

	Number of infected cells / ml virus and % of total possi					ible infection	1	
	No Sr	<u>pin</u>	Spinoculate 1000g		Polybrene		<u> 1000g +</u> Polybrene	
Stock	Original inoculum	% Total ^a	Original inoculum	% Total ^a	Original inoculum	% Total ^a	Original inoculum	% Total ^a
020723	6.8×10 ⁴	33	2.3×10 ⁵	94	0	n/a	4.3×10 ⁴	75
050118A	1.5×10 ⁴	30	3.0×10 ⁵	100	0	n/a	0	n/a
050118B	1.6×10 ⁵	43	4.0×10 ⁵	100	0	n/a	0	n/a
050118D	1.5×10 ⁵	58	1.7×10 ⁵	100	0	n/a	0	n/a
050118F	9.3×10 ⁴	37	3.2×10⁵	91	0	n/a	0	n/a
050118G	6.7×10 ⁴	70	2.5×10 ⁵	55	0	n/a	1.6×10 ⁵	100

Table 4.4 Effects of spinoculation and polybrene on various HHV-7 virus stocks

* % Total = infected cells/ml virus in original inoculum / total infected cells/ml virus achieved by the particular inoculation method × 100. 100% means there were no detectable infectious virus left in the supernatant. For simplicity the number of cells infected by the remaining virus supernatant is not displayed. Also see notes for Table 4.3





Figure 4.7 Effect of spinoculation on HHV-7 infection. a) 10^5 Sup-T₁ cells were inoculated with 0.1 ml of virus stock 020723 for 1 hour either at 37°C (no spin), with spinoculation (1000g, 20°C), with polybrene (8 ug/ml), or with both spinoculation and polybrene. After inoculation period, leftover virus supernatant was removed and added to fresh Sup-T₁ cells (striped bars). Fresh medium was added to the originally inoculated cells (white bars). All cells were harvested after 3 days, and the number of infected cells determined by IFA with an anti-HHV-7 antibody. Results represent the mean of multiple measurements and the error bars represent the standard deviation. b) Experiment was repeated for 5 different HHV-7 stocks, only data for spinoculated cells (white bars) and incubated cells without spinoculation (shaded bars) are shown.

4.2.5 RNA amplification

4.2.5.1 Reasons for RNA amplification

Despite attempts (described in sections 4.2.3 and 4.2.4) to improve the infectious titre of HHV-7 virus stocks, the highest titres achieved were only 2-4×10⁵ infectious units/ml virus (with spinoculation), whereas at least 5×10^6 cells were needed to give enough RNA for each array hybridisation. When spinoculation was used on 5×10^6 cells (MOI of ~0.05), no CPE was visible at 48 hpi. When gene expression of these cells were analysed, the mean fluorescence signal and pass-rate for the HHV-7 probes in the Cy5 channel were low (Table 4.5), and the expression pattern probably represented the majority of uninfected cells rather than the infected cells. Given the limited virus titre, the only available option was to decrease the number of cells used in infection to 1×10^5 cells, so that a higher MOI of 2.5 (section 4.2.4.2) could be achieved. Since the amount of total RNA extracted from such a low cell number $(0.5 - 2.5 \mu g)$ were below the amount normally required for DNA microarray analysis $(50 - 200 \mu g)$, amplification of the RNA was required. Conventional methods produced 3'-biased, antisense amplified RNA (aRNA), ranging in size between 300-3000 nucleotides, with a peak at 600 nucleotides (Figure 4.9b). These 3'-biased aRNAs were too short to encompass some of the HHV-7 array probes, which were designed to target the 5'-end of HHV-7 ORFs (Figure 3.2). Therefore sense strand RNA amplification was needed to produce viral aRNA from the 5'-end.

4.2.5.2 Sense-strand amplification

Total RNA from 1×10^5 SupT1 cells infected by HHV-7 was amplified by the sense strand RNA amplification method (Figure 4.8, also see section 2.7.2.2). Briefly, first strand cDNA was synthesised from the 3'-end of mRNA by reverse transcription using a mixture of oligo-dT and random primers, in a reaction to favour full length cDNA synthesis. A poly-dT tail was added to the 3'-end of the cDNA, which was used for the annealing of a T7 primer. The 3'-end of the cDNA was then 'filled in' by Klenow fragment to generate a double stranded T7 promoter sequence. From this T7 promoter site, sense strand amplified RNA copies were produced by *in vitro* transcription using T7 RNA polymerase. This method produced amplified sense strand RNA (aRNA) of high yield, with an average of 5.6 μ g aRNA from 250 ng total RNA. Preliminary results from 12 mRNA purifications (using a column to bind polyA-tailed mRNA, section 2.7.1.3) suggested that total RNA in SupT1 cells was made up of, on average, 1.8% of mRNA, therefore sense strand RNA amplification using a combination of oligo-dT and random primers achieved on average a 1250-fold amplification of mRNA into aRNA. The yield was much higher than that attained when oligo-dT primer was used alone (i.e. no random primers), which only produced on average 2.8 μ g aRNA from 2 μ g total RNA (77-fold amplification). The aRNA products ranged from 100 to 4000 nucleotides in length, peaking at 200-500 nucleotides (Figure 4.9c), which was sufficient for recognition by the HHV-7 and human gene probes located at the 5'-end of transcripts.

Hybridisation of the fluorescent-labelled aRNAs (converted to cDNAs by reverse transcription) to the custom-made HHV-7 microarrays (see Chapter 3) produced satisfactory results. For example, an array (S19) was hybridised to Cy5-labelled aRNA (500 ng) from HHV-7 infected cells (48 hpi) and Cy3-labelled reference aRNA. The cDNAs bound to the array probes and gave good fluorescent signals, with 71.9% and 68.6% of the probes (viral and human) passing the signal-to-noise filtering criteria (see section 3.2.4.2) in the Cy5 and Cy3 channels, respectively (Table 4.5). All the HHV-7 probes had signal-to-noise ratios (SNR) above the filtering threshold, a pass rate of 100% in both channels, while the blank spots containing only buffer had low SNRs and had a pass rate of 0%. The Cy5 mean signal of the HHV-7 array probes was more than 5 times higher than the all-probes mean. It was unlikely for this difference to be due to a bias introduced by RNA amplification (Schindler et al., 2005; Wilson et al., 2004), because such difference was not observed in reference RNA, which was also amplified. Rather, the higher HHV-7 signals may reflect a large population of HHV-7 aRNAs in the Cy5-labelled 48 hpi sample. Moreover, the signal and pass-rate of HHV-7 probes were much (approximately 30 times) higher in the aRNA array than in the unamplified RNA array (Table 4.5), indicating that the aRNAs contain more copies of the HHV-7 transcripts. The aRNA and unamplified RNA were extracted from 1×10^5 and 5×10^6 infected SupT1 cells, respectively. Therefore the reduction of cells used in infection.

followed by sense strand RNA amplification (using a combination of oligo-dT and random primers), had improved the detection of HHV-7 transcripts.

		Cv5	<u>. </u>	Cy3			
Sample	Spot group	Mean signal intensity ^a	Mean SNR⁵	Mean signal intensity ^a	Mean SNR⁵	Cy5 SNR pass rate (%)	Cy3 SNR pass rate (%)
	HHV-7 blanks	74.0	1.21	388.6	1.46	0.0	0.0
Messenger RNA ^d	HHV-7 ORFs	115.5	1.90	816.4	3.13	10.2	100.0
	All probes ^c	383.9	4.85	1293.8	4.96	76.6	70.5
	HHV-7 blanks	56.9	0.93	194.3	0.73	0.0	0.0
Amplified RNA [°]	HHV-7 ORFs	3645.3	57.07	903.6	3.49	100.0	100.0
	All probes ^c	644.3	10.58	873.8	3.35	71.9	68.6

Table 4.5 Example arrays hybridised to messenger RNA or amplified RNA

* Mean signal intensity = mean of the background-subtracted signal intensities of all the spots in the particular spot group, in one channel

^b SNR = signal-to-noise ratio = (background-corrected signal)/background

° All spots containing array probes were included; but empty spots were excluded

^d Messenger RNA from 5×10⁶ HHV-7 infected cells (48hpi) and Reference RNA were converted into Cy5- and Cy3-labelled cDNA, respectively

 Messenger RNA from 1×10⁶ HHV-7 infected cells (48hpi) and Reference RNA were sense strand-amplified and then converted into Cy5- and Cy3-labelled cDNA, respectively



Figure 4.8. Sense-strand amplification of RNA (adapted from SenseAmp, Genisphere) Messenger RNA were amplified by: 1) first strand cDNA synthesis by reverse transcription using oligo-dT and random primers; 2) poly-dT tailing of cDNA by terminal deoxynucleotide transferase; 3) annealing of a T7 template oligo, followed by T7 promoter synthesis using the 'filling in' property of Klenow fragment; 4) *In vitro* transcription of sense strand aRNA from the T7 promoter by T7 RNA polymerase.



Figure 4.9 From 1 ×10⁵ SupT1 cells, total RNA and amplified RNA (aRNA) were analysed by Agilent Bioanalyser. a) total RNA; b) antisense aRNA; next page: c) sense strand aRNA amplified using oligo-dT and random primers; and d) sense strand aRNA amplified using oligo-dT primer only. The two ribosomal RNA species (18S and 28S) can be seen in total RNA (both graph and gel diagram), but absent in aRNA. [total RNA] and [aRNA] denotes the concentration. The sizes of aRNA products ranged between 100-4000 nucleotides.





Figure 4.9 (continued)

4.3 Summary

This chapter described the work needed to achieve a reproducible high multiplicity of infection (MOI) of HHV-7 virus in SupT1 cells. It was found that culturing HHV-7 virus in a small culture volume of 6 - 10 ml produced virus stocks of higher titre than in larger culture volumes. Virus infectivity was enhanced 2.1- to 3.8-fold by spinoculation at 1000 g for 1 hour at room temperature, but the presence of polybrene (8 µg/ml) had an inhibitory effect on HHV-7 infection. Further improvement of MOI was achieved using a smaller number of cells per infection, and subsequently using sense strand RNA amplification to obtain sufficient RNA for microarray analysis. As a result, amplified RNA from 1×10^5 SupT1 cells infected with HHV-7 at an apparent MOI of 2.5 was hybridised onto the microarray, which produced good signals, resulting in a 100% pass rate for HHV-7 probes, as compared to 10.2% for unamplified RNA from cells infected at a lower MOI. Therefore a system has been successfully developed for the HHV-7 infection of SupT1 cells that provides sufficient materials for microarray analysis.

CHAPTER 5

GENE EXPRESSION PROFILE OF HHV-7

5.1 Introduction

5.1.1 Classification of herpesvirus gene expression

The expression of herpesvirus genes is tightly regulated during infection (section 1.1.5) and was first classified into immediate early (IE or α), early (E or β) and late (L or γ) by pioneering work on HSV-1 infected cell polypeptides (ICP) in the 1970s, using inhibitors that block de novo protein synthesis (Honess and Roizman, 1974) and DNA replication (Honess and Watson, 1977). This classification approach is based on the following observations: 1) the expression of IE genes, which are mainly involved in transcriptional regulation of other viral genes, initiate as soon as the virus enters the host cell or reactivates from latency, and is independent of *de novo* protein synthesis; 2) E genes which encode enzymes and factors for DNA replication, are expressed in the presence of inhibitors of viral DNA replication, but are blocked by inhibitors of *de novo* protein synthesis; 3) expression of structural protein-encoding L genes require both protein synthesis and viral DNA replication, thus are repressed by both types of inhibitors. This approach has been used to classify genes of both human and animal herpesviruses, coupled with various detection methods such as northern blotting, reverse transcription-PCR and microarrays (Table 5.1). One disadvantage of this method, however, is that the effects of inhibitors on the cell may influence the viral transcription program, for example, the inhibitor for viral DNA replication, phosphonoacetic acid (PAA) (at or above 200 µg/ml) completely prevented growth of uninfected cells (Honess and Watson, 1977).

More recently, the use of microarrays for expression profiling without the use of metabolic inhibitors has been applied in the temporal mapping of genes in Kaposi sarcoma-associated herpesvirus (KSHV) (Jenner et al., 2001; Paulose-Murphy et al., 2001) and murine gammaherpesvirus 68 (MHV-68) (Ahn et al., 2002) (Table 5.1). This method relies on the pattern recognition ability of cluster analysis (Section 1.2.3.4) to group together viral genes that are co-regulated at certain stages of the infection. Transcription profiling of KSHV (Jenner et al., 2001) and MHV68 (Ahn et al., 2002) described that genes with similar functions were co-expressed. In KSHV, many DNA replication-related genes were upregulated between 10 - 24 hours post infection (hpi),

while structural genes were co-expressed between 48 – 72 hpi (Jenner et al., 2001). Microarrays had also been coupled with statistical tests to identify viral genes that were differentially up- or downregulated at different stages of the infection (Chambers et al., 1999). The transitions between the different stages of viral infection were sufficiently clear-cut to allow the distinction between genes involved in DNA replication and structural assembly, although some genes may sometimes be detected at the apparently 'wrong' stage of infection. Expression profiling not only allows the classification of genes, but also enables the expression to be monitored in a more authentic infection. In this chapter, the use of expression profiling together with inhibitor-based assays to assign HHV-7 genes into their temporal classes is described.

Virus *	Profiling ^b	Inhibitors	Detection	Reference ^c
Alphaherpes	sviruses			
HSV-1	N	Y	Radio-labeled protein gels	(Honess and Roizman, 1974)
HSV-2	N	Y	Northern blotting	(Easton and Clements, 1980)
VZV	N	Y	Protein gels	(Lopetegui et al., 1985)
BHV-1	N	Y	Northern blotting	(Seal et al., 1991)
EHV-3	N	Y	Northern biotting	(Sullivan et al., 1990)
Betaherpesv	/irus o s			
HCMV	N	Y	Northern blotting	(Wathen and Stinski, 1982)
HCMV	N	Y	Spotted microarrays	(Chambers et al., 1999)
HHV-6	N	Y	Reverse transcription-PCR	(Mirandola et al., 1998)
HHV-7	N	Y	Reverse transcription-PCR	(Menegazzi et al., 1999)
MCMV	N	Y	Northern blotting	(Keil et al., 1984)
Gammaherp	oesviruses			
KSHV	Y	N	Membrane arrays	(Jenner et al., 2001)
KSHV	Y	N	Spotted microarrays	(Paulose-Murphy et al., 2001)
KSHV	Y	N	Quantitative RT-PCR	(Fakhari and Dittmer, 2002)
MHV-68	Y	Y	Membrane arrays (Ahn et al., 2002)	
MHV-68	Y	Y	Spotted microarrays	(Ebrahimi et al., 2003)
Undefined s	ubfamily		-	· · · ·
CCV	N	Y	Northern blotting	(Huang and Hanson, 1998)

Table 5.1 Selected publications on temporal mapping of herpesvirus gene expression

* HSV-1, Herpes simplex type 1; HSV-2, Herpes simplex type 2; VZV, Varicella-zoster virus; BHV-1, Bovine herpesvirus 1; EHV-3, Equine herpesvirus type 3; HCMV, Human cytomegalovirus; HHV-6, Human herpesvirus type-6; HHV-7, Human herpesvirus type-7; MCMV, Murine cytomegalovirus; KSHV, Kaposi Sarcoma-associated herpesvirus; MHV-68, Murine gammaherpesvirus; CCV, Channel Catfish virus (Ictalurid herpesvirus 1)

^b Gene expression profiling involving the use of whole transcriptome microarrays or quantitative RT-PCR to profile the change in viral gene expression over time, other studies are qualitative detection of viral transcripts based on metabolic inhibitor assays c Not a comprehensive list

5.1.2 Temporal regulation of HHV-7 gene expression

The genome of human herpesvirus type-7 (HHV-7) has been sequenced and is predicted to encode 86 ORFs (Megaw et al., 1998; Nicholas, 1996). Functional annotations of these ORFs have been assigned mainly by homology to protein products in other herpesviruses (Table 1.3). Like all herpesviruses, HHV-7 is able to establish a persistent infection in host cells, achieved by the balance between latency and lytic replication (section 1.1.11). A model for studying latent infection of HHV-7 has not been developed, and its lytic replication has only been briefly investigated (Menegazzi et al., 1999). It is believed that lytic replication of HHV-7, like many herpesviruses, proceeds through a temporally regulated transcription programme of lytic gene expression (Roizman, 2001a). The timing of a gene's expression within this programme gives clues to its function (section 1.1.5). For example, genes that are expressed early include enzymes and factors required for DNA replication while those expressed later, after DNA replication has occurred, generally encode structural proteins of the virion.

There has only been one published study on the transcription of HHV-7 genes (Menegazzi et al., 1999), which analysed the expression of 15 ORFs during lytic replication in SupT1 cells using reverse transcription-PCR (RT-PCR). Of these, 10 ORFs (U10, U14, U18, U31, U39, U41, U42, U53, U73, U90) have been categorised as IE genes due to their expression in the presence of inhibitors of *de novo* protein synthesis (cycloheximide, CHX or emetine). U19 and U18/20 transcripts could be detected in the presence of PAA, which blocks viral DNA polymerase, but not in the presence of emetine, hence was classified as E genes. The unspliced mRNAs of U17 and U66 were IE transcripts, whereas their spliced versions were E and L transcripts, respectively. U100 was shown to have multiple splice sites, giving both IE and L transcripts when spliced and the unspliced mRNA showed E transcription kinetics.

Of HHV-7's 86 predicted genes, the expression of less than 20% has been analysed by only one study. Therefore, the expression pattern of many HHV-7 genes, and how this relates to the ordered series of events during lytic replication, remain largely unknown. The attention given to this virus is far less than the other members of the

betaherpesvirus subfamily, human herpesvirus type-6 (HHV-6) and human cytomegalovirus (HCMV). This is mostly due to the lack of confirmed disease association for HHV-7. Nevertheless, HHV-7 is an interesting candidate for study because of its low pathogenicity. This virus infects more than 90% of the world's population and persists in the host without apparent ill effects (section 1.1.8). We still do not know whether it really exists in a commensal symbiosis with human, or whether it is an opportunistic pathogen. HHV-7 reactivates HHV-6 from latency hence may contribute to complications in the immunocompromised (section 1.1.9). Alternatively, the virus may actually benefit the host in certain instances by competing for the CD4 receptor on T-cells with human immunodeficiency virus-1 (HIV-1) (Lusso et al., 1994), and downregulating surface CD4 to prevent super-infection by HIV-1 (section 1.1.10).

To gain more understanding about HHV-7 infection of T-cells, the viral lytic replication programme could be reconstructed by the large scale measurement of viral transcription using microarray technology (section 1.2.2), coupled with the pattern identification using cluster analysis (section 1.2.3.4). This chapter aims to use the custom-made HHV-7 microarray (section 3) to analyse HHV-7 gene expression during lytic replication in SupT1 cells.

5.2 Results

5.2.1 HHV-7 infection of SupT1 cells

Custom-made host-pathogen microarrays were used to monitor changes in the expression of HHV-7 genes during in vitro infection of SupT1 cells (a CD4⁺ T-cell line) over a 72-hour period. Cells were infected with cell-free HHV-7 virus at a multiplicity of infection (MOI) of approximately 2.5, with spinoculation at 1000 g. Cells were harvested at each time point: 8, 12, 16, 20, 24, 38, 48, 72 hours post infection (hpi) for RNA extraction and analysis by microarrays. The level of viral infection was confirmed by immunofluorescence assay (IFA) of selected samples using an anti-HHV-7 monoclonal antibody (Figure 5.1). Percentage of HHV-7-specific antigen-bearing cells increased with time, peaking at 48 hpi with approximately 80% of the cells infected (Figure 5.2b). The percentage of infected cells declined between 48 and 72 hpi. probably due to the proliferation of uninfected cells. The large percentage of infection was important for the microarray analysis, as this meant that the gene expression patterns detected would be representative of HHV-7 infected cells, as opposed to uninfected ones. The virus inoculum was removed after spinoculation, hence the infection should be synchronous (i.e. infection of all cells initiated within the first hour). However, it might be possible that virus removal was not 100% efficient and a smaller fraction of cells were infected at later times, causing asynchronous infection, which may explain the increase in the percentage of infected cells over time (up to 48 hpi). It was also possible that the infected cells began expressing HHV-7 antigen at different times. due to different cells being at different stages of infection. Nonetheless, the effects caused by this were minimal in the gene expresson analysis results (section 5.2.3), which measured the relative rather than absolute abundance of genes across time.



Figure 5.1 Immunofluorescence assay of HHV-7 infected Sup-T1 cells. Mock-infected (a, f); or HHV-7infected cells harvested at 8 hours post infection (b, g); 24 hpi (c, h); 48 hpi (d, i); 72 hpi (e, j) were fixed on slides and infection detected using primary antibody targeting HHV-7 antigen. Immunofluorescence was visualised using FITC-conjugated secondary antibody (f-j). DAPI counterstain showed total number of cells in each corresponding fields of view (a-e).



Figure 5.2 a) Number of HHV-7 probes passing the SNR filtering criteria during time course; b) percentage of infected cells at selected time points. Percentage of infected cells was monitored by immunofluorescence assay and was calculated as: (number of green fluorescent cells / total number of cells) × 100, results shown are the average of four fields of view per time point. Dotted lines indicate standard deviation.

5.2.2 HHV-7 gene transcription increased during infection

Signal intensities of the HHV-7 array spots measured at 8 hpi were low. Half of the HHV-7 transcripts were not detectable at this time point, as the signals were below the signal-to-noise ratio (SNR) filtering threshold. As the infection progressed, the number of detectable viral transcripts increased(Table 5.2 and Figure 5.2a). Such increase in the number of detectable viral transcripts had been observed previously (Jenner et al., 2001), and probably represented the gradual induction of viral genes during the progression of viral infection.

-	Time post- infection (hours)	Number (%) of HHV-7 probes passing SNR filtering ^a
•	8	49 (54.4%)
	12	80 (88.9%)
	16	83 (92.2%)
	20	82 (91.1%)
	24	77 (85.6%)
	38	77 (85.6%)
	48	88 (97.8%)
	72	77 (85.6%)

Table 5.2 Number of HHV-7 transcripts detected increased with time.

* Determined using SNR filtering criteria described in section 3.2.4.2

There was also a general trend of increasing abundance (fluorescent intensity) of HHV-7 transcripts over time (compare panels a-h in Figure 5.3), which coincided with the rising percentage of HHV-7 antigen-positive cells (Figure 5.2). The reason could be that as the number of infected cells at the same stage of lytic cycle expanded over time, more copies of the viral mRNAs were produced and passed the filtering threshold. However, although the intensities gave a general picture of the trend in viral transcription, intensities of individual genes cannot be directly compared between different samples. Comparisons between two-colour arrays need to be made using the Cy5/Cy3 ratios (explained in section 1.2.2), and will be described in later sections. However, the intensities of different genes can be compared within an array, showing that some gene probes had higher signals than others. The intensities of probes U27, U36, U42, U81, U90 and X5 were high relative to the other probes in almost every array; while signals of probes U53.5 and U56 were prominent in all arrays from 38hpi onwards. Of these, probes U36, U42 and X5 have been shown to be of slightly lower specificity, producing just above-background signals on arrays hybridised to mock-infected samples (section 3.2.4.2), which may explain their higher-than-average signals. The intense signals might also be the result of higher hybridisation efficiency of these array probes; or perhaps these genes were indeed expressed at higher levels. The HCMV homologue of U90 (HCMV UL122/3) and U27 (UL44) are known to be expressed in high abundance (Chambers et al., 1999; Geballe et al., 1986; Gibson, 1983; Stinski, 1978).



b) 20,000 15,000 X5 10,000 U42 U36 5,000 U27 0 HHV7_U28 HHV7_U35 HHV7_U4 HHV7_Blank HHV7_DR6 HHV7_U48 HHV7_Blan HHV7_Blan HHV7_U10 HHV7_U12 HHV7_U15 HHV7_U20 HHV7_U32 HHV7_U38 HHV7_U42 HHV7_U47 HHV7_U50 HHV7_U54 HHV7_U59 HHV7_U66 HHV7_U82 HHV7_X3 HHV7_U18 HHV7_U25 HHV7_U30 HHV7_U53 HHV7_U24 HHV7_U56 HHV7_U63 HHV7_U69 HHV7_U70 HHV7_U73 HHV7_U76 HHV7_U79 HHV7_U86 HHV7_U91

Figure 5.3 Cy5 signal intensities of HHV-7 array spots on microarrays hybridised to labelled cDNA from a) 8 hpi, b) 12 hpi, c) 16 hpi, d) 20 hpi, e) 24 hpi, f) 38 hpi, g) 48 hpi, h) 72 hpi. Data are the average of triplicate spots on each array; have been background-subtracted, filtered and subjected to total intensity normalisation and Lowess normalisation.



Figure 5.3 (cont.)





Figure 5.3 (cont.)



Figure 5.3 (cont.)
5.2.3 HHV-7 expression profile during lytic infection of SupT1 cells

After \log_2 transformation, normalisation and mean-centring (section 1.2.3) to enable cross-array comparison, microarray data were analysed by hierarchical clustering to arrange time points (from left to right) and genes (from top to bottom) according to similarity of the gene expression patterns (Figure 5.4 and 5.5). Two cluster analyses were performed, the first one (100%-presence) was with more stringent criteria, including only array spots that passed filtering (section 3.2.4.2) in all arrays (Figure 5.4). The result was therefore unaffected by missing data, although a fewer number of genes were included. The second cluster analysis (80%-presence) included genes with missing values in up to 1 out of 8 arrays. This may affect the stability of clustering pattern (de Brevern et al., 2004), but allowed more genes to be analysed (Figure 5.5).

5.2.3.1 Clustering (100%-presence) of HHV-7 genes

HHV-7 gene expression during a 72-hour time course infection of SupT1 cells were analysed by hierarchical clustering, which arranged the 8 time points into three main clusters from left to right (Figure 5.4): i) 8-12 hpi, ii) 16-24 hpi, and iii) 38-72 hpi. Within the 16-24 hpi cluster, the expression profiles at 16 and 20 hpi were found to be most similar to each other, whereas 24 hpi had the most distant profile. Profiles of 38 hpi and 48 hpi were more similar to each other than to that of 72 hpi. The ordering of the 8 time points on the cluster therefore reflected a gradual change in HHV-7 gene expression throughout the progression of infection. The largest shift in overall gene expression occurred between 24 to 38 hpi, and the second largest from 12 to 16 hpi, indicating major changes in viral gene expression between these times, which will be discussed in more detail later.

Hierarchical clustering also arranged the genes into three main clusters (I, II and III) from top to bottom, some of which can be viewed as two sub-clusters (IIA and IIB; IIIA and IIIB) (Figure 5.4). The mean gene expression profiles of the different clusters were also visualised in graphical form (Figure 5.6). Genes in the top cluster (I) mainly

showed decreased expression over time, followed by an increase between 48 and 72 hpi (Figure 5.6a). The middle cluster (II) was made up of two sub-clusters, the first of which (IIA) consisted of genes that showed early (8 - 12 hpi) increase in expression, followed by a decline (Figure 5.6b). Genes in second sub-cluster (IIB) had higher expression levels at 8hpi (Figure 5.6c), therefore were probably induced earlier in infection than those in IIA. Genes in cluster IIIA had a similar profile to IIA, except that the major rise in expression took place slightly later (between 12 and 16 hpi) and the decline after 24 hpi was not as rapid (Figure 5.6d). Finally, the bottom cluster (IIIB) consisted of genes whose expressions increased more slowly, and at a later stage (24 and 38 hpi) in infection (Figure 5.6e).

Although each gene probe was spotted in triplicate on the array, some genes with lower abundance only appeared once or twice on the cluster diagram, because one or two spots have failed the filtering criteria. However, the replicates were closely located in the cluster, indicating the replicate probes gave reproducible results, detecting similar expression profiles for the same genes.

Figure 5.4 (next page)

Cluster analysis of HHV-7 gene expression over a 72-hour infection of SupT1 cells (MOI=2.5). Only 80 viral gene spots (37 ORFs) with no missing data were included in analysis to give this 100%-presence cluster diagram. Time point samples (i.e. array slides) were clustered horizontally; viral genes were clustered vertically into three major cluster groups I, II and III. A value between -3.0 to +3.0 was calculated for each gene at each time point, depending on the relative level of gene expression. The values were presented on a colour scale (-3.0, green, downregulated; 0.0, black, unchanged; +3.0, red, upregulated). Gene names were colour-coded by their predicted function: green, transactivators; blue, DNA replication; orange, capsid & assembly; pink, tegument; red, envelope, grey, others; black, unknown.



Figure 5.4 Cluster analysis results (100%-presence)

5.2.3.2 Clustering (80%-presence) of HHV-7 genes

The clustering analyses in section 5.2.3.1 included only array data points that passed the data filtering criteria across all 8 arrays, therefore only 37 ORFs were included. In order to analyse the expression of more viral genes, the clustering parameters were amended so that gene spots that passed filtering in more than 80% of arrays (7 out of 8 arrays) could be included in clustering analysis (Figure 5.5). This adjustment allowed 31 additional genes to be clustered, therefore a total of 68 out of 90 (76%) HHV-7 probes were analysed by hierarchical clustering. The remaining 24% were excluded from clustering analysis due to low signals over the entire time course, which may be the result of probe design or low abundance transcripts.

The overall clustering pattern in this 80%-presence clustering (Figure 5.5a) was slightly different to that of the 100%-presence clustering (Figure 5.4), but the majority of genes were assigned to the same sub-clusters as before. The reason for this slight difference was probably because as the clusters were formed by the agglomerative average-linkage method, the inclusion of the extra 31 genes altered the new distances calculated between clusters (section 1.2.3.4), resulting in the sub-clusters being positioned differently. The expression profile at 12 hpi was now more similar to 16 hpi than to 8 hpi as suggested in the 100%-presence cluster. This indicated the gene expression pattern actually changed gradually from 8 through 12 to 16 hpi, unlike the major shift between 24 and 38 hpi, which was observed in both 100%- and 80%-presence clusters. More changes were observed in the arrangement of gene clusters. Cluster IIA was now more similar to clusters IIIA and I than to IIB; whereas cluster IIIB was the most distant. This illustrated the characteristic of the agglomerative nature of average-linkage hierarchical clustering (section 1.2.3.4). In light of this, it was important not to focus on the top-tobottom transitions, but to interpret both 100%- and 80%-presence clustering results in the form of the smaller and more robust sub-clusters. In the rare cases where genes were allocated to different sub-clusters by 100% and 80% clustering (for example U95), the assignment in the 100% cluster (cluster I) was considered to be more accurate.

Figure 5.5 (next page) a) Cluster analysis of HHV-7 gene expression over a 72h-infection of SupT1 cells (MOI=2.5). 168 viral gene spots (68 ORFs), with missing values (grey squares) in up to one time point sample were analysed to give this 80%-presence cluster diagram. Time point samples were clustered horizontally; viral genes were clustered vertically. Expression data at 0 hpi was not included in cluster analysis but presented on the side for comparison. b) Zoomed image of Cluster groups I, IIIA, IIA; c) Zoomed image of Cluster group IIB; d) Zoomed image of Cluster group IIIB. See also legend for Figure 5.4



Figure 5.5a Clustering results (80%-presence)



Figure 5.5b







Figure 5.5d



Figure 5.6 Mean expression ratios [log₂(Cy5/Cy3)] of 100%-prescence genes in a) cluster I; b) cluster IIA; c) cluster IIB; d) cluster IIIA; and e) cluster IIIB. Error bars (dotted lines) represent standard deviation.

5.2.3.3 HHV-7 genes of similar functions were co-expressed

HHV-7 genes were grouped into 4 functional classes by their predicted functions. These were transcription regulation, DNA replication & metabolism, DNA packaging & capsid formation, and virion assembly. These were colour-coded on the cluster diagram (Figure 5.4 & 5.5), together with genes of other functions that did not belong to these four groups, and genes of unknown function. Often genes with similar functions were clustered together. The majority of transcription regulators were assigned into cluster IIB; genes involved in DNA replication into clusters IIA and IIIA; structural genes for virion assembly into cluster IIIB. The most likely explanation for this clustering pattern was because viral genes with similar functions were involved in the same stage of the infection cycle, hence co-expressed at certain times during infection (Figure 5.7). This finding was consistent with the expression profiling of KSHV (Jenner et al., 2001; Paulose-Murphy et al., 2001) and MHV-68 (Ahn et al., 1999) using microarrays.



Figure 5.7 Mean expression ratio of HHV-7 genes by predicted functional group. The mean expression ratios of predicted transactivators (green), genes involved in DNA replication and metabolism (blue), in DNA packaging and capsid assembly (yellow), and envelope & structure (red) were taken from hierarchical clustering after mean-centring, and plotted over time. Error bars (dotted lines of corresponding colours) represent standard deviation. HHV-7 ORFs induced at various time points were colour coded and listed below the graph.

5.2.3.4 Immediate-early transactivators

Cluster IIB consisted of 15 genes, the expression of which were upregulated at the earlier stages of infection, and started to decline from 12 hpi onwards, as shown by the colour change in cluster diagrams (Figure 5.4 & 5.5) and the gene expression graph (Figure 5.7). Six of these (U44, U66, U84, U85, U86, U90 and U91) were detected at high levels at 8 hpi, indicating their induction were earlier than 8 hpi, meaning these were the earliest expressed genes detected in this time course. To determine the induction times for these genes, earlier time points would be required. It was possible that these genes may display IE expression kinetics, but confirmation would require protein synthesis inhibitor treatment (section 5.2.4). The abundance of two genes U86 and U90 were particularly prominent. Previous studies on HHV-7 have classified U90 (Menegazzi et al., 1999) as well as its HHV-6 (Mirandola et al., 1998) and HCMV homologues UL122/ie2 (Wathen and Stinski, 1982) as IE genes. No previous data are available on the expression of HHV-7 U86, but the expression of its HCMV homologue (UL123 or ie1) has been shown to be high at the immediate-early stage of infection, decreased during the early phase, and increased at later times (Stamminger et al., 1991; Stenberg et al., 1989). In HCMV, ie1 and ie2 are often called the major immediateearly (MIE) locus, encoding at least two products IE2 (the major regulatory protein) and IE1 (the accessory protein), which are responsible for mediating the activation of viral gene expression and regulating host gene expression (Spector, 1996; Stenberg, 1996). The MIE genes are expressed at relatively high levels compared to the genes of the other immediate-early (UL36-38) loci (Chambers et al., 1999), homologous to the IE-B (U17-U19) region in HHV-6 and HHV-7. Our results showed that this may also be true in HHV-7: U86 and U90 probes gave higher signals than U17 - U19 (Figure 5.3). Of the HHV-7 IE-B genes, U19 was assigned to cluster IIA (Figure 5.5a), whereas U18 was excluded from clustering analysis due to low signal. Interestingly, the expression profile of U17 was more similar to U86 and U90 than to U19, as it was assigned to cluster IIB in the 80%-presence cluster (Figure 5.5b).

Some transcription regulators were clustered elsewhere. For example, U95 was assigned to cluster I, showing a 'trough-shaped' expression profile, with high expression ratios at 8 hpi, which declined from 8 hpi to 48 hpi. This was followed by a

increase between 48 and 72 hpi (Figure 5.6a), possibly representing the beginning of a secondary infection. U42 was clustered with DNA replication genes, and U54 with the structural genes. These will be discussed in the relevant sections. Overall, the microarray-derived expression profile conformed to the known data on betaherpesvirus IE gene expression.

5.2.3.5 DNA replication and metabolism

The majority HHV-7 genes with predicted roles in DNA replication were found in clusters IIA and IIIA. These included DNA polymerase (U38), DNA polymerase processivity factor (U27), single-stranded DNA-binding protein (U41), and primase (U43). Helicase (U77), a second component of the heterotrimeric helicase-primase complex, was found in both clusters I and IIIB, whereas the third member, helicaseprimase associated factor (U74), was not included in clustering. In addition, several genes with nucleotide metabolism and repair functions, hence indirectly involved in DNA replication, were also clustered with the DNA replication genes. These included ribonucleotide reductase (U28), dUTPase (U45), phosphotransferase (U69), alkaline exonuclease (U70), uracil-DNA glycosylase (U81) and U55A, a gene homologous to HCMV UL84, of which the product was believed to facilitate HCMV oriLyt-dependent replication (Sarisky and Hayward, 1996). A gene expression transactivator (U42) was also clustered with the DNA replication-related genes. Its homologue in HCMV (UL69) may participate in DNA replication by upregulating the expression of DNA replication-related genes. The presence of HCMV UL69 product dramatically increases the transactivation of the TRS1 or IRS1 promoters (Romanowski and Shenk, 1997), which cooperate with IE1491aa and IE2579aa to transactivate E genes (Iskenderian et al., 1996; Romanowski and Shenk, 1997; Stasiak and Mocarski, 1992) of DNA replication functions (Pari et al., 1993). UL69 mutant virus growth was delayed under high MOI conditions due to an impact on DNA replication (Hayashi et al., 2000). We have found that HHV-7 U42 was highly expressed at the beginning of DNA replication stage (12 hpi), consistent with its probable function in transactivating enzymes and factors for viral DNA replication.

The expression of these DNA replication-related genes started low at 8hpi, then increased rapidly from 8 to 16 hpi (Figure 5.7), suggesting viral DNA replication

probably initiated during these 8 hours. These genes were subsequently downregulated between 24 and 38 hpi (Figure 5.7), suggesting that HHV-7 infection switched from a DNA replicating stage to a virion assembly stage between 24 and 38 hpi. Therefore expression profiling identified two major stages of HHV-7 infection cycle in human T-cells. After the reduction in expression between 24 and 38 hpi, some of these genes in cluster IIA (U19, U36, U42, U45 U69, U70) showed a second climb between 48 and 72 hpi (Figure 5.8b). This could be due to the beginning of a second round infection in new cells by the virions produced from the first round infection and may reflect asynchrony in the culture system after 48 hpi.

However, some DNA replication-related genes were assigned to other clusters. For example, only one each of the U43 and U70 array spots was assigned to cluster IIA, the other two were found in cluster IIB, a possible mis-assignment due to the missing values at 8 hpi, therefore the cluster IIA placement was probably more accurate. UL112/113, the HCMV homologue of HHV-7 U79, was suggested to have roles in DNA synthesis due to its ability to recruit replication-fork proteins to nuclear sites associated with viral transcription and DNA synthesis (Ahn et al., 1999; Penfold and Mocarski, 1997). However, hierarchical clustering assigned U79 to cluster I due to its 'trough-shaped' expression profile (Figure 5.6a). Although the functional significance of placement in cluster I was unclear, the relatively high abundance of U79 at 8hpi suggested that it was induced as early as the ORFs predicted as IE genes, and that it may have previously unknown functions in the immediate-early stages of infection. Another unexpected finding was that U55B had a very different transcription profile to U55A and was placed into cluster IIIB with the structural genes, despite the two genes were related (amino acid identity of 25%). The hybridisation signals on the microarrays were probably specific, as although the two ORFs were similar by amino acid sequence, their nucleotide sequences alignment matched no longer than 6 consecutive bases (Figure 5.9). Probes for U55A and U55B found no Blast match with the other ORF. Both HHV-7 U55A and U55B were suggested to have DNA replication-related functions, due to their homology to HCMV UL84, but our results showed that U55B may be involved in virion assembly stages, different to U55A. Perhaps the most surprising was that U73, believed to encode the origin binding protein, had an expression profile more similar to the structural genes in cluster IIIB than to the DNA

replication-related genes. However, the expression of U73 had two peaks (Figure 5.8a), suggesting second round of infection may have begun as early as 38 hpi.

5.2.3.6 Capsid assembly and Virion structure

Cluster IIIB was mainly made up of genes that were expressed at relatively low level during early stages of infection (8 - 24 hpi), followed by a major increase between 24 and 38 hpi (Figure 5.6e). Many of these genes are thought to encode structural proteins. These included genes involved in the formation of capsids, such as major capsid protein (U57), components of intercapsomeric triplex (U56, U76), and capsid protein located at the tip of hexons (U32); genes that encode virion envelope glycoproteins, gB (U39), gH (U82) and gM (U72); and other structural proteins such as glycoproteins (U23, U24, U100), large tegument protein (U31), virion proteins (U34, U50), antigenic tegument proteins (U11, U14), phosphoproteins (U10, U37), and a membrane protein (U46). In addition, genes involved in DNA packaging and capsid assembly, such as major capsid scaffold protein (U53.5), tegument proteins (U30, U64) were also assigned to this cluster. These findings suggested that HHV-7 DNA packaging and virion assembly initiated some time between 24 and 38 hpi during HHV-7-infection of SupT1 cells. The fact that genes encoding capsid, tegument and envelope proteins were clustered together indicated that proteins required for the formation of capsid, tegument and envelope were not expressed in three consecutive steps but simultaneously (Figure 5.7). The coexpression of all structural genes as one group has been observed in KSHV (Jenner et al., 2001).

Clustered with the structural genes was U54, predicted to encode a transactivator by homology to HCMV UL82. ppUL82 is an abundant tegument protein able to activate the ie1/ie2 promoter-enhancer in transient transfection assays (Liu and Stinski, 1992), but also has activities independent of ie1/ie2 expression (Baldick et al., 1997). UL82 expression starts early but peaks at late times in the infection (Gibson, 1983), consistent with our transcription profiling results on HHV-7 U54. HCMV UL82 is believed to act as a general transactivator of gene expression rather than as a specific activator of ie1/ie2 gene expression (Mocarski, 2001). It is unknown what particular genes are

activated by HCMV UL82, but our data suggested that HHV-7 U54 may be a transactivator of structural genes.



Figure 5.8 Mean expression ratio of a) HHV-7 U73 and b) selected cluster IIA genes during time course infection. Error bars (dotted lines) indicate standard deviation.

l えきある Chineshinesher of と目して payment book つけつつけ () いうてき

U55B Probe55B U55A Probe55A	GGGTTTGCGTCACTATGACCTTTTCGTTTTTGAATCTTATCAGACTAGGAAAAACTCCCA 	953 14 955 49
U55B Probe55B U55A Probe55A	TCTGCTCTTGTTCTTTAACATCATGGTTTGGGATGATGACGATAGAGCTGTTAATTTTTA TCTGCTCTTGTTCTTTAACATCATGGTTTGGGATGATGACGATAGAGCTGTTTAATTTTTA TTTTCTTTAAAGTAACAATTATCTTGTATCAG-CAGGGATGACGTAA-TGTTAAAAATTG TTTTCTTTTAAAGTAACAATTATCTTGTATCAGACAGGGATGACGTAA-TGTTAAAAATTG * * ** * * * * * * * * * * * * * * * *	1013 74 1013 108
U55B Probe55B U55A Probe55A	ACAACGGAAATCCAAAAAACAAGACTTTTGATGGATAAATGTGTCTATTCAGATTTTTTA ACAACGGAAATCCAAAAAACAAGACTTTTGATGGATAAATGTGTCTATTCAGATTTTTTA GCAGGGCAAAAATATTGACAAAGTTTATCGATTGTATCATAGGAAAAACATTA GCAGGGCAAAAATATTGACAAAGTTTATCGATTGTATCATAGGAAAAACATTA ** * **** * * * * *** * * **** *** ***	1073 134 1066 161
U55B Probe55B U55A Probe55A	AATGTAGTTTTAGTTGTTGTTCTTCCATTAAAGGACGAGCAGAAAAAAAGGAGTCAGAAT AATGTAGTTTTAGTTGTTGTTCTTCCATTAAAGGACGAGCAGAAAAAAAGGAGTCAGAAT ATTTTGATCACTTTTTTTTTTTTTTAGAACGTGACCCACCAAAGTTACGCATTCTGCAT ATTTTGATCACTTTTTTTTTTTTTTTTTAGAACGTGACCCACCAAAGTTACGCATTCTGCAT * * * * * * * * * * * * * * * * * * *	1133 194 1124 219
U55B Probe55B U55A Probe55A	TCCTTGAATTTATTTCAGCGCTCAAAAGTAAAACTGGTTCAGTTTTGATTGGACATACT TCCTTGAATTTATTTCAGCGCTCAAAAGTAAAACTGGTTCAGTTTTTGATTGGACATACT TGTTTTCAGTTCCATCGGTAATTAGCAACAGTACAGGTCTTTCGTGCGAAAGTATGCAGG TGTTTTCAGTTCCATCGGTAATTAGCAACAGTACAGGTCTTTCGTGCGAAAGTATGCAGG * ** * ** ** ** ** * * * * * * * * *	1193 254 1184 279
U55B Probe55B U55A Probe55A	CGTCAAATAATGTAACATTAT-CTTTACCCTGGAATTCTTCTTTGTATGGAATTCTA CGTCAAATAATGTAACATTAT-CTTTACCCTGGAATTCTTCTTTGA TAATCTCTGCGGGAGAAAAATTGA-GTTCTGGCTGCAAATCATTAAAATCCACAGTAAAC TAATCTCTGCGGGAGAAAAATTGAAGTTCTGGCTGCAA	1252 302 1243 317

Figure 5.9 Multiple sequence alignment by ClustalW, showing matching sequences between HHV-7 ORF U55A and array probe 55A (blue text); between ORF U55B and array probe 55B shown as (red text); between all four sequences (marked with *).

5.2.3.8 Classification of HHV-7 genes from clustering pattern

The 68 HHV-7 genes were assigned to classes A, B, or C, based on clustering results (Table 5.3). Thirteen genes in clusters I and IIB, which showed a decreasing relative expression over time, were assigned to class A. However, probes with missing values at 8hpi were probably mis-assigned, and were therefore manually placed into Class B, which consisted of 24 genes that had low expression levels at 8 hpi, and were upregulated during 12 - 24 hpi (clusters IIA and IIIA). Finally, 25 genes that were induced later in the time course (cluster IIIB) were placed into class C. Ten genes displayed expression profiles that could belong to more than one class, thus the classification of these await further analysis using metabolic inhibitors (section 5.2.4 - 5.2.5). Genes in classes A, B and C may be equivalent to IE, E and L genes, respectively, but confirmation of this would require the use of metabolic inhibitors to block *de novo* protein synthesis and DNA replication (section 5.2.4 - 5.2.5).

Our classification of HHV-7 genes was compared to previous temporal mapping studies of HHV-7, HHV-6 and HCMV genes (Table 5.3). Two of the class A genes in this study (U17, U90) have been previously analysed by RT-PCR with metabolic inhibitors and were reported to be IE-genes in unspliced form in HHV-7 (Menegazzi et al., 1999) and in HHV-6B (Mirandola et al., 1998). Of the class B genes, however, only one (U19) has been classified as and E-gene in both HHV-7 and HHV-6 (Menegazzi et al., 1999; Mirandola et al., 1998). U41 has been shown to be an IE-gene in HHV-7 (Menegazzi et al., 1999) and an E gene in HHV-6 (Mirandola et al., 1998); while U42 was classified as an IE-gene in both studies. U91 has been given both IE- and Ekinetics in unspliced and spliced form, respectively, in HHV-6A (Mirandola et al., 1998). Four genes (U14, U31, U39, U73) classified as class C in this thesis were shown to be IE-genes by Menegazzi et al. Our expression profiling results were also compared to a study of HCMV that used microarrays to detect transcripts in the presence of metabolic inhibitors (Chambers et al., 1999). Results were slightly more comparable, but not without discrepancies. Of the 24 class B genes in this study, the HCMV homologues of only six genes (U17, U38, U40, U41, U51, U81) have been classified as E-genes, while eight (U27, U42, U45, U55A, U66, U68, U69, U70) and two (U36, U62) were thought to be E-L and L genes, respectively (E-L genes were defined as late genes

with reduced expression, and L genes with undetectable expression in the absence of viral DNA replication). The class C genes were more consistent with the HCMV results, with the HCMV homologues of six (U11, U25, U31, U54, U64, U75) and four (U30, U56, U57, U72) HHV-7 genes also classified as L- and E-L genes, respectively. Nonetheless, five class C genes (U4, U14, U37, U39, U76) had E-kinetics in HCMV. Chambers *et al* only identified one IE gene (UL122) that has a counterpart in HHV-7 (U86), which was found in class A in our study. Another class A gene (U79) however, was considered to be an E/E-L gene in HCMV (Chambers et al., 1999).

These findings suggested that data obtained from microarray experiments are more comparable to each other than to RT-PCR studies. However, the reason for this may be simply because microarray studies generate more data to be compared. Our assignment of HHV-7 genes into classes A, B and C were based on the transcription profile of virus-infected cells at a high MOI of 2.5, in a relatively synchronous, natural infection *in vitro*; whereas treatment with metabolic inhibitors have damaging effects on the cells that may influence viral transcription (Honess and Watson, 1977).

5.2.3.9 Predicting functions for previously uncharacterised genes

Ten (U37, U44, U47, U58, U59, U62, U63, U68, U75, U84) of the HHV-7 genes analysed by clustering had no known function. The roles of these 10 genes could be predicted from the time at which they were induced (Figure 5.7), and from the clustering pattern (Figure 5.4, 5.5), because genes with similar functional roles were often grouped together by clustering analysis. U84 was induced before 8hpi and had similar expression pattern to the class A transactivators, therefore may itself be a transactivator involved in the immediate-early stages of infection. Several genes (U2, U44, U62, U68, U63) may have roles in viral DNA replication, as they were expressed at highest levels during 12 - 24 hpi and were clustered with DNA replication-related genes. Four ORFs (U59, U37, U63, U75) may encode for structural products or proteins involved in DNA packaging, inferred from the late induction (after 24 hpi) of their gene expression, which was typical amongst structural genes.

5.2.3.10 Summary of transcription profiling

In summary, monitoring HHV-7 gene expression by microarrays followed by transcription profiling using hierarchical clustering has revealed the regulated pattern of HHV-7 gene expression during infection of SupT1 cells. It has also allowed the classification of HHV-7 genes into classes A, B and C, based on their expression patterns, which also represented three stages of viral infection cycle. Figure 5.7 showed the proposed HHV-7 infectious cycle in SupT1 cells in vitro. After virus attachment and internalisation, expression of class A genes began within 8 hours, which include many genes believed to act as IE transactivators. Most genes involved in DNA replication and metabolism were found in class B. These were mostly induced between 8 and 16 hpi, and their expression started to decline after 24 hpi, to be taken over by the transcription of class C genes, which were involved in DNA packaging, capsid assembly and virion formation. The second round of DNA replication began between 48 and 72 hpi, suggesting the beginning of second round infection with the virions released from first round infection took place as early as 36 hpi (= 48 - 12) and by 60 hpi (= 72 - 12). Our results therefore suggested that the duration of infectious cycle of HHV-7 in SupT1 is between 36 to 60 hours, shorter than the 4-day cycle in cord blood lymphocytes previously suggested (Black et al., 1997). Analysis of infected cells at a time point between 48 and 72 hpi (e.g. 60 hpi) may facilitate the identification of a more precise duration.

Genes in classes A, B and C also roughly corresponded to genes previously classified as IE, E and L genes, respectively, in betaherpesviruses. Therefore expression profiling provided insight into the co-regulated expression of viral genes during different stages of its infection cycle in T-cells, and was able to classify genes into their temporal classes with results comparable with published findings.

5.2.4 Temporal classification of HHV-7 genes using metabolic inhibitors

To classify HHV-7 genes into kinetic classes of immediate early (IE), early (E) and late (L) genes, SupT1 cells were treated with metabolic inhibitors for 1 hour before infection by HHV-7. Cells treated with 200 μ g/ml cycloheximide (CHX) to block *de novo* protein synthesis were harvested at 8 hpi; cells treated with 500 μ g/ml phosphonoacetic acid (PAA) to block viral DNA replication were harvested at 24 and 48 hpi. The expression profiles of inhibitor-treated cells were analysed by the custom-made host-pathogen microarrays followed by clustering analysis (80% presence) in the same way as the untreated cells (section 5.2.3).

5.2.4.1 Immediate-early and Early genes

At 8 hpi, treating with CHX probably had little effect on viral gene expression patterns, as the expression profiles for both CHX-treated and untreated cells at 8hpi were highly similar and clustered together (Figure 5.10a). CHX was added to block protein synthesis in order to differentiate the IE genes that were independent of protein synthesis from the E and L genes that required de novo protein synthesis for their transcription. There could be two possible reasons for the lack of differentiation between CHX-treated and untreated cells at 8 hpi. First, the CHX might be ineffective in blocking protein synthesis in the HHV-7 infected SupT1 cells. Second, the expression of protein synthesis-dependent genes had not started or was not detectable by 8 hpi. The first reason was unlikely, because the presence of CHX indeed inhibited the expression of many proposed E and L genes (section 5.2.5). The second reason was probable, as it had been described in section 5.2.1 (Figure 5.4 - 5.5) that the majority of viral early transcripts only became detectable some time between 8 and 12 hpi. In order to prove this and to reveal the distinction between expression patterns in CHX-treated and untreated cells, a later time point for CHX-treated cells would be required. A 16 hpi sample has been attempted, but most cells have been killed in the presence of CHX by this time, therefore it was not possible to extract sufficient RNA for array analysis. In future experiments, a CHX-treated sample should be harvested at 12 hpi, to

determine if more genes differentially expressed between treated and untreated cells could be detected. Moreover, at 8hpi the abundances of many genes were below the signal-to-noise filtering threshold thus were excluded from analysis. Therefore we were unable to differentiate between IE and E genes using our microarrays. The lack of sensitivity for identifying IE genes by microarrays has been noticed in previous microarray study of HCMV (Chambers et al., 1999).

5.2.4.2 Early and Late genes

Since there were little difference between CHX-treated and untreated cells at 8 hpi, also 8 hpi samples contained a lot of missing data, clustering analysis was next performed without the 8 hpi samples, and focussed on the later samples, in order to optimise the number of genes analysed (Figure 5.10b). HHV-7 genes were classified into IE/E or L genes according to their response to PAA treatment. The expression levels of 15 genes were inhibited by PAA treatment (Figure 5.10b, 5.12), hence they were classified as viral DNA replication-dependent, or L genes. The differential expression of 8 of these L genes (U23, U46, U53.5, U54, U55B, U56, U57, U82) between treated and untreated cells were statistically significant, as determined by Significance analysis of microarrays (SAM) with false discovery rate set at 3.0%, or 0.54 gene (Figure 5.11). All remaining genes were classified as IE/E genes (Table 5.3), as their expression ratios were not reduced in the presence of PAA hence their transcription were independent of viral DNA replication (Figure 5.12).

Figure 5.10 (next page) a) Hierarchical clustering analysis of HHV-7 gene expression at 8, 24, 48 hours post infection of SupT1 cells in the presence and absence of metabolic inhibitors that block de novo protein synthesis (cycloheximide, CHX); and viral DNA replication (phosphonoacetic acid PAA). b) Hierarchical clustering analysis excluding 8 hpi samples; right panel, zoomed image of the cluster containing genes downregulated in the presence of PAA.

Figure 5.10a



202







b)



Figure 5.11 Significance analysis of microarrays (SAM) was used to identify statistically significant downregulated HHV-7 genes in the presence of phosphonoacetic acid (PAA) during infection of SupT1 cells. a) SAM scatter plot of observed (y-axis) vs expected (x-axis) expression values of genes significantly downregulated by PAA treatment (red dots) and unchanged genes (black dots). b) Genes significantly downregulated by PAA were clustered by hierarchical clustering.



Figure 5.12 Mean expression ratio of HHV-7 IE/E genes (blue dotted) and L genes (red dotted) differentiated by clustering analysis of their expression patterns in response to treatment of phosphonoacetic acid (PAA). Mean expression of genes significantly downregulated (solid red) by PAA and genes that showed no significant difference (solid blue), as determined by Significance Analysis of Microarrays (SAM) are also shown.

5.2.4.3 Comparison of expression profiling vs metabolic inhibitor assay

Of the 15 genes classified as late genes using PAA, 9 were assigned as class C genes by transcription profiling, and 4 genes (U10, U46, U50, U63) were found in an extra class in addition to class C (Table 5.3). The use of a DNA replication inhibitor therefore supported the suggestion that class C genes were exhibiting late gene kinetics. It also confirmed the class C assignment of the 4 ambiguous genes. However, two class A genes (U68, U86) were also shown to be inhibited by PAA, although the differences were not statistically significant. Therefore the distinction of L genes from IE/E genes using PAA treatment gave generally consistent results with expression profiling. The degree of consistency was improved by the application of SAM, which ensured a low false discovery rate. Nonetheless, the statistical stringency of PAA-treatment coupled with SAM may also have resulted in a high false negative rate, hence had a lower sensitivity in identifying L genes compared to the profiling method.

				This thesis		Published results				
		E Bradiated function	transcription	metabolic	HHV-7	HHV-6A	HHV-6B	HHV-6B	homologue	HCMV
	URF	Predicted function	profiling ^a	inhibitors ^b	RT-PCR °	RT-PCR ^d	RT-PCR ⁴	Array *	ORF ^f	Array ^g
	DR1	US22 gene family	A	IE/E				L		
	DR6	US22 gene family	A	IE/E				L		
	U2	US22 gene family	В					L	UL23	
	U3	US22 gene family							UL24	
	U4	related to U7 exon 2	С					L	UL27	E
	U7	US22 gene family, related to U4						L	UL27	E
	U8	US22 gene family		IE/E				L	UL29	L
	U10	nuclear phosphoprotein	C	L	IE				UL31	L
	U11	Structural phosphoprotein	C						UL32	L
	U12	G-protein coupled receptor, chemokine receptor		IE/E					UL33	E
	U13									
	U14	HCMV UL25/35 gene family, tegument phosphoprotein	C		IE			L	UL35	E
	U15		B/C	IE/E						
2	U17	IE-B, transactivator, US22 gene family	В	IE/E	IE (unspliced); E (spliced)	E	IE		UL36x1	E
07	U18	IE-B, homologue to HCMV IE protein			IE	E	E		UL37x3	
	U19	IE-B protein	В	IE/E	E	E	E	Ĺ	UL38	
	U20	probable membrane glycoprotein, Ig gene family?			E	E	E			
	U21	probable membrane glycoprotein		IE/E					ł	
	U23	probable membrane glycoprotein, EHV-1 gJ homologue	С	L						
	U24	Glycoprotein	С							
	U24A	contains hydrophobic domain								
	U25	US22 gene family, transactivator	С					L	UL43	L
	U26							L		
	U27	DNA polymerase processivity factor	В						UL44	E-L
	U28	Ribonucleotide reductase (large subunit)	В	IE/E					UL45	
	U29	Minor capsid protein (mCP)							UL46	E-L
	U30	Capsid assembly, myosin	С						UL47	E-L
	U31	Large tegument protein	С		IE	E	E	L	UL48	L
	U32	Capsid protein (located on tips of hexons)	С					L	UL48.5	
	U33	Virion protein		IE/E					UL49	E-L

		This thesis		Published results				HCMV	
OPE	Dradiated function	transcription	metabolic	HHV-7	HHV-6A	HHV-6B	HHV-6B	homologue	HCMV
URF		profiling *	inhibitors ^b	RT-PCR °	RT-PCR ^d	RT-PCR ^d	Array *	ORF ^f	Array ⁹
U34	Virion protein?	С						UL50	
U35	Role in DNA packaging	В	IE/E				L	UL51	
U36	Probable virion protein	В	IE/E				L	UL52	L
U37		С						UL53	E
U38	DNA polymerase	В	IE/E				L	UL54	E
U39	Envelope glycoprotein B (gB)	С	IE/E	IΕ	IE	IE	L	UL55	E
U40	Transport protein TP	В	IE/E					UL56	E
U41	Major DNA-binding protein,	В	IE/E	iE	E	E	L	UL57	E
U42	Transactivator	В	IE/E	ΙE	IE	IE	L	UL69	E-L
U43	Component of DNA helicase-primase complex, primase	В	IE/E				L	UL70	
U44		A	IE/E					UL71	
U45	dUTPase	В	IE/E				L	UL72	E-L
U46	Membrane protein	B/C	L				L	UL73	E-L
U47		A	IE/E					UL74	
U48	Envelope glycoprotein H (gH)						L	UL75	E-L
U49	Fusion protein							UL76	
U50	Virion protein	B/C	L			}		UL77	E
U51	G-protein coupled receptor	В	IE/E				L	UL78	E
U52	opioidR homologue						L	UL79	
U53	Protease/assembly protein			IE	E	E	L L	UL80	
U53.5	Major capsid scaffold protein	с	L						
U54	Tegument protein transactivator	с	L				L	UL82/83	L
U55A	Replication function?	В	IE/E				L L	UL84	E-L
U55B	Replication function?	С	L				L	UL84	
U56	Capsid protein	с	L				L	UL85	E-L
U57	Major capsid protein (MCP)	с	L				L	UL86	E-L
U58		в	IE/E				L	UL87	
U59		с	L				L	UL88	
U62		В						UL91	L
U63		B/C	L				L	UL92	L
U64	Role in DNA packaging, tegument protein	с	IE/E				L	UL93	L

		This t	hesis	Published re		ed results		HCMV	
ODE	Producted function	transcription	metabolic	HHV-7	HHV-6A	HHV-6B	HHV-6B	homologue	НСМ∨
URF	Fredicied function	profiling *	inhibitors ^b	RT-PCR °	RT-PCR ^d	RT-PCR ^d	Array *	ORF ^f	Array ⁹
U65	Tegument protein						L	UL94	L
U66	Late spliced gene (U60/U66) DNA packaging	В	IE/E	IE/L			L	UL89x2	E-L
U67			IE/E					UL95	
U68		В	L				L	UL96	E-L
U69	Phosphotransferase	В	IE/E				L	UL97	E-L
U70	Alkaline exonuclease	В	IE/E				L	UL98	E-L
U71	Myristylated tegument protein						1		
U72	Integral membrane protein (gM)	С					L	UL100	E-L
U73	Origin-binding protein (OBP)	C		IE	IE	IE	L		
U74	Component of DNA helicase/primase complex						L	UL102	
U75		C					L	UL103	L
U76	Virion protein?	С						UL104	E
U77	Component of DNA helicase/primase complex	С	IE/E				L	UL105	E
U79	HCMV replicaton, spliced (UL112/UL113)	A	IE/E				IE/E	UL112/113	E/E-L
U81	Uracil-DNA glycosylase	В	IE/E					UL114	E
U82	Envelope glycoprotein L (gL)	С	L .				IE/E	UL115	
U84	Spliced in HCMV	A	IE/E					UL117	
U85	probable membrane glycoprotein, related to OX-2	A							
U86	IE-A protein, HCMV IE2 homology	A			ļ]	IE/E	UL122	IE, L
U90	IE-A transactivator	A	IE/E	IE	IE	IE	IE/E		
U91	probable membrane glycoprotein	A	IE/E		IE (unspliced); E (spliced)	E (unspliced); IE (spliced)			
U95	US22 gene family,	A	IE/E				IE/E		
U100	Homology to HHV-6 gp82/105	С	L	E/L	L L	L	IE/E		
DR1'	US22 gene family	A							
DR6'	US22 gene family	A							
X1	•••								
X3									
X5		В							
X7		A	IE/E						

Table 5.3 Classification of HHV-7 genes in this thesis and previous studies

* Transcription profiling of HHV-7 genes based on clustering pattern of time course samples (8,12,16,20,24,38,48,72 hpi); class A, genes in clusters I and IIB; class B, genes in cluster IIA and IIIA; class C, genes in cluster IIB

•

^b Statistical analysis of infected in the presense or absense of metabolic inhibitors: cyclohexide (8 hpi) and phosphonoacetic acid (24,48 hpi)

° Temporal classification of HHV-7 genes using RT-PCR and metabolic inhibitors (Menegazzi et al., 1999)

^d Temporal classification of HHV-6A and HHV-6B genes using RT-PCR and metabolic inhibitors (Mirandola et al., 1998)

* Temporal classification of HHV-6B genes using arrays and metabolic inhibitors (Ohyashiki et a., 2005)

¹ HCMV homologues of HHV-7 (Nicholas, 1996; Megaw et al., 1998)

.

^a Temporal classification of HCMV genes using microarrays and metabolic inhibitors (Chambers et al., 1999)

•

5.2.5 Temporal classification of HHV-7 genes by RT-PCR

5.2.5.1 RT-PCR conditions

To verify microarray results, reverse transcription polymerase chain reaction (RT-PCR) was performed for a selection of ORFs. RT-PCR reactions were carried out using the OneStep kit (Qiagen, UK), which contained a mixture of reverse transcriptase and heat-activated HotStar DNA polymerase (Qiagen, UK). Firstly, RNA was reverse-transcribed into cDNA at 50 °C for 30 min, followed by heating at 95 °C for 15 min to inactivate the reverse transcriptase. The heating also activated the DNA polymerase for 35 cycles of PCR amplification reaction of the cDNA. However, it was found that the reverse transcriptase was not completely inactivated by heating, because if RNA was added after the heating step, PCR products were still formed (Figure 5.13). In contrast, using HotStar DNA polymerase (Qiagen, UK) alone produced no products with RNA samples. The HotStar DNA polymerase was capable of amplifying a product from viral DNA (Figure 5.13), but not from total RNA or amplified RNA (aRNA) in the absence of reverse transcription, indicating that there was no DNA contamination in the RNA samples. Some Taq-only reactions yielded products for some ORFs (Figure 5.14), although some of these bands (U17, U90) corresponded to spliced mRNAs therefore they were probably cDNAs from the 1st strand synthesis step in the RNA amplification reaction, rather than DNA contamination.

Initially 50 ng of aRNA was used as template, which resulted in strong product bands but failed to differentiate the expression of a predicted L gene (U56) between untreated or PAA-treated cells at 48 hpi; or the expression of a predicted IE gene (U17) between untreated and CHX-treated cells at 8 hpi (Figure 5.13). This could be either because PAA and CHX treatments made no difference to the expression of these genes, or because at this high template concentration, the transcripts were amplified to saturation, masking any differences. The latter was possible because if the inhibitors were not 100% efficient, transcripts that were supposedly inhibited by PAA or CHX may be present at a reduced level, as it has been observed in the array results (section 5.2.4). The aRNA has already been amplified approximately 1000fold by *in vitro* transcription (IVT) (section 4.2.5), so the RT-PCR was effectively the second round of amplification. Unlike IVT, PCR amplifies exponentially and as a result, the transcripts may have been amplified to saturation, masking the original differences in abundance. In an attempt to prove and rectify this, RT-PCR was performed using a serial dilution of template (from 50 ng to 0.5 pg for U56 in 48hpi samples; or 50 ng to 1 ng for U17 in 8hpi samples), to check if a difference could be observed between inhibitor-treated and untreated samples at lower template concentrations (Figure 5.13). The optimum template concentration for differentiating U56 transcript abundance between PAA-treated and untreated cells at 48hpi was found to be 500 pg/ul. This demonstrated that U56 was indeed an L gene, whose expression was reduced by PAA. For U17 expression at 8hpi, sensitivity to CHX was observed at the template concentration of 10 ng/ul. A template concentration of 1 ng/ul was estimated for 24 hpi samples. These template concentrations were used in subsequent RT-PCR reactions for 12 HHV-7 ORFs.

5.2.5.2 Classification of HHV-7 gene expression by RT-PCR

Expression of the U17 gene was not affected by PAA thus it was not a late gene. CHX treatment reduced the intensity of the two smaller bands and caused the largest band to disappear. The bands were gel purified and sequenced. It was found that the middle band corresponded to genomic sequence hence was an unspliced transcript, whereas the smallest band was a spliced transcript (Appendix C). RT-PCR results thus showed that both spliced and unspliced versions of U17 were reduced by CHX treatment, thus U17 was considered an E gene in both forms. This was slightly different to the previous RT-PCR study, which suggested U17 produced unspliced IE and spliced E transcripts (Menegazzi et al., 1999). The large U17 band produced mixed species when sequenced. U90 expression was not affected by CHX or PAA thus was most likely expressed as an IE gene. The transcription product was smaller than the original probe, which was revealed to be a spliced product by sequence verification (Appendix C). This agreed with previous data (Menegazzi et al., 1999), although Menegazzi *et al* also detected an unspliced form of U90 as an IE transcript.

RT-PCR results showed U27, U39 and U41 to be IE genes, with expressions only slightly reduced by CHX and unaffected by PAA. This was consistent with previous RT-PCR study (Table 5.4) but not with expression profiling using microarray in this study (Table 5.3), which classified U27 and U41 to be class B genes, and U39 to be a class C gene, even though low signals were detected at 8hpi. Both U27 and U41 are thought to be involved in DNA replication. U27 is the DNA polymerase processivity factor and U41 the single-stranded DNA binding protein. From their functions one would predict E kinetics for their expression. However, our data suggested that some low level expression occurred during the IE stage of infection without the need for transactivators, but their expressions were highly accelerated after the IE products had accumulated. The alternative explanation was that CHX was not 100% effective in blocking *de novo* protein synthesis, therefore a low level 'leaky' expression took place.

U66 transcript was most abundant at 24hpi, with or without PAA treatment. It was also detected in untreated cells at 8hpi, but was inhibited by CHX. By 48 hpi the abundance had decreased, therefore U66 displayed a classic E gene profile. This was consistent with microarray classification (Table 5.3) but not with previous results using RT-PCR (Table 5.4). U38, which encodes the viral DNA polymerase, also displayed patterns indicative of E gene (Figure 5.14), consistent with microarray results but no previous RT-PCR data is available for comparison.

Expression of ORFs U10, U50, U53.3 U56 and U82 were all reduced by PAA treatment, suggesting L-kinetics (Figure 5.14). This was consistent with microarray data (Table 5.3). Where replicates of the genes U10 and U50 were present in more than one cluster, RT-PCR results helped towards the final decision of their classification as L genes.


Figure 5.13 RT-PCR using specific primers for HHV-7 ORF a) U82, b) U56, c) U17; template: HHV-7 DNA as positive control; water as negative control; amplified RNA (aRNA) or total RNA (T-RNA) purified from HHV-7 infected SupT1 cells at a) 48hpi, b) 48hpi, c) 8hpi. Reactions were performed with either reverse transcriptase (RT), heat-inactivated reverse transcriptase (hRT), or Taq DNA polymerase (Taq). b-c) template aRNA was serial diluted, and were purified from cells treated with cycloheximide (CHX); phosphonoacetic acid (PAA); or untreated (--).



Figure 5.14 Agarose gel electrophoresis of RT-PCR products a-b) RT-PCR of aRNA from HHV-7 infected SupT1 cells using specific HHV-7 primers; c) Taq only, control PCR reaction without reverse transcription. RNA template: C8, CHX-treated 8hpi; N8, untreated 8hpi; P24, PAA-treated 24hpi; N24, untreated 24hpi; P48, PAA-treated 48hpi; N48, untreated 48hpi; DNA, HHV-7 DNA control; the 500bp fragment of DNA ladder contains 25ng DNA per lane.

Table 5.4 Temporal classification by RT-PCR results compared with microarray results and previous publication

		RT-PCR (this thesis) *						Microarrays (this thesis)			
ORF	Predicted function	8h	8h	24h	24h	48h	48h	Kinetic	Transcription	PAA treatment	inhibitors ^e
		СНХ	ND	PAA	ND	PAA	ND	Class ^b	profiling °	& microarray ^d	in indicated
U17	IE-B, transactivator, US22 family	+	+++	+++	+++			E	В	IE/E	IE/E
U27	DNA polymerase processivity factor	++	+++	+++	+++			IE	В	IE/E	
U39	Envelope glycoprotein B (gB)	++	+++	n/a	n/a	+++	+++	IE	С	IE/E	IE
U41	Major DNA-binding protein,	++	+++	n/a	n/a	++	+++	IE	В	IE/E	IE
U90	IE-A transactivator	++	+++	+++	+++			IE	A	IE/E	IE
U38	DNA polymerase	-	-	n/a	n/a	++	+++	Е	В	IE/E	
U66	DNA packaging,	-	++	+++	+++	+	+	Е	В	IE/E	IE/L
U10	nuclear phosphoprotein	-	+	n/a	n/a	-	+	L	С	L	IE
U56	Capsid protein,	++	+++	n/a	n/a	+	++	L	С	L	
U50	Virion protein	n/a	n/a	n/a	n/a	+	+++	L	B/C	L	
U53.5	Major capsid scaffold protein	n/a	n/a	n/a	n/a	+	+++	L	С	L	
U82	Envelope glycoprotein L (gL)	n/a	n/a	n/a	n/a	•	+	L	C	L	

* Arbitrary scale of band intensity on agarose gel: +++, strong; ++, medium; +, weak; - negative; n/a, not tested; CHX, cycloheximide (200 µg/ml); PAA, phosphonoacetic acid (500 µg/ml)

*Kinetic class assigned based on RT-PCR results: IE, immediate-early gene unaffected by CHX; E, early gene inhibited by CHX but not PAA; L, late gene inhibited by PAA

* Transcription profiling of HHV-7 gene expression using microarrays without metabolic inhibitors

^d Microarray analysis of HHV-7 gene expression in the presence of PAA

• Menegazzi et al., 1999

5.2.6 Host gene expression

The expression of cellular genes were analysed during HHV-7 infection of SupT1 in vitro using the custom-made microarrays. Clustering analysis showed that most host genes remained relatively unchanged throughout the time course (Figure 5.15). Any changes in cellular gene expression were of small magnitude relative to the mock infected cells (less than two-fold). Two clusters of genes can be tentatively identified (Figure 5.15). However, because of the small dynamic changes in gene expression, the significance of these genes was difficult to determine. Therefore to robustly identify differentially regulated genes, Significance Analysis of Microarrays (SAM) was used to compare cellular gene expression between mock-infected cells and HHV-7-infected cells at early (8-16 hpi) and late time points (48 and 72 hpi), and to identify significantly differentiated expression, using at a false discovery rate of 3%. Compared to mock-infected cells, 22 and 17 cellular genes were found to be significantly upregulated at early (8-16 hpi) and late (48-72) times post-infection, respectively, by HHV-7, whereas 12 and 29 cellular genes were found to be significantly downregulated at early and late times, respectively, (Figure 5.16). However, only one of the upregulated genes and 7 of the downregulated genes at late times had a fold-change above +1.5 or below -1.5. At early times only one gene had elevated expression levels above a fold change of +1.5 (Table 5.5). Some of these genes are involved in biological processes that have been identified in previous microarray studies of viral infections. For example, cholesterol biosynthesis was induced by HIV (van 't Wout et al., 2003). The expression levels of many genes involved in interferon-mediated response, apoptosis, protein synthesis, cytokine production and signalling, are increased by various viruses (Table 1.4). Changes in genes involved in transcription and cell cycle regulation have also been described previously (Table 1.4). Interestingly, hydroxysteroid (17-beta) dehydrogenase 12 was slightly upregulated in HHV-7 infected cells. Although the role of this gene in HHV-7 infection is undetermined, a related enzyme (3- beta-hydroxysteroid dehydrogenase) is encoded by vaccinia virus and may play a role in inhibiting host inflammatory response (Reading et al., 2003). However, in general we observed rather small fold-changes in the expression of only a small number of genes. in most published array studies of cellular trancription during viral infection, usually a larger number (up to hundreds) of host genes were up- and downregulated and very often

with much larger fold-changes. In comparison, our microarray data showed HHV-7 induced very little change in host gene expression.

Table 5.5 Significantly diffrentiated cellular genes in HHV-7 infected SupT1 cells

Gene	GenBank	Mean fold change	Function
	Accession		
Upregulated at early times	A1202996	+1.58	Apoptosis
Ubiguitin-conjugating enzyme E2A (RAD6	AA045751	+1.02	DNA repair
Ankyrin repeat and SOCS box-containing 3	<u>H40739</u>	+1.00	Protein synthesis
Killer cell lectin-like receptor subfamily D, member 1	<u>R78286</u>	<u>+1.00</u>	NK cells and T- cells cytotoxicity
Hydroxysteroid (17-beta) dehydrogenase 12	<u>N95053</u>	+0.93	unknown
Interleukin 11	AI079412	+0.92	Hematopoiesis
Apoptosis antagonizing transcription factor	<u>AI439571</u>	+0.91	Apoptosis
Diacylglycerol kinase_gamma 90kDa	BE468027	<u>+0.89</u>	Lipid metabolism
Transcription elongation factor A (SII), 1	<u>AI421762</u>	<u>+0.85</u>	Transcription
S100 calcium binding protein A3	N68995	+0.84	Cell differentiation
KIAA0053 gene product	AA262297	<u>+0.84</u>	GTPase activator
Joined to JAZF1	<u>AI378171</u>	+0.83	Transcription
v-ros UR2 sarcoma virus oncogene homolog 1	<u>W24201</u>	<u>+0.83</u>	Tyrosine kinase signalling
Regulator of G-protein signaling 14	AA428897	+0.82	G-protein signaling
Aldehyde dehydrogenase 9 family, member A1	AA010649	<u>+0.82</u>	Metabolism
<u>Homeo box A10</u>	<u>Al187278</u>	<u>+0.78</u>	Transcription
BCL2 binding component 3	<u>Al821703</u>	<u>+0.78</u>	Apoptosis
Leukotriene A4 hydrolase	<u>AA987920</u>	<u>+0.77</u>	Biosynthesis
Serine (or cysteine) proteinase inhibitor, clade D, member 1	<u>H72143</u>	<u>+0.77</u>	Peptidase activity
Centaurin beta 2	AA237088	<u>+0.76</u>	GTPase activator
Chromodomain helicase DNA binding protein 2	N90649	<u>+0.73</u>	Transcription
Associated molecule with the SH3 domain of STAM (AMSH) like protein	<u>W95888</u>	<u>+0.70</u>	Protein synthesis
Upregulated at late times			
Pirin	N29255	+1.98	Transcription
Associated molecule with the SH3 domain of STAM (AMSH) like protein	W95888	+1.46	Protein synthesis
Transcription elongation factor A (SII)	AI421762	+1.28	Transcription
dependent regulator of chromatin, subfamily a, member 3	BF51373	+1.28	Chromatin maintenance
Ankyrin repeat and SOCS box-coontaining 3	H40739	+1.19	Protein synthesis
Hydroxysteroid (17-beta) dehydrogenase 12	N95053	+1.13	unknown
Programmed cell death 1 ligand	AI202996	+1.12	Apoptosis
Succinate dehydrogenase complex	AA281913	+1.00	Carbohydrate metabolism
CD47 antigen (Rh-related antigen, integrin- associated signal transducer)	AA418709	+0.99	Signal transduction, cell adhesion
Centaurin_ beta 2	AA237088	+0.97	GTPase activator
Mitogen-activated protein kinase	AA236814	+0.91	Heat shock
protein')	AA283840	+0.90	Protein trafficking
Jun D proto-oncogene	Al339181	+0.88	Cell cycle
Isocitrate dehydrogenase 3 (NAD+) beta	AI566177	+0.85	Carbohydrate metabolism
TAF1 RNA polymerase II_TATA box binding protein (TBP)-associated factor	A1806456	+0.83	Transcription
Macrophage stimulating_pseudogene 9	AA262037	+0.81	
19A24 protein	BE46767	+0.72	
Downregulated at early times			
60s Acidic Ribosomal Protein P0		-1.43	Translation
Cytochrome P450 family 11A1	AA142879	-1.31	Cholesterol metabolism
AA400583	AA400583	-1.29	
40s ribosomal protein s18	/ 01-00000	-1.16	Ribosomal RNA, protein
40s ribosomal protein s3A		-1.15	Ribosomal RNA, protein biosynthesis
Glucocorticoid receptor alpha mRNA	AA736674	-1.03	Transcription
Human triosephosphate isomerase		-0.99	Metabolism
Integrin_alpha 2 (CD49B_alpha 2 subunit of VLA-2 receptor)	AI538195	-0.92	Cell adhesion

GS2 gene	AI290788	-0.86	
Low density lipoprotein recept (familial hypercholesterolemia)	AA621897	-0.82	Cholesterol metabolism
Leukotriene A4 hydrolase	AI521212	-0.81	Biosynthesis
Homeo box A5	AI223317	-0.73	Transcription
Downregulated at late times			
Cytochrome P450 11A1	AA142879	-3.10	Cholesterol metabolism
60s Acidic Ribosomal Protein P0		-2.39	Translation
40s ribosomal protein s18		-2.10	Ribosomal RNA
Histone 1_ h2bk	AA280257	-2.09	Histone
Guanine nucleotide-binding protein		-1.58	G-protein signaling
Branched chain keto acid dehydrogenase kinase	W78018	-1.58	Protein synthesis
Human triosephosphate isomerase		-1.55	Metabolism
Death associated protein 3	AA858184	-1.25	Interferon-mediated inflammation
Peptidyl-prolyl:cis-trans isomerse		-1.23	
Heterogeneous nuclear ribonucleoprotein C	AI096954	-1.21	Ribonucleosomes
Lymphoid enhancer-binding factor 1	AF086339	-1.19	
TATA box binding protein	AA470112	-1.19	Transcription factor
Tubulin gaunachain		-1.12	
Aa459022	AA459022	-1.09	n/a
Low density lipoprotein recept (familial hypercholesterolemia)	AA621897	-1.07	Cholesterol metabolism
Peroxiredoxin 6	AA016074	-1.05	Oxidative Stress
Human ribosomal protein L30		-1.01	Protein synthesis, mitochondrial
Interleukin 18 (interferon-gamma-inducing factor)	Al394321	-1.01	Interferon-mediated inflammation
Muscleblind-like 3 (Drosophila	AA460747	-0.99	
UL16 binding protein 1	AI830832	-0.98	Cytokine signaling
Homo sapiens transcribed sequeuence	AA748610	-0.95	Cell cycle
Ai245910	AI245910	-0.94	
Tripartite motif-containing 29	AA293775	-0.93	
Aw449304	AW449304	-0.92	
SWI/SNF related_ matrix associated, actin dependent regulator of chromatin, subfamily a, member 4	AI279006	-0.92	Transcription regulation
Postmeiotic segregation increased 2-like 5	AJ015925	-0.82	
AT-hook transcription factor A	AA904466	-0.82	
SH2 domain protein 2A	AA613070	-0.80	Signal transduction
C-type lectin superfamily member 12	4478515	0.74	

* Mean \log_2 fold change of the triplicate spots on custom-made host-pathogen arrays

There were three possible reasons for this lack of change in host gene expression during HHV-7 infection. Firstly, the relatively small number of cellular probes (approximately 1000) on the custom-made array may have meant that genes that underwent transcriptional changes were not represented on the array. However, this was not likely, because when the same samples were hybridised onto the Human cDNA Array v.2 (Human Genome Mapping Project, MRC, UK), which contained approximately 11,000 human gene probes, the clustering results were similar to the that obtained using custom-made arrays. Secondly, sense strand RNA amplification, which favoured the 5'-end of mRNA transcripts, may have introduced a bias that prevented the detection of subtle changes in cellular transcription. However, the signals of the host gene probes were sufficient for detection, and the pass-rates were adequate (Figure 5.17). Therefore sense-strand RNA amplification had produced enough cellular RNA for hybridisation. Sense-strand amplified RNAs have been analysed by quantitative RT-PCR and was shown to preserve known expression ratios (Goff et al., 2004) and the relative changes in viral transcription levels were clearly conserved after amplification (section 5.2.3). However, it was unknown whether the differences in cellular mRNA abundance were affected by sense strand mRNA amplification in this study. If this was true, however, it would be very difficult to analyse host gene expression during HHV-7 infection using microarrays, because RNA amplification was necessary to ensure a high MOI and synchronous infection.

A third possible reason for the lack of differential expression may be that HHV-7 caused no major changes in the expression of host genes during infection of SupT1 cells *in vitro*. This was also difficult to comprehend: how the SupT1 cells could show no response to a virus that infected 80% of the culture population (section 5.2.1), replicating its genome and producing virions (section 5.2.3). However, if true this may reveal details about the ability of the HHV-7 virus to infect 90% of the world's population without apparent pathologies (section 1.1.9) and would represent the extreme of herpesvirus mutualism with human hosts. So far this phenomenon has not been observed in any other virus. All microarray studies of viral infections have shown that host gene expression was extensively altered during infection (section 1.2.4.3, Table 1.4). It will be interesting to study, if true, why HHV-7 induced no large changes in cellular transcription.



Possibly downregulated genes

Possibly upregulated genes

Figure 5.15 Cluster analysis of host gene expression in SupT1 cells mockinfected and infected with HHV-7 over a 72-hour period, measured using custom-made microarrays, with two clusters representing genes possibly upregulated (red rectangle) and downregulated (green rectangle) by HHV-7



Figure 5.16 Scatter plot of observed (y-axis) vs expected (x-axis) expression ratios of cellular genes from HHV-7 infected cells and mock-infected cells using Significance Analysis of microarrays (SAM). Significantly upregulated genes (red dots), significantly downregulated genes (green dots) and unchanged genes (black dots)





Figure 5.17 a) Background-subtracted Cy5 intensities and b) percentage pass rates of cellular probes (white bars) and HHV-7 probes (striped bars) on the custom-made microarrays hybridised to amplified RNA extracted from HHV-7-infected SupT1 cells at various time points during infection.

5.3 Summary and Discussion

This chapter described the analysis of HHV-7 gene expression using custom-made microarrays, demonstrating the power of microarrays as a large-scale analysis tool capable of deciphering the whole transcriptome of the virus. Transcription profiling revealed the coordinated expression of viral genes during lytic infection of T-cells in vitro. The regulated induction of HHV-7 genes from 8 to 72 hours post infection reflected the sequential stages of viral infection cycle, consistent with the classic model of temporal regulation of herpesvirus gene expression. From the profiling data we have also estimated the HHV-7 lytic cycle to be between 36 to 60 hours long, predicted functions for previously uncharacterised genes, and identified the possible roles of certain transcription regulators. Microarrays also enabled the assignment of 68 HHV-7 genes into temporal classes A, B, and C (corresponding to IE, E and L). A final consensus of temporal classes (Figure 5.18) was achieved from the results of transcription profiling (section 5.2.3); inhibitor treatment coupled with microarrays analysis (section 5.2.4) and RT-PCR (section 5.2.5). Genes of the same temporal class were sometimes adjacently situated in the viral genome, suggesting the possibility of polycistronic transcription (Figure 5.18).

Temporal classification of the HHV-7 genes varied between the different methods. This was probably due to the different features of the dataset emphasised in the different methods. The transcription profiling used (section 5.2.3) was a comparison approach, measuring the relative changes in expression over time. It revealed the temporal regulated patterns of transcription in the absence of metabolic inhibitors, therefore revealing viral gene expression with minimal artificial intervention. The use of microarrays coupled with inhibitor treatment (section 5.2.4) was a rapid method for identifying statistically significant genes that were differentially expressed between treated and untreated cells. The microarray technique was not very sensitive for the detection of low abundance transcripts, especially in 8 hpi samples, therefore was unable to distinguish between IE and E genes. Nevertheless, its high throughput nature allowed many genes to be analysed simultaneously and rapidly. Low abundance genes that require further investigation using other methods was rapidly identified for follow up in future experiments.

RT-PCR, on the other hand, was a highly sensitive qualitative approach capable of detecting very low abundance mRNAs (section 5.2.5). It provided a simple positive/negative answer to whether a gene was transcribed in the presence of metabolic inhibitors. However, it has been suggested that 35 PCR cycles were too sensitive and may lead to detection of leaky E transcription under IE conditions and misassignment of kinetic classes (Pellett, 2001). For example, U42 and U73 of HHV-6, classified as IE genes by qualitative RT-PCR (Mirandola et al., 1998), were found to be dependent on *de novo* protein synthesis by quantitative RT-PCR (Rapp et al., 2000). This may explain the inconsistencies between microarray-based methods and RT-PCR in the classification of IE and E genes (Table 5.3 - 5.4).

In summary, RT-PCR was more sensitive for low abundance transcripts and has the advantage of revealing transcription kinetics for different splice forms. Microarrays offered faster, larger scale temporal classification of herpesvirus genes in the presence or absence of toxic metabolic inhibitors. However, the strength of microarray technology was its capacity to disclose the relative changes in gene expression, resulting in a global picture of the molecular events during HHV-7 infection of T-cells.

The microarray data also showed that HHV-7 infection induced very little changes in host gene expression, with only 8 genes showing significantly differentiated expression above a fold change of 1.5 (section 5.2.6). However, further experiments are required to validate this finding.



Figure 5.18 Temporal map of HHV-7 genome (RK strain). Predicted functions of the ORFs are colour-coded:

metransactivators; replication functions capsid & assembly; tegument; envelop othe unknown.

Array probes of approximately 300 bp were designed for each ORF (small black rectangles); also for ten intergenic regions (small grey rectangles) to measure the expression level of their corresponding ORF. HHV-7 ORFs were classified as IE (green block), E (blue block) or L (red block) genes by expression profiling using arrays; also as IE/E (turquoise block) or L (red block) by arrays with inhibitor (PAA). Kinetic classification of some ORFs were confirmed by RT-PCR (marked with *).

CHAPTER 6

CONCLUSIONS & FUTURE DIRECTIONS

6.1 HHV-7 gene expression

This thesis described the production of a sensitive, specific and reproducible hostpathogen microarray for the detection of human herpesvirus type-7 (HHV-7) mRNAs, and its application in expression profiling of the viral transcriptome during human T-cell infection in vitro. Analysis of the microarray results has revealed that HHV-7 genes were co-ordinately expressed in an ordered pattern, which reflected the sequential stages of viral lytic replication cycle (transcription activation, replication, virion assembly) that is common amongst herpesviruses (Pellett, 2001; Roizman, 2001a; Roizman, 2001b). HHV-7 genes were classified into three temporal classes A. B or C in the absence of metabolic inhibitors, therefore represented viral lytic infection without any external intervention. These classes generally mirrored the herpesvirus immediate-early (IE), early (E) and late (L) genes, respectively. This correlated well with previous assignments of herpesvirus genes into their temporal classes by transcription profiling (Jenner et al., 2001; Paulose-Murphy et al., 2001). A selection of class B and C genes have been confirmed to be E and L genes, respectively, by the use of phosphonoacetic acid (PAA) to block viral DNA replication, hence L gene expression. The confirmation of HHV-7 IE classification, however, would require further work, due to cellular toxicity of cycloheximide (CHX) at 16 hours post infection (hpi) and low expression level of early genes at 8 hpi. A useful experiment to do would be to compare HHV-7 gene expression at 12 hpi in the presence and absence of cycloheximide, in order to distinguish between viral IE and E genes.

Nonetheless, we have shown that transcription profiling using microarrays allowed the temporal classification of herpesvirus genes consistent with metabolic inhibitor assays. Microarray analyses offer higher throughput, reproducibility, speed and a means to measure genome-wide viral gene expression in their 'natural' state, thus have potential advantages over metabolic inhibitor treatment assays. The HHV-7 host-pathogen microarrays we created could be used to study the transcriptome of clinical HHV-7 isolates from latently-infected adults, or from children with primary infection, and these can be compared with the results obtained in this study using the DC strain.

Being the third member of the human betaherpesviridae family, unlike human cytomegalovirus, HCMV (Chambers et al., 1999) and human herpesvirus type-6, HHV-6 (Ohyashiki et al., 2005), no extensive viral gene expression studies have been performed on HHV-7. This thesis has made a step to complete that picture, showing that HHV-7 gene expression patterns were similar to the other betaherpesviruses, and that the three human betaherpesviruses all follow similar lytic replication events *in vitro*. However, the genomes of HCMV and HHV-6 contain extra ORFs that are not found in HHV-7, and it would be interesting to compare the gene expression profiles between these betaherpesviruses. This could be performed using the custom-made microarrays, as array probes for HHV-6 ORFs and a subset of HCMV genes have been generated in our laboratory as separate projects.

In addition, findings in this study also supported the predicted functions of HHV-7 ORFs (Megaw et al., 1998; Nicholas, 1996), and suggested functional classes of some previously uncharacterised genes. We have also estimated the time of action of some transcriptional regulators, hence indicating their possible roles in particular stages of the viral lytic infection cycle. HHV-7 U42 and U54 may be responsible for regulating the transcription of genes encoding enzymes for viral DNA replication and structural proteins, respectively. The functional significance of these two transactivators could be examined by gene knock out experiments, using gene silencing techniques such as RNAi (Hannon, 2002; Kuwabara and Coulson, 2000). RNAi has been used to study host-pathogen interactions in bacterial (Agaisse et al., 2005; Walduck et al., 2004) and viral infections (Lieberman et al., 2003; Wang et al., 2003). The impacts on viral DNA replication and virion assembly upon RNAi silencing of HHV-7 U42 and U54 may provide useful information on the functions of these transcriptional regulators, and thus improve understanding of how herpesviruses regulate their replication events during infection.

We have demonstrated the power of microarray transcription profiling in the study of large genome viruses. So far these techniques have only concentrated on herpesviruses, while no global gene expression profiling studies have been published for the other virus families with large dsDNA genomes, such as the poxviridae and the adenoviridae. Gene expression in poxviruses are also temporally regulated (Broyles, 2003), thus is particularly suited for transcriptional profiling using microarrays. Although the genome of vaccinia virus has been fully sequenced (Goebel et al., 1990) and a microarray has been generated for vaccinia genes (Wenli et al., 2004), it has not yet been used for the transcriptome profiling of the vaccinia virus.

6.2 HHV-7 culturing

This thesis has also documented the optimisation of HHV-7 quantification methods and protocols for maximising the infectious titre of the virus in SupT1 cells in vitro. Previous attempts to improve HHV-7 infectivity by methods such as ultracentrifugation (Ablashi et al., 1998) were without success. We have shown that growing HHV-7 in small volumes was an efficient method for maximising the infectious titre of the virus stocks harvested (section 4.2.4.1). This is also the first report on the effects of spinoculation and polybrene on HHV-7 inoculation efficiency. The multiplicity of infection (MOI) of HHV-7 in T-cells in vitro was reduced in the presence of polybrene (8 µg/ml), but was improved by spinoculation at 1000 g, although the increase was not as high as for some other viruses (Pietroboni et al., 1989). Therefore the effects of spinoculation and polybrene on HHV-7 infectivity were similar to that on its closest relative HHV-6 (Black et al., 1989), but not to the more distant gammaherpesvirus HHV-8 (Inoue et al., 2003), of which the infectivity was increased by both spinoculation and polybrene. These findings would benefit future studies requiring the infection by cell-free HHV-7 virus at a high MOI, particularly gene expression analysis, which has been hindered by the low infectivity of HHV-7 (Pellett, 2001).

RNA amplification has been applied to analyse limited clinical samples such as those extracted by laser capture microdissection or flow cytometry-sorted cells. We have demonstrated that the RNA amplification technique could also be used to amplify RNA from low cell numbers in order to maximise the MOI in virology studies. This may be useful for other viruses with low infectivity-to-particle ratios in cell-free virus form, for example human T-cell leukaemia virus type 1 (HTLV-1) (Sasada et al., 2005), human and murine cytomegaloviruses (Ho et al., 1993; Hodgkin et al., 1988; Hudson, 1988; Osborn and Walker, 1968), and retroviruses (Bahnson et al., 1995;

Chuck et al., 1996; Ho et al., 1993; Kotani et al., 1994). This technique may also benefit gene expression studies on slow growing micro organisms such as *Plasmodium falciparum*, and *Mycobacterium tuberculosis*. However RNA amplification adds cost and time to the experiment therefore may not be suitable for some applications.

6.3 Host genes expression by HHV-7

We have found little change in the expression of cellular genes during HHV-7 infection of SupT1 cells in vitro. This could be due to technical limitations or HHV-7 may be very efficient at evading host response. Confirmation requires further work on validating the use of sense-strand RNA amplification technique on cellular mRNAs. The sense-strand RNA amplification method used in this thesis can be used to amplify RNA from samples that are known to contain significantly different amounts of certain species of cellular mRNA. However, if HHV-7 truly induces so little response in the host during infection, more analysis on how HHV-7 achieves this may provide very useful information about host-pathogen interactions and viral evasion of host immune response. In particular, silencing of candidate HHV-7 genes may give clues to which, if any, of the HHV-7 genes are responsible for the virus's ability to establish such silent infection in the host cell. Moreover, this naturally nonpathogenic virus may be a very useful potential gene therapy vector. However, before HHV-7 can be developed into a gene therapy vector, more studies are required to characterise HHV-7 cell tropism, also the mechanisms for cell attachment, entry and expression of viral-encoded antigens on the cell surface.

Appendix A – Sequence alignments of HHV-7 array probes

HHV7DR1 & DR1'

HHV7_RK_DR1 HHV7_RK_DR1' H1_020206 A1 020228	ATGACTGCTGCAACCACAGAACATTTTGCTCTCCGCGCGCACTCAATCGTTACTGGTGG ATGACTGCTGCAACCACAGAACATTTTGCTCTCCGCGCGCACTCAATCGTTACTGGTGG 	60 60 42
HHV7_RK_DR1 HHV7_RK_DR1' H1_020206 h1_020238	CTGCTTCTGGGACGACACAAGCTCAGTTTGGTATGCAACT-ACGTCACAG-CTCATCGCC CTGCTTCTGGGACGACACAAGCTCAGTTTGGTATGCAACT-ACGTCACAG-CTCATCGCC CTGCTTCTGGGACGACAAAGCTCANTTGGTATGCAACT-ACGTCACAG-CTCATCGCC	118 118 100
HHV7_RK_DR1 HHV7_RK_DR1' H1_020206	AAC-AGTTACTGCCGCTGCCGTGGCCCGAACAGGAATTCCCCAACTTGA-CCCGGCCCC AAC-AGTTACTGCCGCTGCCGTGGCCCGAACAGGAATTCCCCCAACTTGA-CCCGGCCCC AAC-AGTTACTGCCGCTGCCGTGGCCCGAACAGGAATTCCCCCAACTTGA-CCCGGCCCC	176 176 158
A1_020228	AACCAGTTACTGCCGCTGCCGTGCCCCGAACAGGAATTTCCCCAACTTGAACCCGGCCCC	101
HHV7_RK_DR1 HHV7_RK_DR1' H1_020206 A1_020228	CTACTCCAATCTCCGCAACOGT-GTCGC-TCACCATCTCCATCGCGGCTGGCCAGCGGCA CTACTCCAATCTCCGCACCOGT-GTCGC-TCACCATCTCCATCGGGCGGGCCAGCGGCA CTACTCCAATCTCCGCAACOGTFGTGGCATCACCATCTCCATCGGGCGGGCAGCGGCA CTACTCCAATCTCCGNCNCCCCCGT-GTACTCCCATCTCCNTTNCGGGTGG	234 234 218 151
HHV7_RK_DR1 HHV7_RK_DR1' H1_020206 A1_020228	CA-CAACACAtgtaagctaccgtacatctctttca-caaacccaaggctcacatagagac CA-CAACACATGTAAGCTACCGTACATCTCTTTCA-CAAACCCAAGGCTCACATAGAGAC CNACAACTCATGTAAGCTACCGTACATCTCTTTCATCAAACCCAGGGCTCACATAAAGAC	292 292 278
HHV7_RK_DR1 HHV7_RK_DR1' H1_020206 A1_020228	aagcacaagctcgcgcaatgacattaaaacctcccatcattgtcctttcctgtcgctttg AAGCACAAGCTCGCGCAATGACATTAAAACCTCCCATCATTGTCCTTTCCTGTCGCTTTG AAGCACAAGCTCG-CAATCACT-	352 352 299
HHV7DR6 & DR6'		
HHV7_RK_DR6 HHV7_RK_DR6' H2_020425 A2_020206	ATGAGTGCAGAAATGCTCCGCGCTGTTCAGCTCCAGCCAAGACGCCGGGGACATTCCTCA ATGAGTGCAGAAATGCTCCGGCGTGTTCAGCTCCAGCCAAGACGCCGGGGACATTCCTCA 	60 60 39 60
HHV7 PK DP6	₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽₽	114
HHV7 RK DR6'	TCTCCCACTTCCCCTCCACTCGAAGGAGAGCCCCAGTCCCAAGAGACTCCAATCG	114
H2_020425 A2_020206	TCTCCCACTTCCCCTCCACTCGAAGGAGAGCCCAGTCCCAAGAGACTCCAATCG TCTCCCACTTCCCTNTCGNANNTTCGTNNAGGAGAGGCCCAGTCCCAAGAGACTCCAATCG	93 120
HHV7_RK_DR6 HHV7_RK_DR6' H2_020425 A2_020206	АGCААСАӨТСАССААӨЗӨСӨТАДАӨЗСАВАССТАААСССАВАЗСТААААСАТӨЗӨАЗСВА Азсхасхагтасасаабдосогтадазсадасстааасссаядстаааасатөздазсва азсяасаястассаабдосогтадазсадасстааасссаядостаааасатөздазсва азсяасаятассаабдосогтадазсадасстааасссаядостаааасатөздазсва азсяасаятассаабдосогтадазсадасстааассаядостаааасатөздазсва	174 174 153 180
HHV7_RK_DR6 HHV7_RK_DR6' H2_020425 A2_020206	GCTTTATCCCACCGGTCCTTCCTCAACATTTACGCGTGGCTGTCTTTGAGT-CGAGGGTC GCTTTATCCCACCGGTCCTTCCTCAACATTTACGCGTGGCTGTCTTTGAGT-CGAGGGTC GCTTTATCCCACCGGTCCTTCCTCAACATTTACGCGTGGCTGTCTTTGAGT-CGAGGGTC GCTTTATCCCACCGGTCCTTCCTCAACATTTACGCGTGGCTGTCTTTGAGTGCGAGGGTC	233 233 212 240
HHV7_RK_DR6	TCCGCGARARAGTGTACGGATATGCCTTCAGGCACAGA-GGAGAACTCGTAGCATTG-CCA	291
HHV7_RK_DR6' H2_020425 A2_020206	TCCGCGAAAAGTGTACGGATATGCCTTCAGGCACAGA-GGAGAACTCGTAGCATTG-CCA TCCGCGAAAAGTGTACGGATATGCCTTCAGGCACAGA-GGAGAACTCGTAGCATTG-CCA TCCGCGAAAAGTGTACGGATATGCCTTCAGGCACAGNAGGAGAACTCGTAGCATTGANCA	291 270 300
HHV7_RK_DR6	TGGCCGCCTAACTGGAGCCTGGAACTTCACCACGATCCCTATCGAGACGCCAGAGGACAA	351
HHV7_RK_DR6 [.] H2_020425 A2_020206	TGCCGCCTAACTGGAGCCTGGAACTTCACCACGTCCCTTCGAGACGCCAGACGCAGA TGGCCGCCTAACTGGGGCCTGGAACTTCA TGGCCGCCTAACTG	351 299 314
HHV702		
HHV7_RK_U2 A3_020419	TCTGCGCAGACCACGCTTCCAGAGACCATAAGGGGTGTCCGCAACTCTGTAAATGGCGTT 6 	60
HHV7_RK_U2 A3_020419	GTCCACGGAACGGTTAAATGCATAAAATTCACCCCGATCTCCAATAAGCATCAAAGCAGA 7 -TANANCGTTTGNCNAGAANCTTNCGGTCCTTCNCGGAANANTGGCNNCAAGAAACATA 6 * * * * * * * * * * * * * * * * * * *	20 2
HHV7_RK_U2 A3_020419	TAAATCAGAATAGAGCACCAGACGATGACCCAAACTTCCACAAAAGGACAGCAGACAAA 7 CTCN-CGNAAGAGGGCAAAACGCNTCAAGCTCGAGNNCGTCGCGTGTAGAGGNCGCG 1 * ** ** ** ** * * * * * * * * * * * *	80 18
HHV7_RK_U2 A3_020419	ATTTCTGGTTAACTTGCAGACTTCGTCTCCGAAATGCCAAGCATCTCCGAAGAGGCAAA 8 AACAGCGGAGGGGGANAGCNTINNGNN-GANNTCNGGTGTAGGNGNANGGANGTNTG 1 * ** * * * * * * * * * * * * * * * *	40 77
HHV7_RK_U2	CCTCAACTCAGCGCCGGAAGGCCACGCGAGGGAAGTCGATGGTGAGCATATTGCAAAAC 9	00

A3_020419	NCCNAGNGGAGANGCGANAGANNNN-TENGNGGNTCGGGAAGNGGTGCGGTGCGNGGAGG 230	6
HHV7_RK_U2 A3_020419	AAAATGAGCAACGTCTCCGAAAGATACGGTTTCCGCAGCGAAGCTCTGCAAAAACGTG 95 NGGAGGAGTAGCGGTTGTGTGTCGCGCGGCGGCGGAAGNNCTCTGTCCAAAAANCGTG 29 * *** * *** * * * * * * * * * * * * *	8 1
HHV7_RK_U2 A3_020419	GTCC-AGTAGACCTCCGGAACAGGTGATTCAACTGTATTCGATTCCAGATT-AAACAA 10 GRNCCAGTAGACCTCCGGAACAGGTGATTCAACNTGNTATTCGATTCCAGANTTAAACAA 35	14 1
HHV7_RK_U2 A3_020419	ACCOTTCGATGCATTCGACTCAACAGCCGTTTCCACACCGGAAATCCAAAAG-CGGAGAAT 10 ACCNTTCGATGCATTCGACTCAACAGCCGTTTCCACACCGGAATCCAAAAGTCGGAGAAT 41	73 1
HHV7_RK_U2 A3_020419	CTGACAT 1080 CTGAT 416 ****	
HHV703		
HHV7_RK_U3 A4_020413	TCAGTCTCATACTCAAGTTCGTAAACAGTCGTCACAGTATGTTTATGTAACCAAGAACA 78 NCNNGNTNNTNAANCGG 17 + +	0
HHV7_RK_U3 A4_020413	GTAAGTTCTTCAGGACAACACACACCGACGGGGGTACTTTTTGAGCTTATCTTCCTCA 83 NNAATNCCCCCGAGTCANCNCACGGGNATTNC-TGCTNCATGGNCCGCATACCGAACNGA 76	7
HHV7_RK_U3 A4_020413	TATCCCTCCACACCATAAGTCAAAACAAGTTCAAACCCCT-CTGGAAAAAACCAAAG 89 GCGACGCNCATATCAACTCAGTNGATTAACAGTTANAAACCTNNACNGGAAAAAACCAAAG 13 * ** * ** * **** * ***** * ****** * ****	2 6
HHV7_RK_U3 A4_020413	G-САСССТСАЛАТССОДАТТСАДАЛАСАЛАЛАСАТТТТТАДАСТАТТАЛАЛТТТСТАТ 94 GACACCCTCAЛАТССОДАТТСАДАЛАСАЛАЛАСАТТТТТАНДАНСТАТТАЛАЛТТТСТАТ 19	9
HHV7_RK_U3 A4_020413	AAACACGTAGAAAATCCACGAGTTTAGCCTTAGCAGATTCAAAGATTTTCTCTTTTCTTT 10 AAACACGTAGAAAATCCACGAGTTTAGCCTTAGCAGATTCAAAGATTTTCTCTTTTCTTT 	09 6
HHV7_RK_U3 A4_020413	TTTTATCA-GATAAAAACGTTTTCTCAACGTTGGCAACCTTCAAATCCTTTTTCCCGTCA 10 TTTTATCACGATAAAACGTTTTCTCAACGTTGGCAACCTTCAAATCCTTTTTCCCGTCA 31	68 .6
HHV7_RK_U3 A4_020413	ACATGAGC-AACATATTTTCGTATAAACGAATTTTCCTTTTTTAAATCTTTTGCCACAAC 11 ACATGAGNCAACATATTTTCGTATAAACGAATTTTCCTTTTTTAAATCTTTTGCCACAAC 37	.27 6
HHV7_RK_U3 A4_020413	ТСТСАЛАЛСЯССТСТТСАЛАЛТТGАЛGTCAGACCCCAAATCTGCCAT 1175 ТСТСАЛАЛСЯССТСТТСАЛАЛТТGAAGTCAGACCCCAAATCTGCA 422	
HRV7U4		
HHV7_RK_U4 A5_020313	TGTAGCTTTAGTCAGAACTCGATAAAACGGAGAAGTGCACTCTTCAAACCTTAGACTTTG 13 	20 5
HHV7_RK_U4 A5_020313	AAGAAACGCGAGAAAACCGTCTAAAGTCATTGTTGCCTTGTCCCTCCTTTTAAATGAAAC 13 AAAAAAAAAAAAAACAAATINNAAAAAANNNGATCTGTGCCGGTNCCCCTTAAAGTGAAAC 73	80 8
HHV7_RK_U4 A5_020313	GGAGTTTTTCTCCACTAAAACAGCCCATAGTAGACTGTCATCGTAAAATTTTAAGCC 14 GAGNNTCTCCNCAAAACAGCCCATAGTAGACTGTGNTCGTAAAATNCNCACCAGCC 12	137 19
HHV7_RK_U4 A5_020313	TCGAGTTTGGAACAAATCTTCTGAAGTGAGATGACCGCTATCCGCAA-TCATCTCCTCTG 14 TCGAGTTTGGAACAAATCTTCTGAAGTGAGATGACCGCTATCCGCANATCATCTCCTCTG 18	196 39
HHV7_RK_U4 A5_020313	TCAGCTCATTCAAGTTGACGTCTAACTCGCATGGAGTTAAATGCCGTCCATGACAGATAAG 15 TCAGCTCATTCAAGTTGACGTCTAACTCGCATGAGTTAAATGCCGTCCATGACAGATAAG 24	556 19
HHV7_RK_U4 A5_020313	GATGATTCGGTATAACATACGTAAT-GCGATTCCTTAGCGGCCCTTTACAAACATCATAA 16 GGTGATTCGGTATANCATACGTTTTTGCGATTCCTTAGCGGCCCTTTACCAC 30	515 D1
HHV7_RK_U4 A5_020313	тодатталатосат 1629	
HHV7U7		
HHV7_RK_U7 A6_020424	AAAGTCCGGGTCCGGTTCTCGGTAATTTGGATGTGCATAACCAACAGCGCAAAGAAATTC 24 TGGTCGGGTTCTCGGTAATTTGGATGTGCATAACCAACAGCGCAAAGAAATTC 53	160 3
HHV7_RK_U7 A6_020424	GTCTGCTCCTACCAAATTACTCCAATCATAAGTGATGACACTACGTCTACATTTGAAATG 2 GTCTGCTCCTACCAAATTACTCCAATCATAAGTGATGACACTACGTCTACATTTGAAATG 11	520 L3
HHV7_RK_U7 A6_020424	АТААЛАТТСІССАААЛАТАЛІСАСАТАІСТТІЗІСІІССААІСАЛІСАІСАІСТТТАТАСС 25 АТААЛАТТСІЗССААЛАЛТАЛІСАСАТАІСТТІЗІСІІССАЛІСАЛІСАІСАІСАІСТТТАТАСС 17	580 73
HHV7 RK U7	ATTGCGAACTATCTCCATGCCTTCTCCATATGAAGACAGCAGCAACTCAACGGCAAAATC 26	40

HHV7 RK U7	AAAAGACGGTAAATCCGGAGTCGGTTCAATGCAAACATCATT-GCACACAGGAACAATTT	2699
A6_020424	AAAAGACGGTAAATCCGGAGTCGGTTCAATGCAAACATCANTTGCACACAGGAACAA	290

H

HHV708	
HHV7_RK_U8 A7_020313	TTTCCTTCCAATTTGAATCGCTCCAATCAAGCACAAATTTTCCTTAAATCCTTGGTAATG 840 CAATCAAGCNCAAATTCCTTAAATCTGGTA-TG 32
HHV7_RK_U8	GTCATTCAGTTCTGCCAATTGTTG-TTCTGAGTAACCAGCAACGTCAACTGGCTGTGCAA 899
A7_020313	GTCATNNN-TNCTGCCAATNGTTGGTNCTGAGTNNCCCCCN-CGTCA-CTGGCTGTGCA- 88
HHV7_RK_U8 A7_020313	GCCTTAACCACAAATTCTTTGGCCAAATCAAAGACAAGCATTCATT
HHV7_RK_U8 A7_020313	TAAGAAGGGAATTCCCAGGGAAATCAGTAAAACGACATAAATTCTTCAATTCCCTTAACA 1019 TAGAGGCGAATTCCNGGGAATCNTNNCGAC-TAINITCATTCCCTTA-CA 192 ************************************
HHV7_RK_U8	ATCCATCATTTAGTTGTCCAACACAAGGCAAAGTTTTATATTGTTCTGCCATCGTCT 1076
A7_020313	TCNNNTNNTTTTTTGTCCNCCCTGGGCAAGTNTTATATNGTNTGCCTCTCTCNTCNTNC 252
HHV7_RK_U8 A7_020313	CARARTTTTCAT 1089 CCGCGGNNNNTGGCNGCNGGGANCTGCACCTNGGCCNNTNC 294 *
HHV7U10	
HHV7_RK_U10 A8_020502	ATGGCTATAGCAGAAAGCGGAGCGGTTCGAAATAAATGTCAATCTTGGAAAAATCCATCGCT 60 ATTNAAATGAAAGCGNGGNANNGCTTCTTCATGTTATTTCTNGTGTNCTCCT 52 * * ****** * * ****
HHV7_RK_U10	TCCATACAAAAAAATCCTATAAATCTACATGCGTAGACATCTCTCTTTTATGTGGAGTTA 120
A8_020502	ACCANACNAATAANCTANTATANATCCGRTCNNTACNTTCTCTTTTATTGTGGAGTTN 110
HHV7_RK_U10	TTAAAATTTATCATCATCAATAATGAGCAATGTTTTTTACCACCGAAAGGAACGATA 177
A8_020502	NTNAAATACCTNATCATCATTCATCAATATGAGCAATGTTTTTTACCACCGAAAGGAACGATA 170
HHV7_RK_U10	СТТТАССАСААТGGCTTAATCGAACTAAATACTTTAATTATCGATCTGAATCAACAAATT 237
A8_020502	СТТТАССАСААТGGCTTAATCGAACTAAATACTTTAATTATCGATCTGAATCAACAAATT 230
HHV7_RK_U10	ACATCGAAACAGCAGATCTATAGTTGGACGAGCATTACTTTGCCAAAAATATTTTCGACT 297
A8_020502	ACATCGAAACAGCAGATCTATAGTTGGACGAGCNTTNCTTTGCCNTNTTTTCGACT 286
HHV7_RK_U10 A8_020502	AAAGAATTATATTTTATTGTCGCGTCACCAGAATCGGAAAACATAACTCTCAATCCAGCT 357 NGATTATATTTTNTTGTCGCGTCNNN 312 *********** *********
HHV7011	
HHV7_RK_U11	TAAGATTGACGTAATCGGCTGCGTGTGTCACAGTGCCCCAACGAAGAAAACTGCCCAGTT 1860
A9_020424	
HHV7_RK_U11	TAAGCAAATCTTTTGCTACGTTTTTAAATTCTGTTTCAAAG-CCAACTTTGTTAAAAGAT 1919
A9_020424	TTAGCANATCTTTTGCTACGNTTTTTAATTCTGTTTCNAAGGCCAACTTTGTTNAAAGAT 70
HHV7_RK_U11	GTTAAACTAGAATAATCAGTAATCAGTCTTTGGTCTAAGTAATCTTTCAGAATTTTT 1979
A9_020424	GTTANACTAGAATAATCAGTAATCATCTGTCTTTGGTCTAAGTAATCTTTCAGAATTTTT 130
HHV7_RK_U11	TGCACCTTTAGTATTTCTTGCCAAACTTCCTCATAGTCTGGTTTCTTTTTACACAAGGTT 2039
A9_020424	TGCACCTTTAGTATTTCTTGCCAAACTTCCTCATAGTCTGGTTTCTTTTTACACAAGGTT 190
HHV7_RK_U11 A9_020424	тдатдаалаталатссаладаатсалаттуттуталаутуттттасаттуттаатудаа 2099 тдатдалаталатссаладаатсалаттуттуталадтутттасаттуттаатудаа 250 ********
HHV7_RK_U11	ТТТССАЯТТТТАЛСВАТАСАТТСЯЛСТСАЛЛАТССАТСВАТТТТСЭССВАЛТАТСАЛСАЛСТ 2159
A9_020424	ТТТССАЯТТТТАЛСВАТАСАТТСЯЛСТСАЛЛАТССАТСВАТТТТСЭССВАЛТАТСАЛСАЛСТ 310
HHV7_RK_U11	GGTAAAGACGATATGTTTTCAAAAAATCGAGATAAAAAACACTTTGCTTCATCCGAAATC 2219
A9_020424	GGTAAAGACGATATGTTTTCAAAAAATCGAGATAAAAAACACTTTGCTTCATCCGAAATC 370
HHV7_RK_U11 A9_020424	CAAGCAAAGGGTAAATGTGACATTTTCAT 2248 CA 372 **

HHV7012

HHV7_RK_U12 A10_020413 CAGCATGATGAAGAGTACAAGTACAAGTACGAATTATACGTGTATTACGCCAACAGTACGGAAAGCC 180 -----TTACGCCAACAGTACGGAAAGCC 23

HHV7_RK_U12 A10_020413	CANAGACITGANAGCGINATTANCGGANITATGCINACGCIGAIACTICCIGITAGIACI CANAGACITGANAGCGINATIAACGGANITAIGCINACGCIGAIACITICCIGITAGIACI ******	240 83
HHV7_RK_U12	GTTGTCATATGCACTCTGCTAATCTACTACAAATGGACAAAACAGACAATTACTTCTCCCA	300
A10_020413	GTTGTCATATGCACTCTGCTAATCTACTACTACAAATGGACAAAACAGACAATTACTTCTCCCA	143
HHV7_RK_U12	TATCTTATCACACTCTTTATAGTGATTCTTTACATTGACTGTGTTACTTCTCACA	360
A10_020413	TATCTTATCACACTCTTTATTAGTGATTCTTTACATTGACTGGCTGTTACTTCTCACA	203
HHV7_RK_U12	ттдаассдадаадстстсасааассттаатсаддетттдтдтсаатдтдтдстттттдта	420
A10_020413	ттдаассдадаадстстсасааассттаатсаддетттдтдтсаатдтдтдстттттдта	263
HHV7_RK_U12	TACAGTGCGTCCTGCACATACAGTCTGTGCATGCTAGCAGTAATATCCACAATACGCTAT	480
A10_020413	TACAGTGCGTCCTGCACATACAGA	287
RHV7U13		
HHV7_RK_U13	ATGACTTTGAATCAACAGCCATCTAATTGCAGACTAATTACAGCTAACGATCCAGTACTT	60
A11_020419		40
HHV7_RK_U13 A11_020419	GCATCGAATTTTACAATGCAGCCTACTTTTAAAATAGCCGATAAGAAAGTTG ACATGAAATCTGGCTTAATGGAATATGCTCTAATGCTATCGGATGTCTTCTTTTCAATTT *** *** * * * **** * * *** * * * * *	112 100
HHV7_RK_U13 A11_020419	ТАСТСАВАВАТСАТААТТАСАТТВСАВТСАААВАТТТТЭТТСТАТСААВЭТСТТТСА-ТБ ТТТССТВАСАТСВААТЭСААТЭСА-ТБАААВАССТТВАТАВААСААААТСТТТВАСТБ * * * * * * * * * * * * * * * * * * *	171 157
HHV7_RK_U13 A11_020419	CATTOCATTCGATGTCAOGAAAAAATTGAAAAGAAGACATCCGATAGCATTAGAGCAT CAATGTAATTATGATCTCTGAGTACAACTTTCTTATCOGCTATTTTAAAAGTAG ** ** * * *** *** *** *** *** ***	229 211
HHV7_RK_U13 A11_020419	ATTCCATTAAGCCAGATTTCATGTTTTATGACTCAGACGAAGTTCGTTC	289 267
HHV7_RK_U13 A11_020419	TTTTATAA 297 TTAGATGGCTGTTG 281 ** **	
HHV7U14		
HHV7_RK_U14	ATGGAACAGCAAAAAGGATTTTCGATTCCATTTTTTGTCACTGAAGGAAAACTGCAACTTT	60
A12_020308		17
HHV7_RK_U14	GTGCCTGAAATATTACCTCGTATACATACTAAATTTCTTAAAG-ATGTCTTAATTGCC	117
A12_020308	GTGCNNAAAANNATACCTCGTATACAAAANNAAAATTTCTTAAAGGATGTCTTAATTGCC	77
HHV7_RK_U14	GATTCCTACAACTCTGTTAGTTGGGCAAACAGTTTATTCCCATGCC-TATTCAA-ACGC	175
A12_020308	GATTCCTACAACTCTGTTAGTTGGGCAAACAGTNNGATTCCCATGCCGTATTCAAGACGC	137
HHV7_RK_U14 A12_020308	ТТСААСААА - ТТАТОЭТТСТТАТААСАААGTTCAAGTTTTCCCCCCCCGTGATTTTTTA ТССААСААЛАТТАТССТТАТААСАААGTTCAAGATTTCCCCCCCCGTGATTTTTTA * *****	234 197
HHV7_RK_U14	TTCCCAGTAATTCGATTAGCTGTTCATATCA-ATAGGTTCCACACAGGGAAGAAACA	290
A12_020308	TTCCCAGTAATTCGANGGGNNGCTGTTCATATCNCATAGGTTCCACACAGGGAAGAAACA	257
HHV7_RK_U14 A12_020308	GCTGAAAACCA-TGATAGAAATTATGAAAAGCTTGTTTAACACCGAAGAG-GCTATG-CG GCTGAAAACCNATGATAGAAATTATGAAAAGCTTGTTTAANNCCGTTGAGTNNNANGTCG *******	347 317
HHV7_RK_U14	ACGATTCGATGAAGCATTGATGATTCTATTTTCTAATGAGCAAACCAATACTTACATGAC	407
A12_020308	ACGACCC	324
HHV7U15		
HHV7_RK_U15	AAATTTTGAAACTOGTTTGTGTAGAAATAATAGAGGTGGACTCACOGTTTTACAGGTATG	420
B1_020419		1
HHV7_RK_U15	TGTGTGRAATTGAGC-GACGATATAACAATCCTCAGTTTTGTGGACATCTAGATTGAAAAT	479
B1_020419	TGTGTGRAATTGAGCTNACGATATAACAATCCTCAGTTTTGTGAACATCTAGATTGAAAAT	61
HHV7_RK_U15 B1_020419	СТССТТСАТТТСТГОССАТСТСТТААТААТАААСАААССТТСААТСАСТАСАДАСТТТСЯ СТССТТСАТТТСТГОССАТСТСТТААТААТАААСАААССТТСААТСАСТАСАДАСТТТСЯ 	539 121
HHV7_RK_U15 B1_020419	САСАБЛТЮТОБОТТАЛТОТТАЛТССАВАЛАЛАЛАЛАТАЛТСТАТАТОТТОТАТТАЛАЛТССАТ САСАБЛТЮТОБОТТАЛТОТТАЛТССАВАЛАЛАЛАЛАЛТСТАТАТОТТОТАТТАЛАЛТССАТ ****	599 181
HHV7_RK_U15	TTGAAAAAGATATTTTATATAGATTGTCTCCCACTTTAGA-TATAATATTA-GATAAC-GT	656
B1_020419	TTGAAAAAGATATTTTATATAGATTGTCTCCCACTTTAGNATATAATATTAAGATAACCGT	241
HHV7_RK_U15	CATCAATATCTGTAGIGGACACAGCTCGCGAAATTCCTGTAGTCGTTGTCTTCCCAG	714
B1_020419	CCATCCAATATNC-TNGTAGNTGGGACCCCAGCCTCGGCGAANAA	285

. .

нн	v	7	U	1	7
	٠		•	_	

HHV7U17		
HHV7_RK_U17	ACGCTCTTCTTCCCGAGAAGAATGATGGTTTTAGCAGCTGCTGTTGTTACGTCGTTTTT	720
B2_020424		27
HHV7_RK_U17	AGCACTCAGAAAACCGATGACTTGTAACCTTTCATTGCAACATAGGAGATAGCAACCTAT	780
B2_020424	AGCACTCAGAAAACCGATGACTTGTAACCTTTCATTGCAACATAGGAGATAGCAACCTAT	87
HHV7_RK_U17	GAAAACAACAAAAACTTATTAGAACATCATGACACACATTGGAAATGTTTTGACTGCTTTTG	840
B2_020424	GAAAACAACAAAAACTTATTAGAACATCATGACACACATTGGAAATGTTTTGACTGCTTTTG	147
HHV7_RK_U17	CAACTTACATAGGTCTTGCAGCTCGGTTTCACTAGTAGTTCTGAAAGGCATATCCCGCAA	900
B2_020424	CAACTTACATAGGTCTTGCAGCTCGGTTTCACTAGTAGTTCTGAAAGGCATATCCCCGCAA	207
HHV7_RK_U17	AGGCGAAATGATTATTTTCCTACTCTGAGGCCATCTTATGAAGATTTCACACCCACTATT	960
B2_020424	AGGCGAAATGATTATTTTCCTACTCTGAGGCCATCTTATGAAGATTTCACACCCACTATT	267
HHV7_RK_U17	тталсталсалатсясталаадалтсасаяталлассттаттталсаласстталлтт	1020
B2_020424	тталаталсалатсясталаадалтсасаяталалассттаттталсаласстталатт	327
HHV7_RK_U17 B2_020424	TGCGAGTGCATGCCTAAATCGAACTTTTTTGTTCACTTTCGGCCAT 1065 TGCGAGTGCATGCCTA 343	
HHV7018		
HHV7_RK_U18	AGTGATGGAACCACCGGAATTATGACCTTCTAACCAACATTCAAGTGTACTGTTAGCCCG	540
B3_020424		11
HHV7_RK_U18	Адаадсаладтоссатттттодсааттадтостсттаасатаастостстааадтаата	600
B3_020424	адаадсааадтоссаттттттодсааттадтостсттаасатаасттстстааадтаата	71
HHV7_RK_U18	ATTGATCAGCACTCTCRAATAGGTTAATTTATGAGATGTTGTAGTAAGATTTACTAGACT	660
B3_020424	ATTGATCAGCACTCTCRAATAGGTTAATTTATGAGATGTTGTAGTAAGATTTACTAGACT	131
HHV7_RK_U18	CATTTCGCGAAATCCTTCAGTTTTTAAAGACAAAGAAATCAGAAAAGCTGTTCTTACAGA	720
B3_020424	CATTTCGCGAAATCCTTCAGTTTTTAAAGACAAAGAAATCAGAAAAGCTGTTCTTACAGA	191
HHV7_RK_U18	TAAACAACTAGAATTCCCTATTAGCATTTCATTAACATAACACGTGGTGACACAAACAGC	780
B3_020424	TAAACAACTAGAATTCCCTATTAGCATTTCATTAACATAACACGTGGTGACACAAACAGC	251
HHV7_RK_U18	TTCTAATTTATTATCTAGAAATATTATATTCTGGGGACATTTCAAATTTGACATGAGGGG	840
B3_020424	TTCTAATTTATTATCTAGAAATATTATATT	302
HHV7U19		
HHV7_RK_U19 B4_020413	CTGAGACAAGGACAGTAAAAAAACTATTTGTCTTCGCGGGGTCCCAGAAGCTGACACCCA TAAGCTGACACCCA ***********	720 14
HHV7_RK_U19	TT-CTTCGGCAACCCCAACATCACCATCTTTCCCTCAAATGGATAAAATCCGATGTTAA	779
B4_020413	TTTCTTCGGCAACCCCCAAACATCACCATCTTTCCCTCAAATGGATAAAATCCGATGTTAA	74
HHV7_RK_U19 B4_020413	СТАЛАССТТТАЛСТСТССТТТСТССАЛАЛАЛАТСТСАТАТСАСАЛАТТССАЛАТАТСА СТАЛАССТТТАЛСТСССТТТСТССАЛАЛАТССАСТСАТАТСАСАЛАТТССАЛАТАТСА СТАЛАССТТТАЛСТСССТТТСТСССАЛАЛАТССАСТСАТАТСАСАЛАТТССАЛТАТСА	839 134
HHV7_RK_U19	TACACATTATGTGTTGAATGGACACGCCATCCTTCATTATTTCTTGAACTGTTTTTTCAA	899
B4_020413	TACACATTATGTGTTGAATGGACACGCCATCCTTCCATTTT	175
HHV7U20		
HHV7_RK_U20	AATGAATTCAGCGTTAAAATTGTGGTGATAATGAACACTGCGATTCCTATGTGACCAG	898
B5_020502	TTCNNCGTTAAAATTMOTNOGNGATAATGAACACTGCGATTCCTATGTGACCAG	54
HHV7_RK_U20	асалалталдадалалагдстоттталтдтадаттсотссттоттодтадалалдасстал	958
B5_020502	асалалталдадалалтостоттталтдтадаттсотссттоттодтадалалдасстал	114
HHV7_RK_U20	Gacaaagcattaaatttcgttggaaaatctgtgaggaaggcgaaagacgaaatactagtattgccg	1018
B5_020502	gacaaagcattaaatttcgttggaaaatctgtgggaagggaagggaagggaattgccg	174
HHV7_RK_U20	ACAGACAGATTCTTATAATGTGCCAGTTCATAGTTATCCATCAATTCATATTTCCA	1078
B5_020502	ACAGACAGATTCTTATAATGTGCCAGTTCATAGTTATCCATCATCATATTTTCCCA	234
HHV7_RK_U20	АGАТGITCTITAITGATTAITGACAITGICA-ACTITAIGCCACAITAITTGGTAAAAIG	1137
B5_020502	AGAIGITCTITAITGATIAITGACAITGICAIACITTAIGCCACAA	281

HHV7U21	
HHV7_RK_U21	GGGTGAACAGATAATTTGAATATGAATGTAAGACTCGACATCAAGACAGAGTTTATCATA 960
B6_020413	T 1
HHV7_RK_U21	ТСЭСТ-ТСАТСАЭТССТТТКЭТТТТТТАААЭТТСТТАААСААААААТЭАТТАЭТЭЭ 1019
B6_020413	ТСЭСТЭТСАТСАЭТССТТТКЭТТТТТТАААЭТТСТТАААСААААААТЭАТТАЭТЭЭАЭ 61
HHV7_RK_U21	ARANTACTTGATARACAGTTGARAGCCARATATTTGCCARAATCGGTTTCCARTTGCGTTT 1079
B6_020413	ARANTACTTGATARCAGTTGARAGCCARATATTTGCCARAATCGGTTTCARTTTGCGTTT 121
HHV7_RK_U21	CGCTGTTGCCAAAAAAGAAAGCATTGTCTGTCATTGCCGCGACTGCAGAGTGTCTGATTT 1139
B6_020413	CGCTGTTGCCAAAAAAGAAAGCATTGTCTGTCATTGCCGCGACTGCAGAGTGTCTGATTT 181
HHV7_RK_U21	TATCGTCTGAACATTGTTTGCTCAAAGATATGTCGAACAATAGCAAATCATCCACCGTGTAG 1199
B6_020413	TATCGTCTGAACATTGTTTGCTCAAAGATATGTCGAACAATAGCAAATCATCCACCGTAG 241
HHV7_RK_U21	CATTCACATCAAAATCCACAACTGCTAAGGGACAGAAATCCGGATAGAGTTCTCCCATAAA 1259
B6_020413	CATTCACATCAAAATCCACAACTGCTAAGGGACAGAAATCCGGATAGAGTTCTCCATAAA 301
HHV7_RK_U21	TCACAGGAACGCAAAAACAGCAAGATAGTCCACAT 1293
B6_020413	TCACAGGAACGCAAAAACAGCAAGATA 327
HHV7U23	
HHV7_RK_U23 B7_020413	АСТААСТЭСТЭСААТТААААТСАСААОЗААСААААСЗААСТТТТТТАААЭЗЭСТАТТСАА 180 ТТЭСТЭСААТТААААТСАСААЭЗАСАААААСЗААСТТТТТТАААЭЗЭСТАТТСАА 55 **********************************
HHV7_RK_U23	AAGTTGAGGTTTGTCAGTTCTGGAGTTGAAATTGTCAGTGGAAAAATTCAGATGCATT 240
B7_020413	AAGTTGAGGTTTTGTCAGTTCTGGAGTTGAAATTGTCAGTGGAAAAATTCAGATGCATT 115
HHV7_RK_U23	CCCACTGTTTTCATGATAGTAGTAGTGCCACTGGAGAATGTCGAGGTCTTTGTAAAATT 300
B7_020413	CCCACTGTTTTCATGATTAGTAGTAGTGCCACTGGAGAATGTCGAGGTCTTTGTAAAATT 175
HHV7_RK_U23	TAAGGATGTAGATTCGACATTCTCTTTAGTTGCGATAGTCGTAGGTATACTCATAGA 360
B7_020413	TAAGGATGTAGATTCGACATTCTCTTTAGTTGTGCGATAGTCGTAGGTATACTCATAGA 235
HHV7_RK_U23	САСАДАСАТТАЛАТСССТСЭТАСТСТСТАТАТСТДАТТДАСЭТДДААСТААЛАТС 420
B7_020413	САСАДАСАТТАЛАТСЭСТСЭТАСТСТСТЭТАТСТДАТЭТДАТТДАСЭТЭДААСТААЛАТС 295
HHV7_RK_U23	TGAAGTTCTTGACATTTCCAGAGTGGGGCGTACTAGAAACAGTACTCATGTTAACAAAGGT 480
B7_020413	TGAAGTTCTTGACATTTCCAGAGTGGGGCGTACA 328
HHV7U24	
HHV7_RK_U24	ТСАООСТГООТАСССССАТАТСОААСАТТСААОАТОГОТАОААТААОААТСАОАСАТАА 60
B8_020424	ТТТСОТАСССССАТАТСОААСАТТСААОАТОГОТАОААТААОААТСАОАСАТАА 54
HHV7_RK_U24	GATAACCGATATTATGAGACATGCCAGTATTATAAAAGTCCCTCTTCGGGCGCACATTTTT 120
B8_020424	GATAACCGATATTATGAGACATGCCAGTATTATAAAAGTCCCTCTTCGGCGCACATTTTT 114
HHV7_RK_U24	GATTTCAGAACTAGAAGTTGAATTTATTGTTCTTGGTGATAGATTTTCCTGATGCAAAA 179
B8_020424	GATTTCAGAACTAGAAGTTGAATTTATTGTTCTTGGTGATAGAATTTTCCTGATGCAA 171
HHV7U24A	
HHV7_RK_U24A	CTAGGGATATTTAGAGCACCTGATAAATGATAACATATTCTCTG-CTATTG-AAGGACT 57
B9_020314	-TAGGGATATTTAGAGCACCTGATAAATGATAACATATTCTCTGTCTATTGTNNAGGACT 59
HHV7_RK_U24A	ТАТАТА-АТСАТGАТАТGСААТGСААGCCAAAATAAACAGAACAAACAAATAATAAATCC 116
B9_020314	ТАТАТCCATCATGATATGCAATGCAAGCCAAAATAAACAGAACAAACA
HHV7_RK_U24A	CTTTCCAACAATCTTCATAGGTCTCTGGGAAAAAATGGTTGTACTTATAAAACAT 171
B9_020314	CTTTCCTACAATCTTCATTTTTTTTTTANANTCTTTTTTINNCATNATTTATCCNTC 176
HHV7_RK_U24A B9_020314	CTCGTGCNNCTTGNNNTTTTTTCTNATTTINTGTTGTCCNACA 219
HHV7U25	
HHV7_RK_U25	AAAACCATTGTAGACATACAAATGCCCTCTGTCACCTTGTAAAAGAATTGGTCCTCTGCA 720
B10_020419	
HHUT DE HOE	

 HHV7_RK_U25
 AGTTGATTTTTTAACATG-ATGTAACACCAC----CAACAACGATCAACGTTTTTTCACA 774

 B10_020419
 GATNGGATTTTTTAACATGTGATANACACCCNCANCAACGATCAACGTTTTTTCACA 62

HHV7_RK_U25 B10_020419	ACACAAGTAAGTGTTTCGAATTTCTTCGATGTCTTCCTTGAAATTTCTGGGATTGTTTC 8. ACACAAGTAAGTGTTTCGAATTTCTTCGATGTCTTCCTTGAAATTTCTGGGATTGTTTC 1.	34 22
HHV7_RK_U25 B10_020419	TAATGAGTGAAATGTAACTCGCCATTCTTCGGGAAAGGCAATACGAAAGATGTACCTTC 8 TAATGAGTGAAATGTAACTCGCCATTCTTCGGGAAAGGCAATACGAAAGATGTACCTTC 1	94 82
HHV7_RK_U25 B10_020419	CTTACTTAGATGAATTCTTCGAGAAGAGAAGATAAGTTTATTGACACAGCACAAATTCGCT 9 CTTACTTA-GATGA-TTCTTCGAGA-GAGTACGTNTCNCGCCNCAGCNCNNNNCTTCGCT 2 *******	52 39
HHV7_RK_U25 B10_020419	AAGAAATCCAT9 NCATCNNAATTCCCGCNGGCCNGCCNNTGGGCGGGCCCGGGGGGGCNNTGCGGACGTCGNG 2 *** *	63 99
HHV7026		
HHV7_RK_U26 B11_020424	ATAAGATGCCATTGAATGCGTTG-GAAATATTAGCATGCATCTACAGACTATGGCTCTCA 5 TCCATTGAATGCGTTGTGAAATATTAGCATGCATCTACAGACTATGGCTCTCA 5	99 3
HHV7_RK_U26 B11_020424	АССССАТТТСАСАЛАЛАЛАЛАТАЛАЛАССАЛСАСАТТАССАЛАЛАЛАТТСАТССОТСАТТТ 6 АССССАТТТСАСАЛАЛАЛАЛАЛАЛАЛССАЛСАДАТТАССАЛАЛАЛАТТСАТСССОТСАТТТ 1	59 13
HHV7_RK_U26 B11_020424	GCAACACGAGTAATGATAAAGTTATAATTTTTGAGATCGTTGAAAGGACAATAGTTAAACA GCAACACGAGTAATGATAAAGTTATAATTTTTGAGATCGTTGAAAGGACAATAGTTAAACA 	19 73
HHV7_RK_U26 B11_020424	АЛТТСАССССИЛСЕСТАЛЯГСАЛТАССАЛАССТССССТАЛАЛАССАЛАТТАТСАСТАСТТ 7 АЛТТСАССССИЛСЕСТАЛЯГСАЛТАССАЛАССТССССТАЛАЛАССАЛАТТАСАСТАСТТ 2	79 33
HHV7_RK_U26 B11_020424	GTTCAAACAGCGTTTTCTTTATCATAAGATGGAA-ACAGCTTTCTCCCGGAATTACATTC 8 GTTCAAACAGCGTTTTCTTTATCATAAGATGGATTACAGCTTTCTCCCCGGAATTACATTC 2	38 93
HHV7_RK_U26 B11_020424	CCTABARCARGGCCCATAGARARACTTATCATTACATTCCACAT 882 CCTARARCHNGGCCCA 309	
HHV7027		
HHV7_RK_U27 B12_020424	САТААААСТАТСАААТААТGGAACAAAGTTATTTATCGTTTTAGTCGAAAAATGATCTGT 8 	40
HHV7_RK_U27 B12_020424	атсерттатататаласаттсте-сатераталтсяе-сттталалссалатертттт 8 тсерттатататаласаттстерсателаталтсяесттталалссалатерттттт 6 ,	98 5
HHV7_RK_U27 B12_020424	Ссадаттсялттатсяттсясссосттсяссаяталалоталстотастатттстссссала 9 Ссадаттсялттатсяттосссооттсялодотасялоталстотастатттстссссала 1	958 125
HHV7_RK_U27 B12_020424	AGCTTIGCAAAAGCTCTTAGAGGGCTTATTGATTGTTTTCCAAGTTTTCATGTGAAAAG 1 AGCTTTGCAAAAGCTCTTAGAGGCTTATTGATTGTTTTCCAAGTTTTCATGTGNAATAAG 1	L016 L85
HHV7_RK_U27 B12_020424	CCATGGTGGGAGGCTCTTTGGAATCTCGATGTTCGCGA-TGATC-TCGGTGAC-TAT 1 CCGAGTGGTGGGAGGCTNNTTGGAATCTCGATGTTCGCGNATGATCGTCGGTNACGTAT 2	L070 245
HHV7_RK_U27 B12_020424	GGTGTTCTCTGTTGCTACGATCCAT- 1095 GENGTATCGTCGTNNTGTCGTACGAA 271	
HNV7028		
HHV7_RK_U28 C1_020228	ACTITCTAAAATAAGTGGGATGGAATCAAATTCCGAAGCAGAGTTTGTTT	2160 45
HHV7_RK_U28 C1_020228	CTGGCCTAACAAACGATCAACGTGCGGTGCATCATGCATG	2220 101
HHV7_RK_U28 C1_020228	GGCATCACGTAAGCACCACATATCTAAGCCCGAATTTAAGCGCCGGAGAATTTTTGAAAT GGTGNGGCG-ATTTANTGTAGATCTTGGNTCNTNTTANTANATANGTANAT ** ** * * * * **** * ***	2280 151
HHV7_RK_U28 C1_020228	TTCATATAACTGCTTATAAAGAAGAGTTTTTGTGAAGGAGCGTGAATAGCAAAACCGTTT GTAGTTTNTTTGTATGCATTTNGTCNGTGTANATAGCAAAACCGTTT * * * ** ** ** ** ** ** ***	2340 202
HHV7_RK_U28 C1_020228	TTTGTTATATTCATCATGACAAAGACATTTCCTCAAGAGACCCATGTTTTTATTTA	2400 253
HHV7U29		

HHV7_RK_U29 C2_020424	AAACAAAGACACCTTTTCTTCTAAGTAAGAACAACTATCTAACATTGTGTTGATACTGTC AAACAAAGACACCTTTTCTTCTAAGTAAGAACAACTATCTAACATTGTGTTGATACTGTC *******	540 65
HHV7_RK_U29	AACTAATGAAGTTTTATACAGATTAGCTAAAGAAGTCCTCOTGCAGAAAGACACATTGAA	600
C2_020424	AACTAATGAAGTTTTATACAGATTAGCTAAAGAAGTCCTCOTGCAGAAAGACACATTGAA	125
HHV7_RK_U29	GGCATTCATGGCTAAAAACAATACAACTAATCAACAAAAATTTTGGACTCCCGCGTCTGT	660
C2_020424	GGCATTCATGGCTAAAAACAATACAACTAATCAACAAAAATTTTGGACTCTCCGCGTCTGT	185
HHV7_RK_U29	TTCAAGTCTGAACATAAGCATTCCTAAATGGTCTCGTTCAGGTCCTAAAAGTTCGCAAGG	720
C2_020424	TTCAAGTCTGAACATAAGCATTCCTAAATGGTCTCGTTCAGGTCCTAAAAGTTCGCAAGG	245
HHV7_RK_U29	AGTAAAACTTCGCGAAAGCACACACACTCTTTGGGAACACCAAAAGCTAGCAAACTTCTGAT	780
C2_020424	AGTAAAACTTCGCGAAAGCACACACTCTTTGGGAACACCAAAAGCTAGCAAAA	298
HHV7U30		
HHV7_RK_U30	GATAATACATTCTTAACTATAACAGCGAAAATTGAAATTTTTTCTCTTGCGGCTGAAAGT	120
C3_020413		13
HHV7_RK_U30	ATCACAGCAGATAAGATATTGTCATTTGCGAAATTATTACCAATACACAATTATCATTTC	180
C3_020413	ATCACAGCAGATAAGATA	73
HHV7_RK_U30	AACTITATTAAAAATCATATGGTGTTTTTATATATATAAATTATAGTACTCTAGGATTTGCC	240
C3_020413	AACTITATTAAAAATCATATGGTGTTTTTATATATAAATTATAGTACTCTAGGATTTGCC	133
HHV7_RK_U30	AAAAGATTTTCAATAGCTGAAACACTATGTAGAGAATTAAAAAATTTTCCACAGAACATTC	300
C3_020413	AAAAGATTTTCAATAGCTGAAACACTATGTAGAGAATTAAAAATTTTCCACAGAACATTC	193
HHV7_RK_U30 C3_020413	СААТСААССТСТАААТТАСААААСТТАААТААТССАСКАСАСТТТАТСААТТТССААА СААТСААССТСТАААТТАСААААСТТАААТААТССАСКАССТТТАТСААТТТССААА 	360 253
HHV7_RK_U30	CTTATTGAATCCATTCAAGGATTCAAGGCGGAATTGGCATCGCCTATTGGTAAACTTCTA	420
C3_020413	CTTATTGAATCCATTCAAGTATTCA-GGCGGAATTGGCATCGCCTATTA	301
HHV7U31		
HHV7_RK_U31	ATGAGAATCATAGCAGGAAGTACAAATCAAAACGATCCTAAATACGGACCAAGAGCCGGA	60
C4_020502	TGCAGGAAGTACAAATCAAAACGATCCTAAATACGGACCAAGAGCCGGA	49
HHV7_RK_U31	AAGCAATGTATGTCAAATTGTTTTTCTTTCTTGCATACAGTTTATTTGAACGGAATAAAC	120
C4_020502	AAGCAATGTATGTCAAATTGTTTTCTTTCTTGCATACAGTTTATTTGAACGGAATAAAC	109
HHV7_RK_U31	AATGTGTTAAATAAAGAGTCTATTGACATAATCATGGAAAAATGGAGCATTATTGGATAAT	180
C4_020502	AATGTGTTAAATAAAGAGTCTATTGACATAATCATGGAAAATGGAGCATTATTGGATAAT	169
HHV7_RK_U31	ATCAGTACAACGACATTGAAACTCGAAACTGGCAATATCCCAGAATATCGATTTTTTCACA	240
C4_020502	ATCAGTACAACGACATTGAAACTCGAAACTGGCAATATCCCAGAATATCGATTTTTTCACA	229
HHV7_RK_U31	GAAATCCCAAAAAAATTAGTTCTAAT-TTTGGCGAAACAATACATGAATTATCTAGACC	299
C4_020502	GAAATCCCAAAAAAAATTAGTTCTAANATTTGGCGAAACAAA	271
HHV7U32		
HHV7_RK_U32	САЛЛАГТТАВАЛТТТТТОВАЛОТТВАТСТТТТТТТТТТТТТТТТТТТАЛЛАЛАЛАТСОВСАССА	960
C5_020419	Т	1
HHV7_RK_U32	AGTTATTATTTTCTAGCGGGATTAAAAAACGATCGAAACTGTAACAATAATTTTCTATCA	1020
C5_020419	AGTTATTATTTTCTNGCGGGATTAAAAAACGATCGAAACTGTAACAATAATTTTCCTATCA	61
HHV7_RK_U32	TGGCCATGGCCATAGCAAAGATAAAATTTCCACATGGAACTGTTAAAGAACACCCATGAA	1080
C5_020419	TGGCCATGGCCATAGCAAAGATAAAATTTCCACATGGAACTGTTAAAGAACAACCATGAA	121
HHV7_RK_U32	ATATATCTTGATATGTCAGGTTGCTTGCTCTATGAGCGTATCGCAGTGTGATGTCATTAC	1140
C5_020419	ATATATCTTGATATGTCAGGTTGCTCGCTCTATGAGCGTATCGCAGTGTGATGTCAA	178
HHV7 U33		
HHV7_RK_U33	TCCACAATTTGTTTCCAAGTAAGACGTTGAATTTTGTAAAGAGCCAGAATTTTTAGAACC	120
C6_020502		9
HHV7_RK_U33	ТТGCATTGCCTGTACAAAAATACAAATATGCCAAAAGCAAAAAGAAATTTTTTGTCATC	180
C6_020502	АТGCATTGCCTGTACAAAAATACAAATATGCCAAAAGCAAAAAGAAATTTTTTGTCATC	69
HHV7_RK_U33	ATCATTTTATTAAAAATAGCTTTAATCTGATTTCTTGAATCCAGATCACACTTTAACCAA	240

C6_020502	ATCATTTTATTAAAAATAGCTTTAATCTGATTTCTTGAATCCAGATCACACTTTAACCAA 129	
HHV7_RK_U33 C6_020502	АСТТТТАЛАЛАСТТССЭЭТАЭЛАЛТТАТСАЭ-ТТАЛАТСТТТАСТСАЛАЛАСТТТТТАТТ 299 АСТТТТАЛАЛАСТТССЭЭТАЭЛАЛТТАТСНЭЭТАЛАТСТТТАСТСАЛАЛАСТТТТТАТТ 189 	
HHV7_RK_U33 C6_020502	CGTTACGAACATCGATAATACGCAATTTTCAGATT-TGCTGACATCTCCAATAAGAAATA 358 CGTTACGAACATCGATAATACGCAATTTTCAGATTGCGACANNCCATAGAATACGCAAAA 249	
HHV7_RK_U33 C6_020502	ACTGCARATGACTGTACGAACCATGTTTACACAGCGGAGAAATCAGCCGACAAAGCAAAT 418 CGT 252 *	
HHV7U34		
HHV7_RK_U34 C7_020419	ANGAGCGATATACTCAGATTCACTCTCGGGACCAAAAGCCACAAATTCAACTATAAAAGC 360	I
HHV7_RK_U34 C7_020419	СВСАТСТСЭБТТТС-БТСТААТААТВСАСТТТБТСАБТАСАССАСТТБААБААБЭТТТАА 419 СВСАТСТСЭБТТТСТЭГСТААТААТВСАСТТТЭГСАБТАСАССАСТТБААБААБЭТТТАА 61	,
HHV7_RK_U34 C7_020419	ТАССТТСАЛТОТСАЛТСАЛАСАСАСАДАТАЛАСАСТСАЛТБАСАТТТААДАТОТТССАТТ 479 ТАССТТСАЛТОТСАЛТСАЛАСАСАСАДАТАЛАСАСТСАЛТБАСАТТТААДАТОТТССАТТ 121)
HHV7_RK_U34 C7_020419	СТТТСАСАЛТОТТСТСЯСТТТСТТОТТТТАТОССАВСССТОСАВАТААСАЛСАСАТОСААА 539 СТТТСАСААТСТТСТСАСТТТСТТОТТТАТОССАВСССТОСАВСАТААСААСАТОСААА 181)
HHV7_RK_U34 C7_020419	CAGCACCTCCCGTATTTTTAAAAATAAAATCTCGCTTCTTGTTTTGTCAAATCTTCCCAGA 599 CAGCACCTCCCGTATTTTTAAAAATAAATCTCGCTTCTTGTTTTGTCAAATCTTCCCAGA 241	•
HHV7_RK_U34 C7_020419	GACTCAGCAAATATTCCAGGCTATATGCAAAATCCAATTTTAAAAGTATATCACATAAAG 659 GACTCAGCAAATATTCCAGGCTATATGCAAAATCCAATTTTAAAAGTATATCACATAAAG 301)
HHV7_RK_U34 C7_020419	GAAATTTAGGATTTTTAGAAAAAAAAATTTTTATCTGTGACACGAAAGTCTGCGGGACCCA 719 GAAATTTAGGATTTTTAGAAAAAAAATTTTTTATCTGTGACACGAAAGTCTGCGGGAA 358) }
HHV7U35		
HHV7_RK_U35 C8_020419	AATTGAAGTGAGTAGATTTGATTTTTTTTTTTTTTTTT	,
HHV7_RK_U35 C8_020419	ТG-СТСТЮТGАТGТАААТААТТТТАААТАGGTTAAGTGTCTGTATCCAACATTAATAAAAA 176 ТGGCTNCGTGATGTAAATATTTTAAATAGGTTAAGTGTCTGTATCCAACATTAATAAAAA 117	57
HHV7_RK_U35 C8_020419	GTAGTCGGGCATCTAATGAAGGCAGCATTATATCGTACAGTTTTGGTAACATAGGGAGAA 236 GTAGTCGGGCATCTAATGAAGGCAGCATTATATCGTACAGTTTTGGTAACATAGGGAGAA 177	5 7
HHV7_RK_U35 C8_020419	AAGCTATTTCAATTTCAACTCOGGAAGATAATTGTTCATAAAGAATTTCTATATCTGTTT 296 AAGCTATTTCAATTTCAACTCOGGAAGATAATTGTTCATAAAGAATTTCTATATCTGTTT 237	5 7
HHV7_RK_U35 C8_020419	TGGCATTGTCGTTAGACAT 315 TGGCATTGTCGTTA 251	
HHV7U36		
HHV7_RK_U36 C9_020502	CCACCGCAGGCNGCCGCGAATTCACTAGAAAGGGATGGAACTCCGATTC-CTTTTCTATGAA 41 CCACCGCAGGCNGCCGCGAATTCACTAGTGATTTGGAACTCCGATNNTCTTTTCTATGAA 60	
HHV7_RK_U36 C9_020502	CTCAGAACTTTTTAATGAAATTCTTTTATATGCACACTTAGATTCAAGTGGAATAGATTC 10 CTCAGAACTTTTTAATGAAATTCTTTTTATATGCACACTTAGATTCAAGTGGAATAGAATC 120	1 0
HHV7_RK_U36 C9_020502	ТСАТСАТСТТААСАСАААТСССААТАСАСТТСААААТСАААТТААСТСССТТСАААААА	1
HHV7_RK_U36 C9_020502	ТТТАЛАТАТТGАЛДААСТGАЛАЛАЛАТТАСААСТGСТTТАЛАТАТTGАСААССGАТGСАА 22: ТТТАЛАТАТТGАЛДААСТGАЛАЛАЛАТТАСААСТGСТTТАЛАТАТTGАСААССGАTGСАА 24:	1 0
HHV7_RK_U36 C9_020502	TATATGTTCAATTATAAACATCTGTTTACGACATGAAACCGATAAAATGTGGATCTATGA 28: TATATGTTCAATTATAAACATCTGTTTACGACATGAAACCGATAAAATGTGGATCTATGA 30	1 0
HHV7_RK_U36 C9_020502	CTATGCTCTTTTGTGTTACAAATGTAATGCTGCACCTAGAACTCCCCTTGGCTGTCGTAAT 34 CTATGCTCTTTTGTGTTACAAATGTAATGCTGCACCTAGAACTCCCCCA 34	1 9
HHV7U37	!	
HHV7 PK 1137	<u>ል ጥር ል ጥጥ ል ጥጥ ር ል ል ርርጥ ል ር አርርጥ አር ር አርጥ አር ላን አር ለሆነ አር ለሆነ አር ለመስለመመካ እ እ እ እ እ እ እ እ እ እ እ እ እ እ እ እ እ እ </u>	
C10_020424	ALGALIANI LAAGAAGAACAGAGAAGAAGAAGAAGAAGAAAAAAAAAA	

HHV7_RK_U37 C10_020424	ССТТАСААСАЛАБАЛАЛАСТААСТТАТСТТАТСТТТОТСАСТТАЛАБАЛСТССАТТСС ССТТАСЛАСЛАЛБАЛАЛАСТАЛСТТАТСТТТАТСТТТОТСАСТТАЛАБАЛСТССАТТСС СТТАСЛАСЛАЛБАЛАЛАСТАЛСТТАТСТТТОТСАСТТАЛАБАЛСТССАТТСС	120 106
ИНV7_RK_U37 C10_020424	GTTTTCAAATTATTTCCCAGAGTATGAATTGAAAATTCTAAATATGAAGAACTTCCCAATA GTTTTCAAATTATTTCCCAGAGTATGAATTGAAATTCTAAATATGATGAAACTTCCCAATA	180
HHV7_RK_U37 C10_020424	ACCGGTAAAGAACCTATCAAAATTCCATTCGATCTAAGTCTACATCATCAACATACGTGC ACCGGTAAAGAACCTATCAAAATTCCATTCGATCTAAGTCTACATCATCAACATACGTGC	240 226
HHV7_RK_U37 C10_020424	TTAGACTTATCACCATATGCCAATGAACAAGTTTCAAAAAGTGCATGTGTAAATGGCG TTAGACTTATCACCATATGCCAATGAACAAGTTTCAAAAAGTGCATGTGTTAATTGTGGT	300 286
HHV7_RK_U37 C10_020424	асласаласаттесьлеасттеляттегладатестатодтоделтататалателлаттеллас леалелласаттеслаелосттелятостатодтодела	360 326
HHV7038		
HHV7_RK_U38 C11_020424	ATTACCAGATGGAATAACGAAATGTCTATATTGGAAAGGGATGTTTTCTATAGAATCAGT TACCAGATGGAATAACGAAATGTCTATATTGGAAAGGGATGTTTTCTATAGAATCAGT	2760 58
HHV7_RK_U38 C11_020424	AAATAATAAAGATTCGGATGCATCATAGATATGAAATTTTTAAAGGCGCACGCA	2820 118
HHV7_RK_U38 C11_020424	TTCGGCTACTTGAGAAAGCGATGGCCATGTCATATCATTTTTGAGAATGTATTGCTTATC TTCGGCTACTTGAGAAAGCGATGGCCATGTCATATCATTTTTGAGAATGTATTGCTTATC	2880 178
HHV7_RK_U38 C11_020424	TTGATAGAACATTCTCGGTTCCGAGTCACAAAGAGTCTTCATTAAGCCGGGTGCTCCATC TTGATAGAACATTCTCGGTTCCGAGTCACAAAGAGTCTTCATTAAGCCGGGTGCTCCATC	2940 238
HHV7_RK_U38 C11_020424	GTGCATTATACCGCGCGGAAAAATCCGGAGAAAAGTAGATTTGTTTTTTTCTTAGTTCG GTGCATTATACCGCGCGGAAAAATCCGGAGAAAAGTAGATTTGTTTTTTTT	3000 298
HHV7_RK_U38 C11_020424	TACATTCTCTAGATACGGATTAAAGAAGGACACCAGATCCAT- 3042 TACATTCTCTAGATACGGATTAAAGAANGACACCAGATCCATA 341	
HHV7039		
HHV7_RK_U39 C12_020314	GTTTACCATGTGCACTTCTTCAATTGGCATTGGCAAAGTTGTAACAGTTCGATCTAAGAA 	2160 50
HHV7_RK_U39 C12_020314	ATACACAGTGCCAACATCGCGATATGTTGTTTGAAAGTTCTTTTTTTAAAAGTTCT NGGNTNOGTGGNANAATNNCTCTGGATTGTGTTGAATGTAGTCNNTTNNNGTGNNGG *** * ** * * * **** * ***** ****** **	2220 107
HHV7_RK_U39 C12_020314	GACAGAAAACGTGTGTGTCTTCAATCTTTGTTTTGTAAATATCAAAATACCTTCGGTAGT GACGANCGCTCNCTGTGCTTNTCTTTGTTTTGTNNTTTATNNNNCTTCGGTAGT *** ******* *************************	2280 161
HHV7_RK_U39 C12_020314	TTTAATGTTAGAGCCGTAAGACGCGCAAGAAACCTCCCTGTCAAAACGCACTAAATCTGT TTTAATGTTMGAGCCGTNNGACGCGCNNGAC-CCTCCCTGTCACTCGCCTACTTG	2340 215
HHV7_RK_U39 C12_020314	CCC60769CAATTGAACAAATTCGAAATGGTAAATGCTGATTGTGTCCAGTCATGACAAA TCC9G769CC-TTGAACCCTNNNCT9GTNT-TGCTG-TTTTGTCNGTCTGACCCCTC *****	2400 269
HHV7_RK_U39 C12_020314	GTCAGCTTCTGTTTGTAGAGATAGCTGTAAGCTAAAAGTTATAAAAACACTCAGGAATAG NCT	2460 272
HHV7U40		
HHV7_RK_U40 D1_020228	алдадссалдтссалттталдтогтосттосаттологтодалтататттсала 	1860 21
HHV7_RK_U40 D1_020228	ататттттс-тттстсаттатстаатдаалаттсаатсаалссатссаастаатааса ататтттттттстсаттатстаатдаалттсстсаатсаалссатссаастааладаа 	1919 81
HHV7_RK_U40 D1_020228	СТТСАТТАТАСАТАССТТТТААТАААТСТСТАААТАААТ	1978 141
HHV7_RK_U40 D1_020228	ТТАЛАЛТТТТСАТСАСАСАТСАССАЛАТАЛТСТТАЛАСАЛАТСАЛАТТАТАСААС ТТАЛАЛТТТТСАТСАССАСАТСАССАЛАТАЛТОГСТТАЛАСАЛАТСАЛАТТАТАСААС ******	2038 201
HHV7_RK_U40 D1_020228	ССЭТТТТГССТВАААТТЛОТСАТАТСТОЗААТААЭТОЛАСАЭЭЭССАСААААТСЭТААЭ ССЭТТТТССТВАААТТЛОТСАТАТСТОЭААТААЭТОЛЭАСАЭЭЭТСАСААААТСЭТААЭ ССЭТТТТССТВАААТТЛОТСАТАТСТОЭААТААЭТОЛЭАСАЭЭЭТСАСААААТСЭТААЭ	2098 261
HHV7_RK_U40 D1_020228	САСТСТААТТСТАААGCACATTCACTTAACCTTGAACACAAGACACAATGATTGCAAA САСТСТААТТСТАА-GCACATTCACTTAССТМАСАСААCACATATG	2158 310

HHV7U41	
HHV7_RK_U41	TCCTATATCCACAGGCTTTCGATTTCCGTTAACGACAAATTCTTGTAGATTGAATTTTTC 3060
D2_020424	TCAGGCTTTCGATTTCCGTTAACGACAAATTCTTGTAGATTGAATTTTTC 50
HHV7_RK_U41	ТСТТОТТТОТТОСАСААТТТАТСААGATTTCCATGGTCTTCTGCTTCTTTTAAAAAAAC 3120
D2_020424	ТСТТОТТТОТТОСАСААТТТАТСААGATTTCCATGGTCTTCTGCTTCTTTTAAAAAAAC 110
HHV7_RK_U41	ATCTGTACCGTGGAAAAAGAAGCATACAGGCATAAACGATGTGATTTTGGTAATGACTGT 3180
D2_020424	ATCTGTACCGTGGAAAAAGAAGCATACAGGCATAAACGATGTGATTTTGGTAATGACTGT 170
HHV7_RK_U41	TCCGCCGTAATTAATAATGGGTGTTTTAACGGTTGGGAAGAAGTCATTTTCAACCGTTAG 3240
D2_020424	TCCGCCGTAATTAATAATGGGTGTTTTAACGGTTGGGAAGAAGTCATTTTCAACCGTTAG 230
HHV7_RK_U41	ATTAAGTAACAAGGTGAGATCACTACTGATTTTCTTTTTTCCATTAACGATAATGTAGT 3300
D2_020424	ATTAAGTAACAAGGTGAGATCACTACTGATTTTCTTTTTTCCATTAACGATAATGTAGT 290
HHV7_RK_U41	CAGAATTTCTATTAATTTTTGTTCTTTCGGCAGAATGTACAGCCATGCGGGTACATAT 3360
D2_020424	CAGAATTTCTATTAATTTTTGTTCTTTCGGCAGAATGTACAGCCATGCTGCGGTACATAT 350
HHV7_RK_U41 D2_020424	TGGGGCCGAAACAACAGTTTCATTATCATCAGCCAT 3396 TGA 353 **
HHV7042	
HHV7_RK_U42	TGGATCTCGTGCTTCACCGGAATACTTGCGTAAACCAAGGGCTTCCGCGGTGTAGCTAGA 1260
D3_020413	TAAGGCTTCCGCGGTGTAGCTAGA 25
HHV7_RK_U42 D3_020413	GCCAAAAGGTGCTGTTGGTAAATTGGCGAAAAAAGTCAATGCATGC
HHV7_RK_U42	TTCTTCCTGCTTTCTTCAATGATGAGTTCAGCTGTTCTAGTTCTCTGAGACTCAAAGC 1380
D3_020413	TTCTTCCTGCTTTCTTCAATGATGAGTTTCAGCTGTTCTAGTTCTCTGAGACTCAAAGC 145
HHV7_RK_U42	GTCTAATTTTGATTCATCAAGCTGCTTAATAGGAATGATGCGTCTGTGAAAATTGTTTCGT 1440
D3_020413	GTCTAATTTTGATTCATCAAGCTGCTTAATAGGAATGGATGCGTCTGTGAAAATTGTTTCGT 205
HHV7_RK_U42	AAAGCGGCTGACATACTTGTTAGCAGGTTTATGAACAAGAGTTCGTTTAATGGTTTTCAA 1500
D3_020413	AAAGCGGCTGACATACTTGTTAGCAGGTTTATGAACAAGAGTTCGTTTAATGGTTTTCAA 265
HHV7_RK_U42	ACCATATCGTTGCCGCCCTAATACATTCTTTTAACCCCGCGTGGATACAT 1551
D3_020413	ACCATATCGTTGCCGCCCTAATACA 290
HHV7U43	
HHV7_RK_U43 D4_020313	CCARARCOOTGRCGTGGGTRAGAGATACTTATTTGTTAGATGTTTARTGATTATAGTCTT 2220
HHV7_RK_U43 D4_020313	GCAAAAAAAGACTAAGCTTTTAAATTCTATATTTTCACTTCCTTCTGGAACATGAGAA 2278 GCAGGNCCCGGCGCCAGGCGGCNGNGGGANNCGATTNCACTTCCTTC-GGAACAGAGAA 59 *** * * * **
HHV7_RK_U43	AGTAAAAAATCTTTATACAGATTTTCTGTTTCGTATTTAAACAGTGAATAATATCC 2338
D4_020313	AGNNAAAAAT-TCCNNNCAGANNCNNTNNNNNAAANANTAAACAGGAAANATCC 112
HHV7_RK_U43	TCAGACTTAATGGGTAATGCATTTTTTACATAATAGCTGAAATTCAAAGATTCTGCGTCG 2398
D4_020313	TCAGACT-AAGGGNNAGCANACANGANAGCGGAAATTCAAAGATTCGGNGTNG 164
HHV7_RK_U43	САЛАСАЛАТАССОТСТСАЛТТСТТТЮGAЛАЛСТТОСАТТТСТОТОТСТССАТАСАЛАЛА 2458
D4_020313	САЛАСАЛАТАССОТСТСАЛТТСЯGGAЛАЛСТ-GCATTCONGTCGCATACAЛАЛА 217
HHV7_RK_U43 D4_020313	TATACATTTTTTGATGGTTTATATTTAACGATTATTGGAAAAATTAAATGTTCACTAGGG 2518 TANG-ACANNNNGAGGNCTTAATTNACGATTATGGAANATTAAGNTCACNNGGN 270 ** * * * * * ** ** ** ** *** ** *******
HHV7_RK_U43	TTCTTACATAAGATATTTACTATCACGTGCGCGGGGGCATATTCCGTTGCAAACACAGTA 2578
D4_020313	TCTACATGA-ATACATNAGGTGGGGT-CAANANGCNACNNTCAGGNNGTCGCGCCCGTG 326
HHV7_RK_U43 D4_020313	ATAGTCAT 2586 CAGANAGTGCT 337 *
HHV7U44	
HHV7_RK_U44 D5_020413	ATGGGAAATTGCTTTGTAAAAAAAATCTCTTCTGACATCTTCAACTCTAACTACCAGATG 60 TCTACCAGATG 11 ******
HIR 77 DE 1144	ርማርጫ አመርባዊ ጋን በረማርጫር እርጉ በርጉ እርጉ እርጉ እጥበት እርጉ በማሪካ አመርካ በአስተ በእርጉ እንደ የሆኑ እንደ ማይም እንደ እንደ የሆኑ እንደ እንደ

HHV7_RK_U44 D5_020413	СТСТАТТСТВАЗСТСТСАВАЗСААВАЗДАТТТАСТОЗАТТТТТАВАЗАСААААТАТАСА СТСТАТТСТВАЗСТСТСАВАЗСААВАЗДАТТТАСТОЗАТТТТТАВАЗАСААААТАТАСА СТСТАТТСТВАЗССТССАВАЗСААВАЗДАТТТАСТОЗАТТТТТТАВАЗАСААААТАТАСА	120 71

HHV7_RK_U44 D5_020413	GATTTTGGAATTTTAAAAACCGATATTCTCAACTATGAAAGAGACTCCGAAACATTCAAA GATTTTGGAATTTTAAAAACCGATATTCTCAACTATGAAAGAGACTCCGGAAACATTCAAA *********	180 131
HHV7_RK_U44 D5_020413	астттуттусаастоттасстатстаталалаласаласстдасстдаталатттуаттуал астттуттусалотортасстатстаталалаласаласстудортаталттусаттусал ******	240 191
HHV7_RK_U44 D5_020413	СЭСТЭТТАЛАТАЛТЭГССССССТСАСЭТТАЛАЭЛЭСАТТЭЛТЭЛТЭЛТЭЛЭЛЭСССССССССС	300 251
HHV7_RK_U44 D5_020413	GCTAAGAAAATTTTAGAGACTCTGGATGTTGTGTTCATGAAAATTATGATTGGAGAATTT GCTAAGAAAATTTTAGAGACTCTGGATGTTGTGTTCATGAAAA ********************************	360 294
HHV7045		
HHV7_RK_U45 D6_020502	TTTTTCACAGGATTTGAATAAACCGTTTTGAGGAATT-CCAAGTAAAGAAAATGGTCGTT TCCGTTTTGAGGAATNTCCAAGTAAAGAAAATGGTCGTT *******************************	899 39
HHV7_RK_U45 D6_020502	GATAAACCGTCAATTCAGTGTTCACATATTCATTTAATTGCTTTGCATCGATTTTAGGCC GATAAACCGTCAATTCAGTGTTCACATATTCATTTAATTGCTTTGCATCGATTTTAGGCC	959 99
HHV7_RK_U45 D6_020502	ATGTAAGAGAGCTCTCTTTTGTTTCGCTAAAGTTCCAGATATCATCCACATCCGTTACAG ATGTAAGAGAGCTCTCTTTTGTTTCGCTAAAGTTCCAGATATCATCCACATCCGTTACAG	1019 159
HHV7_RK_U45 D6_020502	TAAACAGGAAAAAGTGCAAGTATCTGGACAATAGGCAATTTGTTAAGTGTAATTGTTTCAG TAAACAGGAAAAAGTGCAAGTATCTGGACATAGGCAATTTGTTAAGTGTAATTGTTTCAG	1079 219
HHV7_RK_U45 D6_020502	AGATTTTTTCAGAAAATTATTTTCTTOTTTGTTTCAGAACTTTCGGAGGGTAGACTTTG AGATTTTTTCAGAAATTATTTTCTTGTTTGTTTCAGAACTTTCCTGGAAGGGTAGACTTTG *******	1137 279
HHV7_RK_U45 D6_020502	CAT 1140 CNA 282 *	
HHV7U46		
HHV7_RK_U46 D7_020419	ATGACTCTATACAAGATTGTTTCCAAGCCAATTATATTGCTGGCATTTTTTTT	60 34
HHV7_RK_U46 D7_020419	GTTGTTTTCACAAATGAAGTCGACGGGGGGGGGGGGGGG	120 94
HHV7_RK_U46 D7_020419	ĠĂTĂĊĂŦĂŦĞĂĂĂŦŦĂŦŦŦŦŦĂĂĂĂĂĂŦŦŦŦĊŦĊŦĊŦĂŦŦŦĞĢĂŦĂŦŢĠĢŦĊĂĂĊĂĊĂĊŦŦŦ ĠĂŦĂĊĂŦĂŦĠĂĂĂŦŦĂŦŦŦŦŔĂĂĂĂĂĂŦŦŦŦĊŦĊŦĊŦĂŦŦŦĠĢĂŦĂŦŢĠġſĊĂĂĊĂĊĂĊĂ *****************************	180 154
HHV7_RK_U46 D7_020419	АТТТТАТАТУТСАТТТСАТТАТТТТАААGТАТТGGTGTTTTAAGACTCTC-GCCAA АТТТТАТАТССАТТСАТТАТТТТАААGTATTGGTGTTTTAAGACTCTCAGCCAA ********************************	239 214
HHV7_RK_U46 D7_020419	Адаластоталладодтаттал 261 Адаластод 223 ******	
HHV7047		
HHV7_RK_U47 D8_020502	GGGAAATTCAAAATGTAATGTCCATCGCGGAAAATTATCTAGATAGA	660 51
HHV7_RK_U47 D8_020502	Адаладсалататтсттттстдддатадасодалтдассалалталатстотттала Адаладсал-таттсттттстдддагадасодалтдассалалталалтстотттала ********	720 110
HHV7_RK_U47 D8_020502	ACAGTTCTTAGAGATGGGTATGTCGCAACTCTGAATATATGTTTTTTCAAAAGACTCATA ACATTTCTCAGAGATGGGTATGTCGCAACTCTGAATATATGTTTTTTCAAAAGACTCATA	780 170
HHV7_RK_U47 D8_020502	ATGAAATTTTGCGAATTTCGGCCACATGGAAAAATTCGTTAAATTTACAAACGAGATTTT ATGAAATTTTGCGAATTTCGGCCACATGGAAAAATTCGTTAAATTTACAAACGAGATTTT	840 230
HHV7_RK_U47 D8_020502	ТТСАGАGСАЛАСАСТТААТАСАGGAGTCCATATGATCAGATAACTATAAAAAAGAAAAGA ТТСАGAGCAAACACTTAATACAGGAGTCCATATGATCTTATAACTATAAAAAAAA	900 290
HHV7_RK_U47 D8_020502	AATCAACGTGAACACTAGCATTGAATACATTTTGTTTTCAT 942 AATCAACGTGAACACTAGCATTGA 314 ************************************	
HHV7U48		
HHV7_RK_U48 D9_020413	СТТТТСАТАЛАЛТАЛАGGTAATAAAGACACTGTAATCGTTTTTGTATGCGAAACAGGTTT 	1740 20

 HHV7_RK_U48
 GATATTTGTCTGAAATTTTTGTGCA-TAGATTTTCACAGCTTTCTCTGTTGATGGGTTAA 1799

 D9_020413
 GATATTTGTCTGAAATTTTTGTGCACTAGATTTTCACAGCTTTCTCTGTTGATGGGTTAA 80

HHV7_RK_U48	АЛАЛАССТСТАЛТСССАТСТТАТАТТСАТСАЛАТСАЛТСАТССАЛАТТСАСАСАЛАТСАЛ	L859
D9_020413	АЛАЛАССТСТАЛТСССАТСТТАТАТТСАТСАЛАТСАЛ	L40
HHV7_RK_U48	ATAAATTCGAGACAAAAGTATATCCGAATAAGCATTTGGGAATCTGATAAACCCGAGTTT 1	L919
D9_020413	ATAAATTCGAGACAAAAGTATATCCGAATAAGCATTTGGGAATCTGATAAACCCGAGTTT 2	200
HHV7_RK_U48 D9_020413	CGTTATAATCATGGAAATTAAACGTTATTAAACCAGGTTGAATAATAGT-TTGATTTGTT CGTTATAATCATGGAAATTAAACGTTATTAAACCAGGTTGAATAATAGNCNTGATTTGTT 	L978 260
HHV7_RK_U48 D9_020413	TTTTCATTTACGCAAATTGAAGAATTGAGTATATTCCAGGGTTTCCATCCA	2036 317
HHV7049		
HHV7_RK_U49 D10_020413	АТЭТСТТТАGААТАСТТЭССССССТЭТТАGААGАСЭТАТТЭЭТСААТААТААТСАТТТЭАGA ТАGААСЭСТАТТЭЭСССССТЭТТАGААСЭТАТТЭЭТСААТААТСАТТТЭАGA ************************************	60 34
HHV7_RK_U49 D10_020413	атстасалалалаттстоттотталаатссааттттолалалатталатт	120 94
HHV7_RK_U49 D10_020413	АЛТСТТТТТССАGААGAACTGCATGATTCAAAAATACATGTATATTTTGAAGTTAGATTA ААТСТТТТТССАGAAGAACTGCATGATTCCAAAATACATGTATATTTTGAAGTTAGATTA 	180 154
HHV7_RK_U49	GGGTGCCGTATTCCGGATTGCATTATTGTTTTTAGGCATTTTGGTGAGAAATTATTAAAA	240
D10_020413	GGGTGCCGTATTCCGGATTGCATTATTGTTTTTAGGCATTTTGGTGAGAAATTATTAAAA	214
HHV7_RK_U49	асттесатотаттттттораттсалаастастаттоссалатсялатсятотттса	300
D10_020413	асттесаттотататтттттораттсалаастастттоссалатсялатсяотттса	27 4
HHV7_RK_U49	атт-салалалатабдасасалалаттсалтаттасаддд - Асталдосалстасдас	358
D10_020413	NTTTCAЛАЛАЛТАБДАСАСАЛАЛАЛТСАЛТАТТАСАДДД - Асталдосалстасдас	334
HHV7_RK_U49	AAGCAACAGATTATCTACAGCAATTTGTCATCAAAAATGAAAGCTTATGTAAGGTAAACC	418
D10_020413	AAGCAACCAGAAATCACT	352
HHV7U50		
HHV7_RK_U50 D11_020424	АСТGААСТТGАТGTTTTGCAAGCACATGTACAGACAGAATGCCAGAAATTAAATACAAAT ТатGTACAGACAGAATGCCAGAAATTAAATACAAAT ************************	240 36
HHV7_RK_U50	ТТАЛДАДАСАТТДААААТGCATTGCTTTTAGAAAATCAAAAAATAATTCCTTCTAGTGAA	300
D11_020424	ТТАЛДАДАСАТТДААААТGCATTGCTTTTTAGAAAATCAAAAATAATTCCTTCTAGTGAA	96
HHV7_RK_U50	асасоттстотсттеблавартстстссавосталалстотсасосалотсаслаттаса	360
D11_020424	Асасаттстотсттеблавартстстссавосталалстотсасосалотсасалттаса	156
HHV7_RK_U50	CAAATTGATCCTGCTATTCATTTCACGGAAAATTTCCGACCTGAGATGATAAAAACTTTT	420
D11_020424	CAAATTGATCCTGCTATTCATTTCACGGAAAATTTCCGACCTGAGATGATAAAAACTTTT	216
HHV7_RK_U50	TATAATAATACTCAGATGTGGGGCTATACGTTTGGAGCTTGGTTTTACAAATTAAAGCGA	480
D11_020424	TATAATAATACTCAGATGTGGGGCTATACGTTTGGAGCTTGGTTTTACAAATTAAAGCGA	276
HHV7_RK_U50	GCTTTCTTTACCGATTCTAAATTAAAAAGAATGCTAAAGTTAACATATGTTGATTCTCTC	540
D11_020424	GCTTTCTTTACCGATTCTAAATTAAAAGAATGCTAAAGTTAACATATGTTGATTCTCTC	336
HHV7_RK_U50 D11_020424	ТСТАТТАСАСАБДААТТЭСТЭТСТАТТТСААТТААТЭСАТТЭДААСАААТТАСТАТТТАТ ТСТАТТАСАСАВДААТТЭСТЭТСТАТТТСААТТААТЭСАТТЭДААСАААТТАСТАТТТАТ *******	600 396
HHV7_RK_U50 D11_020424	CCTATGCATGATAATTTAGTGTCCGATTTAGAAGCTGGTTTGTGTCTACTGACTG	660 435
HHV7U51		
HHV7_RK_U51	TTTTTCTOGTTTTTCTTTCTTTGTCTGCTGGTAATAATAGTGGTGAAATTTAATAACAGC	120
D12_020314	TCTCGGNTTCNTNNNTTCNNGCTGG-AATAATAGTGGTAAANNAATAACAGC	51
HHV7_RK_U51	ACCGTTGGTCGAGAATACACATTTAGTACT-TTTTCTGGAATGTTGGTT-TATATTTTAT	178
D12_020314	ACCGTGGTCGGAGAATACACANN-AGNACTGTTTTCTGGAATGNGGGTTANANNNATNNN	110
HHV7_RK_U51	TGT-TACCGGTTAAAATGGGAATGTTAACCAAAATGTGGGATGTATCCACCGAT	231
D12_020314	NGAATACCGGTTAAATGGGGAATGTTAACGGCGGGAAATGTGGGANCTGRATCCACCGAT	170
HHV7_RK_U51 D12_020314	ТАСТС-ТАТААТСТТААТСТТТСТСАДТСАТТТТТСАТТТАТСТТТССОССТОЗСССОТ ТАСПИСТАТИАТСТТИАТСТТТСТСАСТСАТТИТСАТТАТСТТТТССОССТОСССОССОТ *** *** ****	290 228
HHV7_RK_U51	AACGCTGTTGGCACTGGAACGTATCAACAATTTTTCTTTTCTGAGATTAAGGTAAACGA	350

D12_020314	A-CGCTGTTGGCT 240
HHV7052	
HHV7_RK_U52	ТААТТТТТТССАДАДСТСААСТССАСТСТТСАСТСААДАТАТСТТТААСДАГТТСТ 478
B1_020228	
HHV7_RK_U52	GATCCAGATCTTTGTTTAATCATGCAAAAGATCTTAAACCAAAATTCATCAGGTACCTTT 538
B1_020228	GATCCAGATCTTTGTTTAATCATGCAAAAGATCTTAAACCAAAATTCATCAGGTACCTTT 107
HHV7_RK_U52	ССАТТСАСАА-ССАСТАЛАТСАСТСАСТСССАТАТТОТССАЛАСИТСТСТСААСАСНА. 597
B1_020228	ССАТТСАСАЛАССАСТАЛАТСАСТССССАТАТТОТССАЛАСИТСТСТСААСАСНА. 167
HHV7_RK_U52	GATTTGGATGCCGTGAGAAAACATTAGGGTAC-ACACATTCTGGATGATCAAAATCTCTT 656
B1_020228	GATTTGGATGCCGTGAGAAAACATTAGGGTANCACACATTCTGGATGATCAAAATCTCTT 227
HHV7_RK_U52	CTTGTTTTAGATTAAGTAAAGATTCACCGCTTACCAATTTCTTAGTGATATGTAATATATA 716
B1_020228	CTTGTTTTAGATTAAATNAAGATTCACCGCTTACCA-TTTCTTAGGTANATGTTAATTTA 286
HHV7_RK_U52	AATTOGAGGTTGAGTTGTTCGAGATTATGTATTGACCTACTTGTGTCAT 765
B1_020228	NOGROGAAGT-GCGTTGTTC-AGATCGAGNNT-GGGTTGCCGTCGC 329
HHV7U53	
HHV7_RK_U53 B2_020424	АТЭБАЛАСТЭТЭСТАЭТЭССАЭСТТССТАТЭТЭТЭТЭТЭТЭТЭЛЭЛЭЛЭЛЭЛЭЛЭЛЭЛЭЛЭЛЭЛЭЛЭ
HHV7_RK_U53	ААСТТТТАТСТОССААОДАОДАСДАТАСААДААДАААТСААТТССОДЛААТОДТТТДААТ 120
B2_020424	ААСТТТТАТСТОССААОДАОДАСДАТАСААДААДАААТСААТТССОДЛААТОДТТТДААТ 106
HHV7_RK_U53	АТТССАТТСЯАТАТАЛААССАСААТСВАЛАТССССТТАТАGGCACAGTCTCTTCTTTAGG 180
B2_020424	АТТССАТТСЯАТАТАЛАССАСААТСВАЛАТССССТТАТАGGCACAGTCTCTTCTTTAGG 166
HHV7_RK_U53 B2_020424	GATTTACAGCACGGTTTGTTCACGGTTGCCCGTGTTCAATCAA
HHV7_RK_U53	АТТАЛДАЛАЛАТАДСТІДТАЛАЛТСТАЛІСТІДАТАЛССАЛСАСІДДАЛАЛАЛАСТСТІДССА 300
B2_020424	АТТАЛДАЛАЛТАДСТІДСАЛАЛТСТАЛІСТІДАТАЛССАЛСАСІДДАЛАЛАЛАСТСТІДССА 286
HHV7_RK_U53 B2_020424	CCAGATCCCGAAATAGAGTGTTTAAATTCAATTTCCCAGGTTTGTCATTATCGAACAGG 360 CCAGATCCA 295 ******
HHV7053.5	
HHV7_RK_U53.5 B3_020413	ATGAATGGCTC 11 CCCCGGGNCCCCATTGGGGGCGGGCCCCGGGCCGGNGCNANTTTCGGGAATTTTAGGCTT 60 ** ****
HHV7_RK_U53.5	AGCTACTGGCAACTCAGCTAGTCTTCCCGCTCCCAGTTTTTCCGAATGTG 61
B3_020413	AAN-GTINTITTINCNNGCGGNCTTCCCCTTCCNAGGTTTTTTTTCCCGGAAAATGGTG 119
HHV7_RK_U53.5	ТТТТТТТАССGARAGATACАТТТТGĆTCCTTАСТGAA-TGCT 102
B3_020413	ТТТТТТТТПИТТИМСССGGAANAAGAATANCAATTTTTGGCTCCCTTAACTGAAATGCTTN 179
HHV7_RK_U53.5	ACAGCTGGGG-CCCAAAATAAAAACGTAACGCCAGCTGCTCCGATATTTAAAACTGAT 159
B3_020413	ACAGCTGNGGGCCCAAAATAAAAAACGTAACGCCAGCTGCTCCGATATTTAAAACTGAT 239
HHV7_RK_U53.5	GAATATATAACTCCGTACCCAGAATCTCTGAGCAGAATGGAATTAGGAATAGAAT 219
B3_020413	GAATATATAACTCCGTACCCAGAATCTCTGAGCAGAGTGGATTATGGAAATAGAATGAAT
HHV7_RK_U53.5	ТАТСАСАТАССАССТССАТАТТОСТАТСССТСТАТСССТОСАТТТААТТАТАААТСТТАТ 279
B3_020413	ТАТСАСАТАССАССТССАТАТТССТАТСССТСТАТСССТССАТАТТААТАА
HHV7_RK_U53.5	CGTGGTTCCCAAAAACGGTGCGCTCCAACAGATTCGGATGAAGAATGAGCTTTCCAGGA 339
B3_020413	CGTGGTTCCCAAAAACGGTGCGCTCCAACAGAATTCGGCAGTAGTA-AANTCACNTAGNG 418
HHV7_RK_U53.5 B3_020413	GATCCTGACTATACGACAAAGAAAGAAAGAAGCGGTATAGAGAAGATGACGATCGTGAACTC 399 TAANTTGCNNGGCNNNCCCGATCGGACAAGNTNNGCCTTNAGGATNCNTNANTACTTC 476 * ** * * * * * * * * * * * * * * *
HHV7_RK_U53.5 E3_020413	ACTAAAGACAAAAATGATATTAAAGAACTAGTTGATGCAATAGGGATGCTGCGTCATGAA 459 CNTGTGAGCGGGNTCNNGGGNGGGACC 503 * *
HHV7 RK 1754	ርሞርሚልፕናርርሞርጊል ርምል ምምየናዊጊል ሮዲያርጊል ይር - የርጋል ካል ጥንል ል አምታል አጥሪም አርሃጋል እር አል እ እ እ እ 1 ሳ ሳ ሳ

 HEY/KK_U54
 CTCCATGCGCTCAACTATTTGGACAGGAAG-TGATACCAAATTAATGCTAGGAAGAAAAA 1079

 B4_020313
 -----NGCGCTCAACNANN-GGACAGGAAGGNGATACCAAATNAATGCTAGGAAGAAAAA 54

ННV7_F В4_020	2K_U54 AGG0 0313 AGG0 ****	САЛАЛАДАТТАЛТАЛТТАЛДСGAЛАТТGTTTTАЛЛТСТАЛАТСТАСАТСАGATAT 11 САЛАЛАДАТТАЛТАЛТАЛАЛАЛИАЛИАЛТGTTTАЛЛТСТТАЛЛТСТАСАТСАДАТАТ 11 ***********************************	.37 .4
HHV7_F B4_020	RK_U54 ATT 313 ATT ****	TTGAACAGCCAAGGTTACATTGCCACAATCTCTTAGATCGATATACGTAAAGTAGGC 11 T-GAACAGCCAAGGNTACATTGCCACAATCTCTTAGATCGATATACGTAAAGTAGGC 17	.97 73
ННV7_Б В4_020	2K_U54 TGC 313 TGC ****	TGTAAAGCTTCCTTCATCTCTGCACATACACACAAATGTTGATTGTTGCGAA 12 TGNTNGGACCTTCCTTCATCTCTGCACATACACACAAATGTTGATTGTTGCGGGGGGA 23	252 33
HHV7_F B4_020	RK_U54 AGT/ 313 AGT/ ****	ARGCTTAGCCC-TGTCCTTAAGCATCTTATTTCACGAGGTTTTAAAAGGATAGGTGC 13 ANGCTTAGCCCCTGTCCTTAAGCATCTTATTTCACGAGGTTTTAAAAGGATAGGTGC 29	911 93
нну7_б В4_020	2K_U54 TGT0 313 TNN0 * 1	CAATTTTAGTTGTAGAAAATATGAAGACCAGACCACATAACAGGATTCCAT 1365 CAATTTTAGTGNTAGAAAATA-GTAAACCNNACCANANNACGGTATCTCTA 346 ********** ********* * **** **** *** *	
HHV7U5	52		
ННV7_F В5_020	RK_U55A TG 0413	TTCCTAAACCCTTTTCAAG-ATAGGCAAATTG-ATGAAATCTAAGAGGATCCATATTT 9 TACCCTTTTCAAGGATAGGCAAATTGGATGAAATCTAAGAGGATCCATATTT 9	958 52
ННV7_Б В5_020	RE_U55A TCT 0413 TCT +++	TTTAAAGTAACAATTATCTTGTATCAG-CAGGGATGACGTAATGTTAAAAATTGGCAG 1 TTTAAAGTAACAATTATCTTGTATCAGACAGGGATGACGTAATGTTAAAAATTGGCAG 1	1017 112
HHV7_F B5_020	RK_U55A GGG 9413 GGG ***	САЛЛАТАТТВАСАЛАСТТАТСВАТТСАТАССАЛАЛАСАТТАЛТТТВАТСАС 1 САЛЛАТАТТВАСАЛАСТТАТССАТТСАТАССАЛАЛАСАТТАЛТТТВАТСАС 1 ************************************	1077 172
ННV7_F В5_020	RK_U55A TT 0413 TT ***	TTTTGTTTTTTAGAACGTGACCCACAAAGTTACGCATTCTGCATTGTTTTCAGTTCC TTTTGTTTTTTAGAACGTGACCCACAAAGTTACGCATTCTGCATTGTTTTCAGTCC	1137 232
HHV7_F B5_020	RK_U55A ATC 0413 ATC ***	CGGTAATTAGCAACAGTACAGGTCTTTCGTGCGAAAGTATGCAGGTAATCTCTGCGGG CGGTAATTAGCAACAGTACAGGTCTTTCGTGCGAAAGTATGCAGGTAATCTCTGCGGG	1197 292
нну7_5 В5_02(K_U55A AGJ 0413 AGJ ***	алалаттда-дітстідстісалатсатталатссасадталастдаатталтідсі : Алалаттдалідітстідстісала	1256 317
HHV7U	558		
нн∨7_н В6_020	RK_U55B CG 0424	TCACTATGACCTTTTCGTTTTTGAATCTTATCAGACTAGGAAAAACTCCCATCTGCTC 	960 21
ННV7_1 В6_020	RK_U55B TT 0424 TT ***	GTTCTTTAACATCATGGTTTGGGATGATGACGATAGAGCTGTTAATTTTTAACAACGG GTTCTTTAACATCATGGTTTGGGATGATGACGATAGAGCTGTTAATTTTTAACAACGG	1020 81
нн∨7_1 В6_02(RK_U55B AA 0424 AA ++	atccaaaaaacaagacttttgatggataaatgtgtctattcagattttttaaatgtag atccaaaaaacaagacttttgatggataaatgtgtctattcagatttttttaaatgtag	1080 141
нн∨7_1 В6_020	RK_U55B TT 0424 TT **	TTAGTTGTTGTTCTTCCATTAAAGGACGAGCAGAAAAAAAGGAGTCAGAATTCCTTGA TTAGTTGTTGTTCTTCCATTAAAGGACGAGCAGAAAAAAAGGAGTCAGAATTCCTTGA	1140 201
нн∨7_1 В6_020	RK_U55B AT 0424 AT **	TTATTTCAGCGCTCAAAAGTAAAACTGGTTCAGTTTTGATTGGACATACTCGTCAAA TTATTTCAGCGCTCAAAAGTAAAACTGGTTCAGTTTTGATTGGACATACTCGTCAAA	1200 261
HHV7_1 B6_02	RK_U55B TA 0424 TA **	атаатытасаттатстттассствеааттсттсттутатвеаатттсаатстваас атаатытасаттатстттассствеааттсттсттва	1260 302
HHV70	56		
HHV7_1 B7_02	RK_U56 GTC 0419	ТСТАВОСАТСАСТАТСЯССКСА. ТСТАВОСАТСАСТАТСЯСТСС-САСТТАССТАВСТААСТААТТОТТ 5 	39 2
HHV7_ B7_02	RK_U56 GCG 0419 GCG ***	SAAGGCTTCAGAAATGGTGGCAGTAAACTCAATTGATCTCCCTTTTCCAGAGGAAATG 5 SAAGGTTTCCAGAAATGGTGGCAGTAAACTCAATTGATCTCCCCTTTTCCAGAGGAAATG 1	99 02
ННV7_ В7_02	RK_U56 GTC 0419 GTC ***	CCGGTATTTTTGATTTCATACTGGTGTCCAATTTTTGGACGTATTAATAATTAAT	59 62
HHV7_ B7_02	RK_U56 TTC 0419 TTC ***	ССТТСТАСТТТТТТБТААGAГТОЛТААСАССАТТЭТТТТТААGAТАТСССТТАТТТGАА 7 ССТТСТАСТТТТТБТААGAГТGTTAACACCATTGTTTTTAAGATATCCCTTATTTGAA 2	19 22
HHV7_ E7_02	RK_U56 TAT 0419 TAT ***	TAATCGGTGGTTGTAGATAAGACCGTGTAAAGTCCTAAATTTGTGTTGCCAATTAA 7 TAATCGGTGGTTGTAGATAAGACCGTGTAAAGTCCTAAATTTGTGTTGCCCCAATTAA 2	77 82
ннv7_ В7_02	RK_U56 ATG 0419 ATG ***	BACGGTGCGACGG-TATT-GGGATGACGGCATTAGTGAGTTTGCATAAAGTTCCAATG 8 BACGGTGCGACNGGTATTTGGGATGACGGCATTAGTGAGTA	35 25

HHV7U57		
HHV7_RK_U57 B8_020425	TGTGTTTTTTAAAGGTGTTTTCTAAATGCGCTAAACAAGCAGCACTTAATTCAAACGATAT GAGGGGGENNCGCGCCAANGACGNNANNGATCAGATAG * * * * * * * * * * * * * * * * * * *	3660 37
HHV7_RK_U57	ATTGATAGGATGCTTTTCTGAATACTTGGTAACCAGTACGGTTGTTTCTTTAGGCGCAGT	3720
B8_020425	GGGAGCNOCNCGNATNGAGGTGCNNCNCGCANNNNGCGGGGGNNNCTATT-GGACANNAC	96
HHV7_RK_U57 B8_020425	TACGTCGTTTCCTGTAGCGACTCTAGGTAATTGTATATAAAACAGAATTTTTCCCAGTGA NNTNNTCCTNNA-CGACTC-AGGTATGGATANNNAA-ANNANTTCCNNNACAAG **** * ****** ***** ** ** ** ** ** **	3780 147
HHV7_RK_U57 B8_020425	CATTTTGTCTAGGTCGTTGAAACGGATAACATTTGCAGCAACCGCTATGGACGTATGAAA CGTNGAGTGACAAACGGATACATTGACNNCCGTAGACGTANNNAA * * ** * * * * * * * * * * * * * * *	3840 192
HHV7_RK_U57 B8_020425	ARARTCTATCCATTCAGTCCGATTACAGTARATCCCAAGTAATGCTTCGARACTTATGTT NATATATCCATTCAGTCCGATTACAGTNA-TCCCAAGTAATGNTTNNGNNCNTANGN- * * * * * *	3900 248
HHV7_RK_U57	ATAACGA-TCAGAGTCGTCACCGTAATACAATCTTAAGTTTTCAAAAAGTTGTTCAGCAG	3959
B8_020425	ATAACGGGTCAGAGTCGTCACCGTA-TACATCTAGTTCATATGTACNTTGTTANTAT	304
HEV7_RK_U57	TTTGTGTTCTGATATCATCATAACACGTTTGGAGAAACATCTAGTTTTGGGAAAATTTCAG	4019
B8_020425	AAACRAATTNTTTANTT	321
HNV7U58		
HHV7_RK_U58	АТGАТАGАСТСААТАТСТСЯААGАААСТТТААТАЭТАААGAGCTACACCGTTAATCACTGT	60
B9_020308		4
HHV7_RK_U58 B9_020308	GCTAAAAATGTTCCAGTGTTTATTAACTCCTATGATTTAACCGCAGAAGTGGCCCAAAAAT GCNGNATGTNNTGGTACTACCCNATGATNACCGCGANTGGCCAANT ** **** ** *************************	120 50
HHV7_RK_U58 B9_020308	GAAGACGTGCGATTAGCGCGACAAGTTCAAATTTCATTAGAGAAAATAGATGAAG -GAGACGTGCGATTAGCGCGACGAGTINCGNNTTTCTTAGACCCCNCNNNATAGATGAAG ******	175 109
HHV7_RK_U58 B9_020308	TTATAGAATCAATTTTTTCTGCGTCTGGTCCTAGCGTTGAAAATGTAAAAGATCAGGCAA TTATAGAATCCCTTTTTTCTGCGTNTGGTCCTAGCGTTGANTGTANATCCNGGCN- ****	235 164
HHV7_RK_U58	AGTTIGCTITIGTGGGTTTACTGGTCCTGTGA-GTATTCCGTGCTAC	285
B9_020308	-GTTIGCTTTGTGTCGTTTACTGGTCCGTGGANNATNATNCAACAGATAGCCTAC	223
HHV7_RK_U58	TGCGAAGAATGGGATGTCAATTTTTATCTGACAAAATGTAGTTATAATTGCGAAGGCCCG	345
B9_020308	TGCGAAGANTGGGATGT-ATTTTTTCTCTGACC	255
1HV7059		
HHV7_RK_U59 B10_020502	ATGACAGCAAAACAAATTTTTCCAGTTTGGAAAGACTGTACACTTTACGTTAACAATGAG TAACCNNGGANGNTATGCAANGGGAG ** * ** ** *	60 26
HHV7_RK_U59	ACAGCAACTGTGCATG-AAATCCTGAATTCCGAATTTAAGTGAACTGTTACAGTTAAAGAC	119
B10_020502	NNGACTACTNTGCATGGAAATCCTGGAATTCCGAATTTAAGTGAACTGTTACAGTTAAAGAC	86
HHV7_RK_U59	GGAATTTGTATCTATGACAGACCTATGTGTTTTACATTACTGGATGTATAAATCAGAATAT	179
B10_020502	GGAATTTGTATCTATGACAGACCTATGTGTTTACATTACTGGATGTATAAATCAGAATAT	146
HHV7_RK_U59	TTCCAGCATCACGATATATTGGCATGCTTACAGTGAAGTAATTTATGCTTTAACTGGAAT	239
B10_020502	TTCCAGCATCACGATATATTGGCATGCTTACAGTGAAGTAATTTATGCTTTAACTGGAAT	206
HHV7_RK_U59	TATACACTGTGAAAAGATTTCTATTGAATGTGGAATTAAATCCACGGACAATAACATTTT	299
B10_020502	TATACACTGTGAAAAGATTTCTATTGAATGTGGAATTAAATCCACGGACAATAACATTTT	266
HHV7_RK_U59	GTATGAAAAGCCCAAACTGTTTTTACTTCGAGAAAATTTAGCACCAACTGAATTAAGGTG	359
B10_020502	GTATGAAAAGCCCAAACTGTTTTTA	291
HHV7U62		
HHV7_RK_U62 B12_020425	атсалсосасттталассататаласастсястттатсталасасасстт	60
HHV7_RK_U62	ТТААССІТІСАТАСААААААТТТОТСТСААСТСІЛАТТСАТСТТСААСССІСІЛАДААТСТ	120
B12_020425		45
HHV7_RK_U62	ТТТССТАААААААССGАGTTGGTTGGGTGGATGTATGATACGCTCGCGGTGGAAATTTTT	180
B12_020425	ТТТССТАААААААССGAGTTGGTTGCGGTGATGTATGATACGCTCGCGGTGGAAATAT	103

HHV7063		
HHV7_RK_U63 F1_020228	ATGAACAAAAAAAAGATGGACTTGCCTAAATGTCAGTCAA-TCACAG-TTGCTTGTGAAG 	58 32
HHV7_RK_U63 F1_020228	GAGAATGTTCGCAAATGTACAATTTGCATAATCCTCTGACATTTGAAATGGGTTTGGGAA GAGAATGTTCGCAAATGTACAATTTGCATAATCCTCTGACATTTGAAATGGGTTTGGGAA	118 92
HHV7_RK_U63 F1_020228	ACATCTTTATATGTGTTCGGTGTTTTAAGATACATTTTTGCAATATGCTAGAAGACTGCA ACATCTTTATATGTGTTCGGTGTTTTAAGATACATTTTTGCAATATGCTAGAAGACTGCA	178 152
HHV7_RK_U63 F1_020228	ACCTGATAAATA-CTCATGAAGGATGCGTGTGTGTCAAAAACCGGGCTTTTTTATAACGGA ACCTGATAAATAGCTCATGAAGGATGCGTGTGTTCAAAAACCGGGCTTTTTTATAACGGA	237 212
HHV7_RK_U63 F1_020228	TGGATGCCAGCCTATTCACATACCTGTATGGAACCTACTGAAGAGCCAAATATGGAGACC TGGATGCCAGCCTATTCACATNNCNGTATNNNNCCTNCNAGANNCNAN-ATGTNNACC **********************************	297 269
HHV7_RK_U63 ₽1_020228	GTTAATGTAGTTGTAGTGCTGTTATCATACGTTTACAGTTTTTTAATACAAAAAAAA	357 303
HHV7U64		
HHV7_RK_U64 F2_020425	GGATCTTTTTCAGTGTGTATTTTTTAAAGAAAAGCTTGTATATACGGAATGGACAAAA 	180 26
HHV7_RK_U64 F2_020425	ATAAAGTITACCTTATGTGTTACATGATCTTGT-AATTTCTCAAATCTTTTAAGAATGCCTG ATAAAGTITACCTTATGTGTTACATGATCTTGNTAATTTCTCAAATCTTTAAGAATGCCTG	239 86
HHV7_RK_U64 F2_020425	TATTAAAGAAGTAATACATGGGGCATTAATTCTTTCAGTTCCCATAAATAA	299 146
HHV7_RK_U64 F2_020425	асаттттдатасадататтттааттсталалаттатттасссасаттттттдсасдатда асаттттдатасадататтттааттсталалаттаттасссасаттттттдсасдатда ******	359 206
HHV7_RK_U64 F2_020425	TATTGTCATAAAATTATCGGAAATTTTGTCTGGAGCACCTCGCATACAAAAAACAGTGGA TATTGTCATAAAATTATCGGAAATTTTGTCTGGAGCACCTCGCATACA	419 254
HHV7065		
HHV7_RK_U65 F3_020313	TCTCCCTGTTTTCGCTGTTGATAAATTTAAGATTGCTTGC	180 54
HHV7_RK_U65 F3_020313	AAAAAGAATTTOGATTTTOTT-TTTAGCTTGGCCATAAACGGAATACATGCCGGACAGTT AAAAGAANTTTGGATTTTGTCCTTTAGCTNGGCCATNAACGGAATAC-TGCCGGACAGTT **** * *****************************	239 113
HHV7_RK_U65 F3_020313	ТЭСААСАААТТССАТТАААСТААААААААТТАТААТААСАААСЭАТТТЭЭТЭСТАСАТ ТЭСААСАААТТССАТТАААСТАЭААААААТТАТААТААСАА-СЭАТТТЭЭТЭЭТЭСТАСАТ *********************************	299 172
HHV7_RK_U65 F3_020313	ATTAGAATTGGGATCTTTAACGGTAA-CGGATTTACATTCCATACATAGAAT-ACAATAG ATTAGAATTGGGATCTTTAACGGTAAACGGATTTACTTCCNNNTNNTTNCNNTAG ************************************	356 232
HHV7_RK_U65 F3_020313	ТЗААСАТСТССКААТСТСССАССТАТТАССССАААТСТААТТТАТЗАТАСТТСТСААТ ТСЛААСТСТССССААТСТССССАТТАСС **** *******************************	416 263
HHV7066		
HHV7_RK_U66 B11_020314	TCGATCTGTAAAAACATGGGATTTCTGTAAAAACTTGTTGTAGCTCTAGATGTTTTTGA ATNGAAAAACAGGGATTCGANNANGNGAGNNNTAGANGTTTTTGA * *** * ** *** * *** *** *** *** ******	4800 45
HHV7_RK_U66 B11_020314	TATTG-TACTTCTGTATTTTTCTC-TATGATTGCAGTAAATTTGAGCTTTGACAAAGC TATTGGTACTTCTGNGNNTTCNNNANNNATTGCAGTAAATTTGAGCTTNGACAAATGCT ***** ******** ** * *****************	4856 105
HHV7_RK_U66 B11_020314	GCAGT-TCAAGGGGTCGCATATTG-CAATTTTAGATTTTACGTGCCGTTGGCGATC- GCAGTGNCAATGGGGTGCTGCATATTGGCAATTTTAGATTTTACGTGCCGTTGGCGATNC	4910 165
HHV7_RK_U66 B11_020314	ACAAAAGAGATATAAAGGTTTAACAT-GCCTGCAATAT-GCATGTGTAAAGCCAAGTTC- ACAAAAGAGATATAAAGGTTTAACATTGCCTGCAATATTGCATGTGTAAAGCCAAGTTTA ************	4967 225
HHV7_RK_U66 B11_020314	-ТЭСАСТТААТАТААТААТССЯ-ТТТТЭССААААТАТСССССТСЯТТЭСААААСТ ТЭССАСТТА:	5021 284
HHV7_RK_U66 B11_020314	TGAAGAAATCTTTACGTCCTGTTCATGTTTCCAAATTATAGACTGATACGCTTTCTGAAT TGA-GAATNCTTTACN-CCTGT-CATNTTCCAATTATTTTTTTTNNNAATTTNNNTT *** *** ***** ***** ***** * ***** * * ****	5081 339
HHV7_RK_U66 B11_020314	TGCATCTATATCGCATGATCGCAACAT 5108 TNTGCATT 347	
HHV7U67		
--	--	--
HHV7_RK_U67 F4_020313	САТССАСАСАСАЛАТТТСТСТАССТССААТТТАТАААССААСС	60 39
HHV7_RK_U67 F4_020313	GGATGCTAAAATTTTCTCGGAGGCAGTGCAGACCGCACTAACTGTGTGTAAAGCAAC GGATGCTAAAANTNCTCGGANNAAAANGNNGCAGACCGCACTAACTGTGNGTAAAGCAAC	117 99
HHV7_RK_U67 F4_020313	С-GCTССТААТАСАСЭТСТААААСТСЭ-ТТЭАААСАССААСТААТААСТТСТТАСТАЭТА СТЭСТССТААТАСАСЭТСТААААСТСЮЭТИЭЛААСАССААСТААТААСТТСТТАСТАЭТА 	175 158
HHV7_RK_U67 F4_020313	асалатсттоттссатсадаласттсдала-дсалсдастдалд-салатстталтатт асалатстд-тссатсадаласттсдалатдсалсдастдализсалатстлаттики ******	233 216
HHV7_RK_U67 F4_020313	ATGCAGCGTTGGAAAAACTGGCGTCTTCCTTTAATACAGCGGTACCTGTAAAA GNNNATGGATGGGCCNGGAAAACTTN-GNGNTTCCTTAATACAGCGTNACCTTGAAT	287 273
HHV7_RK_U67 F4_020313	CATCCAAAAAGTATTTGTTGCAAAATGTGAGAAAAATGACCAGTGAGAACATCGCTCTAA CATCCAAATGATNT	347 287
HHV7U68		
HHV7_RK_U68 F5_020425	атсятсястосатсялаталагаластатсятссялалатттасялалалалсяттат - тотсястосатсялаталагаласаластатсятссялалатттасялалалаласяттат **********************************	60 59
HHV7_RK_U68 F5_020425	GAGITIGITAAAATTAAAACITIGGIGAAGATCATCCTCTTAGCGITCGACAGCAAATTCAC GAGITIGITAAAATTAAAACITIGGIGAAGATCATCCTCTTAGCGITCGACAGCAAATTCAC	120 119
HHV7_RK_U68 F5_020425	GCTCTCAATCAAAATCTTGTATCAGAAAATCTCGAACAGTCCCAGATAATTACTTCTTTG GCTCTCAATCAAAATCTTGTATCAGAAAATCTCGAACAGTCCCCAGATAATTACTTCTTTG	180 179
HHV7_RK_U68 F5_020425	acaaaaatgttaaaggatcaaaagctgcaggtgaaagggaaaaatgctggtgg acaaaaatgttaaaggatcaaaggtgcaggtggaaggggaaaaatgctggtgg	240 239
HHV7_RK_U68 F5_020425	CTAGAATGTGTAGATTTGGATGACATTTTTGGATACGGCAGCGGAAGTGAAATCCGTCACC CTAGAATGTGTAGATTTGGATGACATTTTGGATACGGCAGCGGAAGTGAAATCCGTCACC	300 299
HHV7_RK_U68	GACAATATAAAAGAAACTTTACTGGCCGGATTAGAATCAGACTAA 345	
	unca	
NHV7U69	**** ****	
HHV7U69 HHV7_RK_U69 F6_020425	атодадсадсттаадасассоссалаатсаалаласасотосладаалтатдсттоста тосадсттаадасасоссалаатсаалаласасотосладаалтатдсттоста	58 56
HHV7U69 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425	ЗОЗ **** АТОЗАЯСАЯСТТААВАСАССССААЛАТСААЛААСАСЭТССААВАЛТАТGCTTCCTA ТGCAGCTTAAGACACCCCCАЛАЛТСААЛАААСАСЭТССААВАЛТААТАGCTTCCTA ***** АЛАЛАЛАЯВДАЛААСАСССССААЛАТСААЛАААССЭТССААВАЛТААТА	58 56 118 116
HHV7U69 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425	атодадсадсттаадасассссаллатсааллаасасотссалдалататдсттеста тдсадсттаадасассссаллатсааллаасасотссалдалатаа. аллалаладдаладаасттаалааладоссттоталаадтааласотааттатттоот аллалаладдаладаасттаалааладоссттоталадоталаасоталаттатттоот аллалаладдаладаасттаалааладоссттотогаладоталаасотааттатттоот ссодаласатсадасстаасалалаалатасстстодосттсадасотодатаасолаттод ссодаласатсадасстаасалалаалатасстстодосттсадасотодатаасолаттод	58 56 118 116 178 176
HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425	атодадсадсттаадасассссаалатсаалаласасотссаадалататдсттсста тдсадсттаадасассссаалатсаалаласасотссаадалаталдонсттсста тдсадсттаадасасссссаалатсалаласасотссаадалаталдонсттсста Алалаладдааладалсттаалаладоссттоталалоталалсоталаттатттодот алалалаладдаласаталалаладоссттототалалоталалсоталаттатттодот сседаласатсадасстаасалалаладосстстедеттсадасотодаталасдааттед сседаласатсадасстаасалалалаласстстедесттсадасотодаталасдааттед алалалаласатсадосстаасалалалаласстсетодосттсадасотодаталасдааттед сседаласатсадосстаасалалалаласстсетодосттсадасотодаталасдааттед алалалаласододостседатедатаседалаласодасствадасодассттате	58 56 118 116 178 176 238 236
HHV7U69 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425	ATGGAGCAGCTTAAGACACCCCCAAAATCAAAAAACACGTCCAAGAAATATGCTTCCTA TGCAGCTTAAGACACCCCCAAAATCAAAAAACACGTCCAAGAAATAAAGCTTCCTA TGCAGCTTAAGACACCCCCAAAATCAAAAAACACGTCCAAGAAATAAAGGCTTCCTA AAAAAAAGGAAAAGAACTTAAAAAAAGGCCTTGTAAAGTAAAACGTAAATTATTTGGT AAAAAAAAGGAAAAGAACTTAAAAAAAATACCTCTGGCTTCAGACGTAGATAACGAAATTGG CCGAAAACATCAGGACCTAACAAAAAAATACCTCTGGCTTCAGACGTGGATAACGAAATGG CCGAAAACATCAGGCCTAACAAAAAAATACCTCTGGCTTCAGGACGTGGATAACGAAATGG CCGAAAACATCAGGCCTGAACAAAAAAATACCTCTGGCTTCAGGACGTGGATAACGAAATGG AAAAAAAACGGGGCCTGGATGATACGAAAACGGTCTGAGACGGGACTTATGTCCAGATCCAT AAAAAAAACGGGGCTGGATGATACGAAAACGGTCTGAGACGGGACTTATGTCCAGATCCAT	58 56 118 116 178 176 238 236 297 296
HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425	ATGGAGCAGCTTAAGACACCCCAAAATCAAAAAACACGTCCAAGAAATATGCTTCCTA TGCAGCTTAAGACACCCCCAAAATCAAAAAACACGTCCAAGAAATATAGCTTCCTA AAAAAAAGGAAAAGAACTTAAAAAAAGGCCTTGTAAAGTAAAACGTAAATTATTTGGT AAAAAAAAGGAAAGAACTTAAAAAAAAGGCCTTGTAAAGTAAAACGTAAATTATTTTGGT CCGAAAACATCAGACCTAACAAAAAAATACCTTCTGGCTTCAGACGTGGATAACGAAATGG CCGAAAACATCAGGCCTGATGATACGAAAAAATACCTTGGGTCTGAGACGGAGCTTATGTCCAGAACGAATTGG CCGAAAACATCAGGCCTGATGATACGAAAACGGTCTGAGACGGACTTATGTCCAGAACGAATTGG CCGAAAACATCAGGCCTGATGATACGAAAACGGTCTGAGACGGACTTATGTCCAGAACCGATCCAT AAAAAAAACGGGGGCTGGATGATACGAAAACGGTCTGAGACGGACTTATGTCCAGATCCAT CTGTAACAGGCCTCCTATGTCATGGAATCTTTGACTGTATCTCCAAAGTTTGAACGAAAAC GGATTGAGTGCATGCACGGAATTTGAGAATTTTATGGATACCAAGGAAAATCGTGTTAAAGT	58 56 118 116 178 238 236 297 296 357 311
HHV7U69 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425	ATGGAGCAGCTTAAGACACCCCCAAMATCAAAAAACACGTCCAAGAAATATGCTTCCTA TGCAGCTTAAGACACCCCCAAMATCAAAAAACACGTCCAAGAAATAAAGCCTTCCTA TGCAGCTTAAGACACCCCCAAMATCAAAAAACACGTCCAAGAAATAAAGGCTTCCTA AAAAAAAAGGAAAAGAACTTAAAAAAAGGCCTTGTAAAGTAAAACGTAAATTATTTTGGT AAAAAAAAGGAAAAGAACTTAAAAAAAAGGCCTTGTAAAGTAAAACGTAAATTATTTTGGT CCGAAAACATCAGGACCTAACAAAAAAATACCTCTGGCTTCAGACGTAGATAAACGAAATTGG CCGAAAACATCAGGCCTAACAAAAAAATACCTCTGGCTTCAGACGTAGATAACGAAATTGG CCGAAAACATCAGGCCTGATGATACGAAAAAAATACCTCTGGGCTTCAGACGTGGATAACGAAATTGG CCGAAAAAAAACGGGGCTGGATGATACGAAAAAAATACCTCTGGGCTTCAGACGTGGATAACGAAATGG CCGAAAACATCAGGGGCTGGATGATACGAAAACGGTCTGAGACGGGACTTATGTCCCAGATCCAT AAAAAAAACGGGGCTCGGATGATACGAAAACGGTCTGAGACGGGACTTATGTCCAGATCCAT AAAAAAAACGGGGCTCGTATGTCATGAAACGGTCTTGACGGGACGTAGCTTGAACGAGAC-T CTGTAACAGACCTCCTATGTCATGAATCTTTGACTGTATCTCCCAAAGTTTGAACGAGAA-T CTGTAACAGACCTCCTCTATGTCATGAATCTTTGACTGTATCTCCCAAAGTTTGAACGAAAAT GGATTGAGTGCATGCACGGAATTTGAGAATTTTATGGATACAAAGAAAACGGTTAAGT	58 56 118 176 238 236 297 296 357 311
HHV7U69 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425	ATGGAGCAGCTTRAGACACCCCCAAAATCAAAAAACACGTCCAAGAAATATGCTTCCTA TGCAGCTTRAGACACCCCCAAAATCAAAAAACACGTCCAAGAAATAAAGCATCCCTA TGCAGCTTRAGACACCCCCAAAATCAAAAAACACGTCCAAGAAATAAAGCATTCCTCA AAAAAAAGGAAAAGAACTTAAAAAAAGGCCTTGTAAAGTAAAACGTAAATTATTTTGGT AAAAAAAAGGAAAAGAACTTAAAAAAAAGGCCTTGTAAAAGTAAAACGTAAATTATTTTGGT AAAAAAAAGGAAAAGAACTTAAAAAAAATACCTTCTGGCTTCAGACGTAGATAACGAATTGG CCGAAAACATCAGGACCTAACAAAAAAATACCTTCGGCTTCAGACGTAGATAACGAATTGG CCGAAAACATCAGGCCTAACAAAAAAATACCTTCGGCTTCAGACGTAGATAACGAATTGG CCGAAAACATCAGGCCTGAACAAAAAAATACCTTCGGCTTCAGACGTAGATAACGAATTGG CCGAAAACATCAGGCCTGAACAAAAAAATACCTTCGGCTTCGAGACGTAGATAACGAATTGG CCGAAAACATCAGGCCTCGATGATACGAAAACGGTCTGAGACGGACTTATGTCCAGAATCGA AAAAAAAACGGGGCCTGGATGATACGAAAACGGTCTGAGACGGACTTATGTCCCAGATCCAT AAAAAAAACGGGGCCTGATGATACGAAAACCGTCTGAGACGGACTTATGTCCCAGATCCAT CTGTAACAGACCTCCTATGTCATGAATCTTTGACTGTATCTCCCAAAGTTTGAACGAAAAT	58 56 118 116 178 176 238 236 297 296 357 311
HHV7U69 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U70 F6_020425	ATGGAGCAGCTTAAGACACCCCAAAATCAAAAAACACGTCCAAGAAATATGCTTCCTA TGCAGCTTAAGACACCCCCAAAATCAAAAAACACGTCCAAGAAATAAAAGCTTCCTA AAAAAAAGGAAAAGAACTTAAAAAAAGGCCTTGTAAAGTAAAACGTAAATTATTATGGT AAAAAAAAGGAAAAGAACTTAAAAAAAAGGCCTTGTAAAGTAAAACGTAAATTATTATGGT CCGAAAACATCAGACCTAACAAAAAAATACCTTGGGTTCAGACGTGGATAACGAAATTGG CCGAAAACATCAGACCTAACAAAAAAATACCTTGGGTTCAGACGTGGATAACGAAATTGG CCGAAAACATCAGGCCTGATGATACGAAAACGTCTGAGACGGGCTTATGTCCAGATCAGATTGG CCGAAAACATCAGGCCTGGATGATACGAAAACGTCTGAGACGGACTTATGTCCAGATCCAGATTGG CCGAAAACATCAGGCCTGGATGATACGAAAACGGTCTGAGACGGACTTATGTCCAGATCCAT AAAAAAACGGGGGCTGGATGATACGAAAACGGTCTGAGACGGACTTATGTCCAGATCCAT CTGTAACAGACCTCCTATGTCATGGAATCTTTGACTGTATCTCCCAAAGTTTGAACGAAGA- TCTGTAACAGACCTCCTATGTCATGGAATCTTTGACTGTATCTCCCAAAGTTTGAACGAAGA- CTGTAACAGACCTCCTCTATGTCATGGAATCTTTGACTGTATCTCCCAAAGTTTGAACGAAGAA- CTGTAACAGATGCATGCACGGAATTTGAGAATTTTATGGATACAAGGAAAAATCGTGTTAAAGT GGATTGAGTGCATGCACGCACAAATTTCTTGTAATTTGGCTTCTATTATAGAAGAAGGACC	58 56 118 176 238 236 297 296 357 311 57 60
HHV7U69 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U70 F7_020425 HHV7_RK_U70 F7_020425	ATGGAAGAGCTTAAGACACCCCAAAATCAAAAAACACGTCCAAGAAATATGCTTCCTA TGCAGCTTAAGACACCCCCAAAATCAAAAAACACGTCCAAGAAATAAAGNCTTCCTA TGCAGCTTAAGACACCCCCAAAATCAAAAAACCGTCCAAGAAATAAAGNCTTCCTA AAAAAAAGGAAAAGAACTTAAAAAAAGGCCTTGTAAAGTAAAACGTAAATTATTTGGT AAAAAAAAGGAAAAGAACTTAAAAAAAATACCTCTGGCTTCAGACGTAGATAATTATTTGGT CCGAAAACATCAGGACCTAACAAAAAAATACCTCTGGCTTCAGACGTGGATAACGAATTGG CCGAAAACATCAGGCCTGAACAAAAAAATACCTCTGGCTTCAGACGTGGATAACGAATTGG CCGAAAACATCAGGCCTGAACAAAAAAATACCTCTGGCTTCAGACGGGACTTATGTCCAGAACGAATTGG CCGAAAACGGGGCTCGATGATACGAAAACGGTCTGAGACGGACTTATGTCCAGAACGAATTGG CCGAAAACGGGGCTCGATGATACGAAAACGGTCTGAGACGGGACTTATGTCCAGATCCAT AAAAAAAAACGGGGCTCGATGATACGAAAACGGTCTGAGACGGGACTTATGTCCAGATCCAT CTGTAACAGACCTCCTATGTCATGAATCTTTGACTGTATCTCCCAAAGTTTGAACGAAAACG GGATTGAGTGCATGCACGGACATGTCAGGAATTTGAGTATATGGGATACGAAGAAACGGAC CTGGTACAGGACTGCCACGGAATTTCGTGAATTTGGGATACCAAGGAAAATCGTGTTAAGT GGATTGAGATAGGATTACGCACGAAATTTCTTGTAATTTGGCTTCTATTATAGAAGAGGGAC TTGGGCAATAGATTACGCACAAAATTTCTTGTAATTTGGGACATTTCAAGAGAGAAAATT CCGGTCTTTTTTATTCTAATAGACAAATTAAACAATCTGGACATTTCAAGAAGAGGACA	58 56 118 176 238 236 297 296 357 311 57 60 117 120
HHV7U69 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U69 F6_020425 HHV7_RK_U70 HHV7_RK_U70 F7_020425 HHV7_RK_U70 F7_020425	ATGGAGCAGCTTAAGACACCCCAAAATCAAAAAACACGTCCAAGAAATATGCTTCCTA TGCAGCTTAAGACACCCCCAAAATCAAAAAACACGTCCAAGAAATATAGCTTCCTA AAAAAAAGGAAAAGAACTTAAAAAAAGGCCTTGTAAAGTAAAACGTAAATTATTTGGT AAAAAAAAGGAAAAGAACTTAAAAAAAGGCCTTGTAAAGTAAAACGTAAATTATTTGGT AAAAAAAAGGAAAAGAACTTAAAAAAAAGGCCTTGTAAAGTAAAACGTAAATTATTTGGT AAAAAAAAGGAAAGAACTAAAAAAAAGCCTTGGAAGAGACGTGGAATAACGAATTGG CCGAAAACATCAGGCCTGAACAAAAAAATACCTCTGGCTTCAGACGTGGATAACGAAATTGG CCGAAAACATCAGGCCTGAAGAAAAAAATACCTCTGGCTTGAGACGGACTTATGTCCAGATCCAG AAAAAAAACGGGGGCTGGATGATACGAAAACGGTCTGAGACGGACTTATGTCCAGATCCAT AAAAAAAACGGGGGCTGGATGATACGAAAACGGTCTGAGACGGACTTATGTCCAGATCCAT AAAAAAAACGGGGGCTGGATGATACGAAAACGGTCTGAGACGGACTTATGTCCAGATCCAT CTGTAACAGACCTCCTATGTCATGGAATCTTTGGCTGTATCTCCCAAAGTTTGAACGAAGA-T CTGTAACAGACCTCCTCATGTCATGGAATCTTTGACTGTATCTCCCAAAGTTTGAACGAAGA-T CTGTAACAGACCTCCTCTATGTCATGGAATCTTTGGCTTCTATTATAGAAGGAGGAC TCGGTCTTTTTATTCCTAATAGACAAATTAAACAATCTGGAAAATCGTGTTAAAAATCGTGTTTAAAGACGAAAATT CGGTCTTTTTATTTCTAATAGACAAATTAAACAATCTGGACATTCAAGAAGGACAAATT CCGGTCTTTTTATTTCTAATAGACAAATTAAACAATCTGGACATTTCAAGAAGGAAAATT	58 56 118 176 238 236 297 296 357 311 57 60 117 120 177 180

HHV7_RK_U70 TTTATCGATGAGAATGTTGAGCTATCTGATTTATATAGCAATATTTATGT-CCGCTTACA 296 F7_020425 TTTATCGATGAGAATGTTGAGCTATCTGATTTATATAGCAATATTTATGTACCGCTTACC 300

HHV7_RK_U70 F7_020425	АGАТGCGAGTCCAAAAGTTGTTAAGAATCTATTTAAAATATTAGAACGAGAGACAAGAGG AGATGCGA	356 308
HHV7071		
HHV7_RK_U71 F8_020425	-ATGGGCTCAAAGTGCTGTAAGACAATACATGGCGGCATCTTCAGCAAGGCCGAAGACAC TATGGGCTCAAAGTGCTGTAAGACAATACATGGCGGCATCTTCAGCAAGGCCGAAGACAC	59 60
HHV7_RK_U71 F8_020425	CTTAGTAGACTATAAAGGAAAAATATATATAATCTTGAAAAAGAATTTTCTGCTTTAAGTGA CTTAGTAGACTATAAAGGAAAAATATTAATCTTGAAAAAGGAATTTTCTGCTTTAAGTGA	119 120
HHV7_RK_U71 F8_020425	ТАСТGААТСТGААGААGАЭТТGСААСТАGАGААGССАСТТСТАААТАААСААGАТТСТАG ТАСТGААТСТGААGААGАЭТТGСААСТАGАGААGССАСТТСТАААТАААСААGАТТСТАG	179 180
HHV7_RK_U71 F8_020425	ССЯТТССЯТЛАСССАДААСАЛАСТТСАЛАЛАТСАЛТССАЛАТАЛ 222 ССЯТТССЯТЛАСССАДААСАЛА 202	
HHV7U72		
HHV7_RK_U72 F9_020502	ACACGATAAACTATCCCCCATCCAAGTTATATCCCTTGTAGATTGATGTTGGGAGT TTCCCCCATCCAAGTTATATCCCTTGTAGATTGATTGATGTTGGGAGT	720 48
HHV7_RK_U72 F9_020502	ТТТТТССБАЛСАБААТАСТТТТАСССАЛСАЛАСТАТАТАБТАТАСТАЛДАТБАЛАЛА ТТТТТССБАЛСАБАЛАСТТТТАСССАЛСАЛАСТАТАТАБТАТАСТАЛДАТБАЛАЛА *******	780 108
HHV7_RK_U72 F9_020502	GCAGATATCCATGAAAACAACATAGCAGATAAGTTGGATAGGATTTAGGTAGAGTTGTGG GCAGATATCCATGAAAACAACATAGCAGATAAGTTGGATAGGATTTAGGTAGAAGTTGTGG 	840 168
HHV7_RK_U72 F9_020502	COTTAGGCTGCGGATGTCATTCCGTATAGAGAGGTTGATGGCTTTCATATCATTGATTTC CGTTAGGCTGCGGATGTCATTCCGTATAGAGAGGTTGATGGCTTTCATATCATTGATTTC	900 228
HHV7_RK_U72 F9_020502	ATAATATGCGCATGGGAATCCCAAATTGGGAAAGTGAACAGCCACTAAATGCACTGTGAC ATAATATGCGCATGGGAATCCCAAATTGGGAAAGTGAACAGCCACTAAATGCACTGTGAC	960 288
HHV7_RK_U72 F9_020502	СТТСЯСАТАССЯТТАЛАСЛАGСАСАGАТТАТАСТТАЛGАСССАЛАТТСТСАТАТТАЛТСАС СТТСЯСАТАССЯТТАЛАСЛАССАСАGАТТАТАСТТАЛGАСССАЛАТТСТСАТАТТАЛТСАС ****	1020 348
HHV7_RK_U72 F9_020502	ATCTACTCTGCTCAATGCCAT 1041 ATCTACTCTGCTCAATGCCAA 369	
HHV7073		
HHV7_RK_U73 F10_020502	ATGGAAACTCAGCTTCAAAATGATCAATTATTTTTGGAATGGTTTGGACAAAATTTGTTG 	60 31
HHV7_RK_U73 F10_020502	GATTGCCACTTTGCACAAAACGTTTCAGTGTATTTGCAAGATGCTTCAATGGTTCATTTT GATTGCCACTTTGCACAAAACGTTTCAGTGTATTTGCAAGATGCTTCAATGGTTCATTTT	120 91
HHV7_RK_U73 F10_020502	АЛААССТТТТСТСКААСАААТААЛААТТАТААСААССССААТССССТСТСКССААЛААССТСТ АЛААССТТТТСТСКААСАААТААЛААТТАТААСААССССААТССССТСТСКССААЛААССТСТ ****	180 151
HHV7_RK_U73 F10_020502	GCATTGATAGAATTCTTGAAAACTGTTTCATATATTGATTCTGTCCTTGTTATTTCCTGT GCATTGATAGAATTCTTGAAAACTGTTTCATATATTGATTCTGTCCTTGTTATTTCCTGT	240 211
HHV7_RK_U73 F10_020502	СЭТААААСТТТТЭСТЭСАGААСТТТТАААТАGАТТСААGАААТАGАТТЭGААCGA1TTC СЭТААААСТТТТЭСТЭСАGААСТТТТТАААТАGАТТСААGААGAATGATTTGAACGA1TTC 	300 271
HHV7_RK_U73 F10_020502	ТАТСТОТАСАВССАААТТААЛСАВССЭТСАААТСААСААДААСАААСТСААТААТТСААСТА ТАТСТОТАСАВССАААТТААЛСАВССЭТСА ********************************	360 300
HHV7074		
HHV7_RK_U74 F11_020314	AATAAACAATATAGAGTACCAATTTATCAATGCCTGTTTTTTAATGCTGAGACT-CAC 	117 19
HHV7_RK_U74 F11_020314	ТССТТССАТАСТТТТТТСЭТТАТТСЗАААТСАСАТТСТЗААААТСТССТТСАТАА ТССТТССАТАСТИТТТТСЭТТАТТСЗАААТСАСАТТССТЗААААТСТССТТСАТАА ТССТТССАТАСНИАТАААТНОСТТАТТСЗАААТСАСАТТССТЗААТАА	173 79
HHV7_RK_U74 F11_020314	GATTTCCATTTCAAAAGAAAAAA - TCTTTATGTGGGATTTGAATGAACAAATAATGAGTA GATTTCCATTTCAAAAGAAAAAATCTTTATGTGGGATTTGAATGAA	232 139
HHV7_RK_U74		292

HHV7_RK_U74 F11_020314	ААСТАТССАGААСАТТТАТТТАСТААСААТТАТТТТАААТТСТААТАТGТАТG	352 255
HHV7_RK_U74 F11_020314	TAGATGTCTGCATAGACAGTAATGCGGTTTTGTACATGCCAGACTTAAGACCAATGATAA AANTTGCATAACGTATCANNGAC * ** * * * * *	412 278
HHV7075		
HHV7_RK_U75 F12_020314	ACTCAGCCGAATCTCTTCTAGCGTATCTTCGGAATCAAGAAGAATAATGCCAATGAAGCC TCTCAGCCGAATCTCTTCTAGCGTATCTTCGGAATCAAGAAGAATAATGCCAATGAAGCC	540 60
HHV7_RK_U75 F12_020314	ТАВАААСССТААТВТТТСТСАСАТАТВСТТТТАТАВВАТТТВССАТТТАСААТТВТССС ТАВААСССТААТВТТТСТСАСАТАТВСТТТТАТАВВАТТВССВАТТТАСААТТВТССС ********************************	600 120
HHV7_RK_U75 ⊮12_020314	ATCTTCCAAATGAAACCATACATCATCATCATTCCCAACCGAAATATCAATTAA-AGGTAA ATCTTCCAAATGAAACCATACATCATCATCATCACCGAAAACTATNATNNGATNNNTNN *******************************	657 180
HHV7_RK_U75 F12_020314	ATTTTTAATGGAAACGTGTTCGGCGTGTTTGTTTAGTTCACAGATATTCCGGACGCTTGA ANGGTTATGGCNGNAGTGC-CGGNG * *** ** *** *** *	717 20 4
HHV7U76		
HHV7_RK_U76 G1_020228	AATCAATCTAGAGGAAATACTTTTTGCTAAGGGTGTCAGCGTA-ACCTTGTTTTTTGGTA GTGNCGGNNTTTGGCATACCGTNNTGGN *** * * * * * * *	1379 28
HHV7_RK_U76 G1_020228	ATGTGACAATTGGTGTTATTCCAATCGTACAAATCCAATCAAT	1439 84
HHV7_RK_U76 G1_020228	CGTTATCGAGATT - TTGAAGGATGTAGTTTATCGTTTGCAAAATGCCTTCCAGGATTACC CGTTATCGAGATTCTTGAAGGATGTAGTTTATCGTTTGCAAAATGCCTTCCAGGATTACC	1498 144
HHV7_RK_U76 G1_020228	TGTTTGATTGCATAATACCAGATATTAAAATATTCTTGAAATGACATGCCTTTACTGCGG TGTTTGATTGCATAATACCAGATATTAAAATATTCTTGAAATGACATGCCTTTACTGCGG	1558 204
HHV7_RK_U76 G1_020228	AACGGAAATGCTTTCAGATGACGTTGCCATTCAGCATTGATCTGCTGGATGGA	1618 264
HHV7_RK_U76 G1_020228	TTCAAGAGAGTGTGTGACAGTGTGTAAAACATTTGCTTCTTAAAGACAGCCGAGTCTCTA TTCAAGAGAGTGTGTGACAGTGTGTAAAACATTTGCTTCTTAAAGACAGCCGAGTCTCTA	1678 324
HHV7_RK_U76 G1_020228	AGAACACTGTGTATAGATTGTCCCTTAAAGAATTCTTTTTTACCGTGTAGCATTTCTATA AGAACACTGTGTATAGATTGTCCCTA	1738 350
HHV7077		
HHV7_RK_U77 G2_020425	TTACCTTTTAGAGTATTATTAATAACCGGTACTGCGGGTGCGGGCAAAACAAGCAGCATT 	300 19
HHV7_RK_U77 G2_020425	САЛАССТТАЯСАЯСТААТАЯТИЗАТИЗТСТАТААСТИСТАССАСТТССАТИЗСТИЗСТСАА САЛАССТТАЯСАЯСТААТАЯТИЗТСЭТТАТААСТИСТАССАСТТССАТИЗСТИЗСТСАА	360 79
HHV7_RK_U77 G2_020425	АЛТСТТАОТОЭТТТАТТСААСАСАААССАААТСТССССААСТСАААСААТТТТТАЛААСА АЛТСТТАСТОЭТТТАТТСААСАСАААССАААТСТССССААСТСАААССААТТТТТАЛААСА *********************************	420 139
HHV7_RK_U77 G2_020425	TTTGGTTTTAATAGTTCACATGTATCTATGAATGAAAGAATTAGTTGTTCAGTAACAACT TTTGGTTTTAATAGTTCACATGTATCTATGAATGAAAGAATTAGTTGTTCAGTAACAACT *********	480 199
HHV7_RK_U77 G2_020425	TTAGATTCGATTGCGGATCAGCAAAAACATGATTTATCTACATATTGGAACGTCATCGCA TTAGATTCGATTGCGGATCAGCAAAAACATGATTTATCTACATATTGGAACGTCATCGCA	540 259
HHV7_RK_U77 G2_020425	GATATAGCGGAAAGAGCTTTGAATGCAGCAAATGGGAAAGCCAAAGTGATACCTGATCTA GATATAGCGGAAAGAGCTTTGAATGCGGCAAATGGGAAAGCCAAAGTGA	600 308
HHV7079		
HHV7_RK_U79 G3_020313	ATGATTGCOGAGGATAGAGAATATGGAACGTTTGAATC-TGTAACCCAGGCTTATCAGCA GNAAGCANGANTGCGGGAGGATAGANANAGGNAGAAGNATCGAACCCAGNGNNANCAGCA * * * *** *** * * *** * * **** * ******	59 60
HHV7_RK_U79 G3_020313	GATCATTAGTCATACTTTACAGCTGAGACGATATGAATTTGAAACTGGGTGCATGATTAT GANNNNANANGTCNNCNACANGCGGAGACGANATNNANNAAACGGGGCAGATAGNNN- ** ** ******* * * **** *** **	119 117
HHV7_RK_U79 G3_020313	GTTTTCTGCTAATTCTGGAAAATGTGAGATGCTCTCTAATGGTTGGATTTCAATGATTTC CCGNNAATCGGAAAAGGGAGA-GCTCTCTAA-GGNNGGATTCAAGANNGC * * *** * *** * *** * **** ******** **	179 165

G3_020313	ANGGACT-CAGAAACCGACACGGCCGGCCCAGGGCAN-GNANCCCNACNCGAGGGGNNG * ***** ********** *************	223
ННV7_RK_U79 G3_020313	GCAGTGCAAAACTTACAGTGCCAGAGGTCATATATTATGTTCTAAAAACATCACTTCAAT GCAACACGGC-CGGGCAAAANNACCCCATCCCCCACCCCCANCANCANCNTCCNN *** * * * * * * * * * * * * * * * * *	299 280
HHV7_RK_U79 G3_020313	TTCTCAGRARARAGGAGARAGGARARAGTTTTGACGATTTGTCATGACARTGGARART NTTTCCGNTTTACNTTTNTT	359 300
HHV7081		
HHV7_RK_U81 G4_020425	TAAAGAATTCGGAGGGGAACATCCTCTCACAGTGCTGAAAGCTAATCCGTGACCTCTACC TTGCTGAAAGCTAATCCGTGACCTCTACC ******************************	480 29
HHV7_RK_U81 G4_020425	АТСЭЭGАТААЭGАТСТТЭЭСССАСТАТТАТСАССТТААТСТСТТСАЭЭСЭААСАТАААТА АГСЭЭGАТААЭGАТСТТЭЭСССАСТАТТАТСАССТТААТСТСТТСАЭЭСЭААСАТАААТА *****	540 89
HHV7_RK_U81 G4_020425	ACTCCAGCTGTGTACATTCTGAGGATCGGGGTAGATAATTAAT	600 149
HHV7_RK_U81 G4_020425	ААССААТТТЭТАТАСАТТТТЭТААТТЭТАСААТЭТСАВААТСТВАТААБТТТААААСТТ ААССААТТТЭТАТАСАТТТТЭТААТТЭТАСААТЭТСАВААТСТВАТААБТТТААААСТТ ******	660 209
HHV7_RK_U81 G4_020425	AAGCCACTTCACATTGATITGAAACGITTCATGTTGTTCTTCAAGAGACATTTTTATTGA AAGCCACTTCACATTGATITGAAACGITTCATGITGTTCTTCAAGAGACATTTTTATTGA	720 269
ННV7_RK_U81 G4_020425	GTRGTCATCTGAAATGTTTTCCCAGCATCCACTGTAGTAATGCCAT 765 GTRGTCATCTGAAATGTTTTCCCAGCATCCA 299	
HHV7082		
HHV7_RK_U82 G5_020419	GAGTTTTTCATAAGAGTATGATTTACACACATTATCTCTACAGGTTAAAAGTGTGTTAAC 	360 17
HHV7_RK_U82 G5_020419	TGAAAGACATCTTTCAAAGTTATTTAAAAAGCCAAGCCAGTTGTGCCCTGATTC TAAAATTTTTAAAGGGGAAANAAAGGTTTAAAGGGGGTTGGGTT	414 73
HHV7_RK_U82 G5_020419	TGAATCTTTAATTAAGGAAAGTAAGGTGTGAGTTGATTAGGATTATT	474 131
HHV7_RK_U82 G5_020419	AAATGTTTCTAAAAAATCTTCATCAATTAAAACTT-GATCTCCCTTGAGATATAGGG TTCTTTNNAANNTCTTTTCATCGNAGTTAAAACTNTGATCTCTCCCTTGAGATATAGGG * *** ** * * *****	531 189
HHV7_RK_U82 G5_020419	AG-TATGATATAATAAGCGAGAAATTATATCCGAATTTGGTATAGGTGCCTGATCAAT ANGTATGATATAATAAGCGAGAAATTATATCCGAATTTGGTATAGGTGCCCTGATCGAATN	588 249
HHV7_RK_U82 G5_020419	AGGTTCCAGAAGGGACAGAGTTTTTGATTGGGTACAATTTAG-AATGTTTTTGATAGAAT NGGTTCCAGAAGGGACGGAGTTTTTGATTGGGTACAATTTAGGAATGTTTTTGATAGAAT 	647 309
HHV7_RK_U82 G5_020419	СТТСТТСТАЛАТТАС-АЛТАЛТАЛСАС-ТТАТТАЛА-ТАЛАС-САТАТАТТТССТТ СТТСТТСТАЛАТТАСТАЛАТАЛСАССАСТАТАТТАЛАЛАЛАССАТАТАТТТССТТТ ******	700 369
HHV7_RK_U82 G5_020419	XGAATGGAGATAAGAAATATAAAAA-GATGTTAGTTTTCAT NTAANNNANGNTGATAGATAANATAATCACTGANGTTAATTGCNGCGTNTCGCTCCGTNA * * **** ** *** * **** * **** **	741 429
HHV7_RK_U82 G5_020419	CGCGTTT 436	
HHV7U84		
HHV7_RK_U84 G6_020502	GITAAAATTGCTGTGAACAATACCCTTAGATTTTGTTGCAAAGTTTGTCGTTGAGAGA 	658 29
HHV7_RK_U84 G6_020502	TTTTCTTT TACAGCCTTAAATATTTTTCTCAATTTAGAATGTTTATAGATGGCATTCG TTTTGNNNGTTACAGCCTTAAATATTTTTCTCAATTTAGAATGTTTATAGATGGCATTCG ****	716 89
HHV7_RK_U84 G6_020502	ТАТАТТТЯГСССССТТААСТААСАТТТСКААТТТКАААТСТТАТТСАGAATTTATT ТАТАТТКЯГСССССТТААСТААСАТТТСКААТТТКАААТСЯТТАТТТСАGAATTTATT *******	776 149
HHV7_RK_U84 G6_020502	ТТТСАGGААТТGTACATTTTATGGCTTTTCATTAAACGGTAACGTTTTTGAAAAACAGT ТТТСАGGAATTGTACATTTTATGGCTTTTTCATTAAACGGTAACGTTTTTGAAAAACAGT	836 209
HHV7_RK_U84 G6_020502	ТАСАТОТАТАТСОАССАТТОССТТТАСТАСАССТССАССТТТТАСААСТОАССТТ ТАСАТОТАТАТСОАССАТТОССТТТАСТАСАССССССТТТТАСААСТОАСТТТАССА ******************************	896 269
HHV7_RK_U84 G6_020502	TGGCCTTACAAACAGTTTTACGTTTTTTGCCCCCTCAT 933 TTGGAAACTTACAAACAGTTA 290 * *********	

HHV7U85	
HHV7_RK_U85 G7_020419	AGTCTTGGCGAAGAATGCAACATGTTAAGCAACCAGCATCGTTAATTGTTCGAGATTTAAA 540 TCCAGCATCGTTAATTGTTCGAGATTTAAA 30
HHV7_RK_U85 G7_020419	TTTCAGCAATGCTGTGATATTGTTAGCATTGAAGTAAATAATGCT-TTCTTTGTCACGAT 599 TTTCAGCAATGCTGTGATATTGTTAGCATTGAAGTAAATAATGCTCTTCTTTGTCACGAT 90
HHV7_RK_U85 G7_020419	GTTTAAATTCATGAACTATTTTTTTCACTCATGTTTTTCA-GGTTTTTGGTATTTATGGTA 658 GTTTAAATTCATGAACTATTTTTTCACTCATGTTTTTCACGGTTTTTTGGTATTTATGGTA 150
HHV7_RK_U85 G7_020419	TGATTTAATATCAGTTCGTAATGTTTAAACCATATTACAGAGCTGAGATT-TTTAGTTGT 717 TGATTTAATATCAGTTCGTAATGTTTAAACCATATTACAGAGCTGAGATTCTTTAGTTGT 210
HHV7_RK_U85 G7_020419	ATCAACAGTACACTGAAATACAATTGTGTTTCCTAGTTCTACTCTATGATGCATTATTGG 777 ATCAACAGTACACTGAAATACAATTGTGTTTCCTAGTTCTACTCTATGATGCATTATTGG 270
HHV7_RK_U85 G7_020419	АДАТААТССАТАТАТСАЛТТАСАДААЛАТССАЛТСАТСТТТАТСВАЛАЛТГССАССТААСАА 837 АТАТААТССАТАТАТСАДААЛАТССАЛТСАТСТТТАТСВАЛАЛТГССАССТААСАА 330
HHV7_RK_U85 G7_020419	AGGCAT 843 AGGCAA 336 *****
HHV7U86	
HHV7_RK_U86 G8_020314	TCTGTTGTTGCGACCCCTGTAATGATGTTGTCGATAGGTTGTATCGAACGAA
HHV7_RK_U86 G8_020314	AAGTTCTCTATTTTTTATAGTCCAAATTGTTAAAATATCAACTGAAGATTGGGGTTGATA 3300 AAAAAGAGNANNNNAAAAATTGGAAACAAAGAAAAAANAANANANGAGGGATAAGA 63 ** * **
HHV7_RK_U86 G8_020314	TGGTAAAGAGATAAAATTTTTCTCCACTTGCTTGAGGATAATTATAATACAATGATTGCAG 3360 TGATAAAAAAAAAAAAGGAGGTNGAGGAANANANTATANTGATG 107 ** **** * * ****
HHV7_RK_U86 G8_020314	TCCTTGATGTGGCAATTCTGATGCAAATCGAAATTGTCCAGTTTCAGTTGTATTTGGAGA 3420 -CGTCGATGTGGCATNNGATGCATCNAATGTCGTNNNNNTGNNNNNT-GGAGA 159 * * ******** * ****** ** ****
HHV7_RK_U86 G8_020314	AGATTCAGAAGTAAAAGTACTTCTTGATTTAAGAGGAATAGCACGTCGCGGATCTTGAA-3479 GATCCCCCGNNTNNTCTGGTTGATTTANGAGGAATAGCACGTCGCGGATCTTGANC 215 * * ** ********
HHV7_RK_U86 G8_020314	-GAGTTTTTGATGACTATTTC-TTGGTTGTTAAAATTTGAATAAGTTATTTCATTTTC 3537 NANGTTTTTGATGTGACTATTTCCTTGGTTGTTGTTTTTGACTCCGTNTTTCATTTTCN 272
HHV7_RK_U86 G8_020314	TOGTTTTGTATGATC-GTGAATTTCATTATATAAACCCGAAAGA-GTCATGGCTTC 3591 TGGTTTGTNTGATCCGTGNAATCCCCCCCATAATATNNCCTTTTTTTTTAGTNTGGCTTC 332
HHV7_RK_U86 G8_020314	ACGAACCACGTCAGCTGTAGTGCTCAT
HHV7U90	
HHV7_RK_U90 G9_020502	AGCTAGAATTTAAAGAAAAAAGTCTAAATATGAAATCTGTACACATATATCATATTTAT 3360 CGTCCT-C 7 * *
HHV7_RK_U90 G9_020502	ATGTAAGAAAATGAAAAAAAAAAAAAACCTGCTGCCGGTTTAAATATCCGGGGGAAGCTCCTTCT 3420 AGGNCNCANCGCNCGGGCGNCNCGCATATNCTCATNNNTCCANGAANNGCCNTTNCT 64
HHV7_RK_U90 G9_020502	ТСААСАТТТЭТЭЭААТАЭЭААСАТТТЭССЭТТЭСААТАТСАТТАЭАТЭАТАТТЭАЭТСА 3480 ТСААСАТТТЭТЭЭААТАЭЭААСАТТТЭССЭТТЭСААТАТСАТТАЭАТЭАТЭАТСА 124
HHV7_RK_U90 G9_020502	AGGAAATCAAAAGAATCCAATTCTTOGAAGAGAGTATTATCAAAGGAAGGGTTGCATTTT 3540 AGGAAATCAAAAGAATCCAATTCTTOGAAGAGAGTATTATCAAAGGAAGGGTTGCATTTT 184
HHV7_RK_U90 G9_020502	AAACTGTCTTGCAATATGTCTCTGATGGTCTCTTCCACGGTGGTATCTGGGTGATATCTG 3600 AAACTGTCTTGCAATATGTCTCTGATGGTCTCTTCCACGGTGGTATCTGGGTGATATCTG 244
HHV7_RK_U90 G9_020502	ТАЛТАЛАЛТТТАТАЛАТТТТАТТАСАGАGAЛАТТАЛАТСТGTCTTTTTGTATTAAAATA 3660 ТАЛТАЛАЛТТТАТАЛАЛТТТТАТТАСАGAGAЛАТТАЛАТСТGTCTTTTTGTATTAAAATA 304 ************************************
HHV7_RK_U90 G9_020502	TTTAGGTTTACCTACTCAATATGTCTTTCGTCATTCGATGCCATGGTTTCAGATGGTTGT 3720 TTTAGATTTACCTACTCAATATGTCTTTCGTCATTCGATGCCATGGTTTCAGATGGTTGT 364
HHV7_RK_U90 G9_020502	CCAACAATCAACATAGGTGAAGTAGCTCCACTTCTTTCCAT 3761 CCAA 368 ****

:

HHV7091	
HHV7_RK_U91 G10_020425	ATGTATACTCTGGAATATGAAAAACGTGTATCAAGGCCAAAACTTACTT
HHV7_RK_U91	ТТЭЗСАЛТАТТАТТТЭТТТТТТТАЛТАЛТААСТЭЗАТСТЭТАЛТТЭТААТТЭЗАЛТЭТ 120
G10_020425	ТТЭЗСАЛТАТТАТТЭТТТТТТТАЛТАЛТААСТЭЗАТСТЭТАТТЭЛАТТЭЛАЛТЭЛААСТ 100
HHV7_RK_U91	СТОТСАЛТТСАЛАGAACGACTTTGAATGCTCAAAATGATAAAACAAGTACTGTTGTTCCA 180
G10_020425	СТОТСАЛТТСАЛАGAACGACTTTGAATGCTCAAAATGATAAAACAAGTACTGTTGTTCCA 160
HHV7_RK_U91 G10_020425	GAATTAACATCTAATTCTCCAGGTTTGTTCAGCTATAAAATGTCATTGTTAAAAAAAA
HHV7_RK_U91	GTTTGTAACTCTGTTAATATGAATATTTTTTCTTGTAGATCAAACAACTGTGACAAATTT 300
G10_020425	GTTTGTAACTCTGTTAATATGAATATTTTTTCTTGTAGATCAAACAACTGTGACAAATTT 280
HHV7_RK_U91	TTCTGCAAGTTCTAAACCAACTCTTAGCAGTAAACA-ACCCGGATGGATACAGGCACTAA 359
G10_020425	TTCTGCAAGTTCTAAACCAACTCTTAGCAGTAAACACCCCGGATGA 327
HHV7095	
HHV7_RK_U95	ATGGAGAGTGGGGATAGTTTCGGAAATAATCACCAAGTCTCATCGAATT-CAGAT 54
G11_020314	ATGGGAGGAGTGGGGATAGNNTCGGAAATAAAAAAAAAAA
HHV7_RK_U95	GCTTTTCAATTCCGGCAATATTCAACAACTGTAGATACTTTTGCATATTCATCAATGGAT 114
G11_020314	NCTNTCNAATTCCGGCAATATTCAACAACTGTAGATACNGCATATTCATCAATGGAT 117
HHV7_RK_U95	CCTTCAAATTGCATGCTTAATGAACA-AACTCATTTAGAAACGGTACTAG 163
G11_020314	NNGCTTCAAATTGTCTATGGGGGGGGGGATGNTTATCATAACTCATTTAGAAACGGTACTAG 177
HHV7_RK_U95	TTTATCCTGTTTACAACATG-TCT-TCCCAGCA-AACAGTGGAGAAAAATCTCTATGCCA 220
G11_020314	TTTATCCTGTGTACAACAGGGTCTGTCCCCAGCCCAACAGTGGAGAAAATCTCTATGCCA 237
HHV7_RK_U95	ATAATATAGGAAATGGAAATGGCAGAGTTATTGTAGACAAGATAGTTTGCAAGGAAACC 280
G11_020314	ATAATATAGGAAATGGAAATGTGC-GAGTTATTGTAGANCAGATAGTGNGGGGGGAAGGC 296
HHV7_RK_U95 G11_020314	AAGGAACAT-ATTTTCCAACATATCAGGGATTTCACTCAAACACTCCTCAAAACATTTT 337 GCNNGNANCNTTATTATATAGGTATINNTINGNNTHTTTTTTTT-CTTTGHNNNTAATCCTT 355 * * * * * * * * * * * * * * * * * * *
HHV7_RK_U95 G11_020314	CTATAGAGAATTTTAATTTTGGCACCAATGTCATCAGGCCTATACCATTTAGAAATGAGG 397 TTNTTTTTTTNNTTT370 * * **** ***
HHV7U100	
HHV7_RK_U100	AGAGAATAGTGTAAGCCTGTAGTTACTGACAAGACTTCAGACAGCCCCTTTACTCATAATT 2280
G12_020314	
HHV7_RK_U100	TTTTCTCTTAATCCGCGCACTCCCAATGAGTTCTCTTTTAGTGGAATGCAGAGTGTTGTT 2340
G12_020314	NCNCCCCCTNCTNCCACCANCNCCAAATACANNAAANNAAAAAAGANNAGNGNGGGNA 69
HHV7_RK_U100 G12_020314	TCTAATAATATAACCCTTAGAAGAATTAAAAGTAAGGTATGGAGAAATTTACTTAAAAAGC 2400 AAAANTANNAAAAAAANNNAANACAANAAAATN-GAAATGAGTGTNNNNTNCATAAANC 128 * ** * ** ** ** ** ** * **** * ****
HHV7_RK_U100	TCTTCAAAGTCGTCATATTGT-AGAAATCAGTGTAACCATTTTTACT-TACTGTGCT 2455
G12_020314	NNNATCAGATNNNCATAGTGNTANAAATCNCCCCGNTAACCATTTTTACNCNACTGTGCT 188
HHV7_RK_U100	TCGTCTGGCCATTCACATTCGCAGGAAAACGATGTACATCTTGTGTATTCTAACAAGCTT 2515
G12_020314	TCGTCTGGCCATTCACATTCGCAGGAAAACGATGTACATCTTGTGTATTCTAACAAGCTT 248
HHV7_RK_U100	TGTCTTTCCTCGGAACTTAGACC-AATATATCTAGTAGAGCTTGCGTTGTGCATGTTTTC 2574
G12_020314	TGTCTTTCCTCGGAACTTAGACCCAATATATCTAGTANAGCTTGCGTTGTGCATGTCTTC 308
HHV7_RK_U100	ATTTATTAAGTTTTCGATTA-TATCTTCTAGAGTTATTGCCGTGCGTGCTTCTTT-CA 2629
G12_020314	ATTTNTTTNNNNGTTTTCGATTAATATCTTCTAGAGTTATTGCCGTGCGTGC
HHV7_RK_U100	AGTGGGACCGTATTGGTACATATCTTTATGAGTAATGCGAAAATTATATATGCAGTTGC 2689
G12_020314	AGTGGGACCGTATTGGA

Second attempt Results:

>CL7 = HHV7 C1 (U28) >CL7 = HHV7 C1 (U28) >EM_VI:<u>AF037218</u> AF037218.1 Human herpesvirus 7 strain RK, complete genome. Length = 153,080 Minus Strand HSPs: Score = 1361 (210.3 bits), Expect = 3.0e-53, P = 3.0e-53 Identities = 273/274 (99%), Positives = 273/274 (99%), Strand = Minus / Plus

Query:	317	ATGGAATCAAATTCCGAAGCAGAGTTTGTTTTGAGATAGAACTGGCCTAACAAACGATCA 258
Sbjct:	40404	ATGGAATCAAATTCCGAAGCAGAGTTTGTTTTGAGATAGAACTGGCCTAACAAACGATCA 40463
Query:	257	ACGTGCGGTGCATCATGCATGCTCCCAGTGCTGAAATTATGGCATCACGTAAGCACCAC 198
Sbjct:	40464	ACGTGCGGTGCATCATGCATTGCTCCCAGTGCTGAAATTATGGCATCACGTAAGCACCAC 40523
Query:	197	ATATCTAAGCCCGAATTTAAGCGCCGGGAGAATTTTGAAATTTCATATAACTGCTTATAA 138
Sbjct:	40524	ATATCTAAGCCCGAATTTAAGCGCCGGAGAATTTTTGAAATTTCATATAACTGCTTATAA 40583
Query:	137	AGAAGAGTTTTGTGATGGAAGCGTGAATAGCAAAACCGTTTTTTGTTATATTCATCATGA 78
Sbjct:	40584	AGAAGAGTTTTGTGATGGAAGCGTGAATAGCAAAACCGTTTTTTGTTATATTCATCATGA 40643
Query:	77	CAAAGACATTTCCTCAAGAGACCCATGTTTTAAT 44
SDJCT:	40644	CAAGACATITCUICAAGAGACCCCATGITITIAT 40677
>CL8 = EM_VI: Plus Score = Ident:	HHV-7 AF0372: Lengt Strand 1422 Lies	C12 (U39) 18 AF037218.1 Human herpesvirus 7 strain RK, complete genome. 2h = 153,080 1 HSPs: (219.4 bits), Expect = 5.3e-56, P = 5.3e-56 = 290/297 (97%), Positives = 290/297 (97%), Strand = Plus / Plus
Query:	35	TCACTAGTGATTTTGGCATTGGCAAAGTTGTAACAGTTCGATCTAAGAAATACACAGTG 94
Sbjct:	60735	TGCACTTCTTCAATTGGCATTGGCAAAGTTGTAACAGTTCGATCTAAGAAATACACAGTG 60794
Query:	95	CCAACATCGCGATATGTTGTTTGAAAATGTAAGTTCTTTTTTTAAAAGTTCTGACAGAAAAC 154
Sbjct:	60795	CCAACATCGCGATATGTTGTTTGAAATGTAAGTTCTTTTTTTT
Query:	155	GTGTGTGCTTCAATCTTTGTTTGTAAATTATCAAAATACCTTCGGTAGTTTTAATGTTA 214
Sbjct:	60855	GTGTGTGCTTCAATCTTTGTTTTGTAAATTATCAAAATACCTTCGGTAGTTTTAATGTTA 60914
Query:	215	GAGCCGTAAGACGCGCAAGAAACCTCCCCTGTCAAAACGCACTAAATCTGTCCCGGTGGCA 274
Sbjct:	60915	GAGCCGTAAGACGCGCAAGAAACCTCCCTGTCAAAACGCACTAAATCTGTCCCCGGTGGCA 60974
Sbict:	60975	
>CL9 = >EM_VI Minus Score : Ident	HHV7 I AF0377 Lengt Strat = 1136	D4 (U43) 218 AF037218.1 Human herpesvirus 7 strain RK, complete genome. th = 153,080 nd HSPs: (176.5 bits), Expect = 4.4e-43, P = 4.4e-43 = 228/229 (49%), Expect = 9.28/229 (9%), Strand - Minus (Plus
Query:	229	AGTAAAAAAATCTTTATACAGATTTTCTGTTTCTTCCGTATTTAAACAGTGAATAATATCC 170
Sbjct:	71003	AGTAAAAAATCTTTATACAGATTTTCTGTTTCTCCGTATTTTAAACAGTGAATAATATCC 71062
Query:	169	TCAGACTTAATGGGTAATGCATTTTTTACATAATAGCTGAAATTCAAAGATTCTGCGTCG 110
Sbjct:	71063	TCAGACTTAATGGGTAATGCATTTTTTACATAATAGCTGAAATTCAAAGATTCTGCGTCG 71122
Query:	109	CAAACAAATACCGTCTCAATTCTTTTGGAAAACTTGCATTCTGTGTCTGCATACAAAAA 50
Sbjct:	71123	CARACARATACCGTCTCAATTCITTTGGAARACTTGCATTTCTGTGTCTGCATACARARA 71182
Query:	49	TATACATTTTTGATGGTTTATATTTAACGATTATCGGAAAAATTAAAT 1
Sbjct:	71183	TATACATTITTIGATGGTTTATATTTAACGATTATTGGAAAAATTAAAT 71231
>CL10	• HHV7	E8 (U57) 218 AF037218.1 Human herpesvirus 7 strain RK, complete genome.
Minu Score Ident	Leng S Stra 542 ities	th = 153,080 nd HSPs: (87.4 bits), Expect = 3.0e-16, P = 3.0e-16 = 110/112 (98%), Positives = 110/112 (98%), Strand = Minus / Plus
Query:	112	AAATCTATCCATTCAGTCCGATTACAGTAAATCCCCAGGTAATGCTTCGAAACTTATGTTA 53
Sbjct:	91585	AAATCTATCCATTCAGTCCGATTACAGTAAATCCCAAGTAATGCTTCGAAACTTATGTTA 91644

Appendix B – Identities of lymphochip clones on custommade microarrays

Plate	Row	Column	Well	Clone ID	Name	GB Acc#
AP-bu01	Δ	1	A1	67037	Homo sapiens transcribed sequence with weak similarity to protein	T70413
	<u>^</u>	•	~	0,00,	ref NP 0602651 (H seriens) hyrothetical protein EL [20378 [Homo seriens]	
		•	••	040400		N72214
AP-hu01	A	2	A2	246430	UDP glycosyltransferase 2 family, polypeptide 54	N/ 3214
AP-hu01	Α	3	A3	67071	interleukin 22 receptor, alpha 1	T70439
AP-hu01	Δ	4	A4	257298	0	N39874
AD 1. 04	2	2		440000	state a headbatase 2 (formaty 24) regulatory subusit B" alaba	T9/122
AP-nu01	A	5	AD	110393	protein phosphatase 2 (tormeny 2A), regulatory suburnt D , alpha	104132
AP-hu01	Α	6	A6	257581	matrix metalloproteinase 12 (macrophage elastase)	N41372
AP-hu01	A	7	A7	111460	cyclin T1	T90767
1001	?			057740		NA1940
AP-nu01	A	8	Að	20//40	Homo sapiens transcribed sequences	1441040
AP-hu01	Α	9	A9	111721	insulin induced gene 2	T91091
AP-hu01	Δ	10	A10	258790	cholecystokinin	N40841
	· ·			440440		T00057
AP-hu01	A	11	A11	112148	8	192007
AP-hu01	Α	12	A12	263341	DKFZP566K1924 protein	N20003
	۸	13	A13	113786	hypothetical protein MGC45562	T77343
1001	?			000000		100400
AP-hu01	A	14	A14	263836	Zinc tinger protein 24 (KOA 17)	N20492
AP-hu01	Α	15	A15	114045	Homo sapiens transcribed sequences	T79505
AP-bu01	Δ	16	A16	264287	CCAAT-box-binding transcription factor	N27463
40 5 04	· · ·		A 4 7	444040		T70507
AP-nu01	A	17	A17	114048	eryuropoleun receptor	179007
AP-hu01	Α	18	A18	264530	Pirin	N29255
AP-hu01	Δ	19	A19	114094	cytochrome P450, family 2, subfamily C, polypeptide 8	T79539
	7		400	005550		N07804
AP-nu01	A	20	A20	200003		N2/004
AP-hu01	Α	21	A21	114807	ATP-binding cassette, sub-family G (WHITE), member 2	T87360
AP-hu01	A	22	A22	266074	caldesmon 1	N31556
AD 1:001	2		400	445044	autorina autoria hinding protein (Constein), sinhe 15 (Coloss)	T07260
AP-nu01	A	23	A23	113014	guarine nucleotide binding protein (G protein), alpha 15 (Gd class)	10/200
AP-hu01	Α	24	A24	268412	MCF.2 cell line derived transforming sequence	N35192
	8	1	B1	358379	associated molecule with the SH3 domain of STAM (AMSH) like protein	W95888
		÷	00	703007		A A 404727
AP-hu01	в	2	82	121301	thyroid hormone receptor interactor 13	AA401737
AP-hu01	в	3	B 3	358433	retinoid X receptor, gamma	W96099
AD 6001	<u> </u>	A	B4	727705	la ctotransferrin	44398686
	-	-		121100		101000000
AP-hu01	в	5	85	358506	collagen, type XVI, alpha 1	W96115
AP-hu01	в	6	B6	728707	Homo sapiens transcribed sequences	AA435905
AD bu01	- -	7	87	350134	transforming acidic colled-coil containing protein 1	AA010120
AF-IIUU I		1	57	009104	anatoming, active concercon containing protein 1	
AP-hu01	в	8	B8	729912	DNA (cytosine-5-)-methyltransferase 2	AA399591
AP-hu01	в	9	B9	359617	aldehvde dehvdrogenase 9 family, member A1	AA010649
AD hu01	- -	10	B10	730346	heme hinding protein 1	AA470002
AF-IIUUI	-	10	BIU	730340	name binding protein 1	
AP-hu01	в	11	811	359769	RAB interacting factor	AA010789
AP-hu01	в	12	B12	730397	TATA box binding protein	AA470112
AD bu01	-	13	B13	360531	versf murine serroms 3611 virel oncorene homolog 1	44015618
	-	13	513	300331		2010010
AP-hu01	в	14	B14	730950	chromosome 3 open reading frame 8	AA421118
AP-hu01	в	15	B15	360723	peroxiredoxin 6	AA016074
AD bu01	- -	16	B16	731995	chomosome 1 open reading frame 16	AA421062
AP-NUUT	B	10	DIO	131223		AA421002
AP-hu01	в	17	817	362942	DIRAS family, GTP-binding RAS-like 2	AA019438
AP-hu01	в	18	B18	731592	protein phosphatase 1, regulatory (inhibitor) subunit 8	AA412525
AD build		10	P10	264557	hypothesisal protein MCC9721	A A022602
AP-nu01	в	19	DIA	304557	nypotnetical protein MGC6721	AAU22003
AP-hu01	в	20	B20	739891	E74-like factor 3 (ets domain transcription factor, epithelial-specific)	AA477878
AP-hu01	в	21	B21	365098	BCI 2/adenovirus E1B 19kDa interacting protein 3-like	AA025195
AD hu04	5		800	740400	HOA bistone family member V	A A 470000
AP-NUUT	D	22	DZZ	740406	HZA historie ramity, member A	MM410223
AP-hu01	в	23	B23	365452	chromosome 3p21.1 gene sequence	AA009799
AP-hu01	в	24	B24	741406	Ras and Rab interactor 1	AA400958
	~			440400		T04 440
AP-nu01	C	1	CI	110430	transcriptional adaptor 2 (ADA2 nomolog, yeast)-like	191410
AP-hu01	С	2	C2	269800	hypothetical protein FLJ10116	N36276
AP-hu01	С	3	C3	120386	0	T95837
AD 5001	č		64	070646	Res Deretein	N/42900
AP-nuu i	U.	4	C 4	270010	Rag D protein	1442033
AP-hu01	С	5	C5	121454	Homo sapiens transcribed sequence	AF143883
AP-hu01	С	6	C6	270939	catenin (cadherin-associated protein), beta 1, 88kDa	N42740
AD 6001	~	7	67	101049	interformer related developmental regulator 1	T07969
	č			121340		191000
AP-hu01	C	8	C8	2/1051	pnosphodiesterase 40, CAMP-specific (phosphodiesterase E3 dunce homolog,	N42967
					Drosophila)	
AP-hu01	С	9	C9	122112	beta-transducin repeat containing	T98478
AP-hu01	C	10	C10	271085	tyrosinese (oculocutaneous albinism IA)	N42770
	č	44	0.0	104000	interleubin 11 recenter einhe	000700
AP-0001	C	33	CII	124086	inteneukin 11 receptor, alpha	RU2/08
AP-hu01	С	12	C12	276501	hypothetical gene supported by AK026880; BC030095; AF052138	N43917
AP_hu01	С	13	C13	124708	vacuolar protein sorting 4B (veast)	R02155
	ž		010	077050	hundhatiaal arataia El (10701	NOOFOO
AP-0001	U	14	G14	∠1/058	nypometical protein FLU 10701	N39282
AP-hu01	С	15	C15	126348	transmembrane 4 superfamily member 4	R06489
AP-hu01	C	16	C16	277138	protein phosphatase 2 (formerly 2A) regulatory subunit B (PR 52), gamma isoform	N40919
AD LOOK	č	17	~~~	107000	T call caute lumphonetic laukemic 4	DOGOEO
AP-nu01	C	17	017	12/080	r-cen acute rymphocyuc reukernia i	600020
AP-hu01	С	18	C18	277359	phosphoglycerate mutase 2 (muscle)	N57500
AP-hu01	С	19	C19	127841	pyruvate kinase, liver and RBC	R08829
	ž	20	000	070057	Aumon prototo pE2 inducible publicar prototo 4	NO4800
AP-NU01	C	20	C20	2/825/	tumor protein pos inducible nuclear protein 1	N94892
AP-hu01	С	21	C21	128065	BH3 interacting domain death agonist	R09650
AP-hu01	C	22	C22	278448	Homo sapiens transcribed sequences	N98717
	ž	~~	022	400400		D00707
AP-hu01	C	23	C23	128129	conagen, type II, alpha 1 (primary osteoarthritis, spondyloepiphyseal dysplasia,	RU9767
					congenital)	
AP-hu01	С	24	C24	279665	E3-binding protein	N49046
AP-hu01	D	1	D1	365837	G-rich RNA sequence binding factor 1	AA026052
	5		5	740040		A A 404 CO 1
AP-Nu01	U	Z	02	/42643	v	AA401504
AP-hu01	D	3	D3	365866	titin	AA025367
AP-hu01	D	4	D4	742747	canicua homolog (Drosophila)	AA400572
	5	-		200705	sharmanana 01 anan sanding frame 6	A A O O O A O ~
AP-NU01	U	5	05	300/85	chromosome 21 open reading trame 6	AAU29407
AP-hu01	D	6	D6	743233	polymerase (DNA directed), delta 3	AA401552

	D	7	D7	376147	RAB5C, member RAS oncogene family	AA039472
AP-hu01	D D	8	D8	743316	0	AA400583
AP-hu01	D	9	D9	376394	protein phosphatase 4 (formerly X), catalytic subunit	AA041409
AP-hu01	Ď	10	D10	752754	hypothetical protein LOC284611	AA482455
AP-hu01	D	11	D11	376405	chromosome 6 open reading frame 4	AA041420
AP-hu01	Ď	12	D12	753278	chemokine (C-X-C motif) ligand 16	AA411655
AP-hu01	D	13	D13	376522	gap junction protein, beta 1, 32kDa (connexin 32, Charcot-Marie-Tooth neuropathy,	AA041502
					X-linked)	
AP-hu01	D	14	D14	753284	glutathione S-transferase M5	AA411662
AP-hu01	D	15	D15	376708	hypothetical protein BC013995	AA046358
AP-hu01	D	16	D16	753969	discs, large (Drosophila) homolog 5	AA478949
AP-hu01	D	17	D17	377282	stearoyt-CoA desaturase (delta-9-desaturase)	AA054956
AP-hu01	D	18	D18	754054	homeo box D9	AA478764
AP-hu01	D	19	D19	382913	v-ral simian leukemia viral oncogene homolog B (ras related; GTP binding protein)	AA084543
AP-hu01	D	20	D20	755663	retinoic acid receptor, beta	AA419238
AP-hu01	D	21	D21	383156	chloride channel 4	AA071201
AP-hu01	D	22	D22	756208	retinoic acid receptor, gamma	AA481863
AP-hu01	D	23	D23	415293	kininogen	W92148
AP-hu01	D	24	D24	756557	thiosulfate sulfurtransferase (rhodanese)	AA481682
AP-hu01	Ε	1	E1	128274	zinc finger protein, subfamily 1A, 5 (Pegasus)	R11517
AP-hu01	Ε	2	E2	280231	coagulation factor C homolog, cochlin (Limulus polyphemus)	N49197
AP-hu01	E	3	E3	128667	chemokine (C-C motif) ligand 14	R16807
AP-hu01	E	4	E4	280758	thymosin, beta 4, Y chromosome	N50611
AP-hu01	E	5	E5	128921	sterol O-acyltransferase 2	R10273
AP-hu01	Е	6	E6	280868	Homo sapiens transcribed sequence with strong similarity to protein pir:S71101	N47532
AP-hu01	E	7	E7	129585	(H.sapiens) S/1101 protein kinase JNK2 isoform beta-1 - human Homo sapiens transcribed sequence with moderate similarity to protein sp:P02795 (H.sapiens) MT2 HUMAN Metallothionein-II (MT-II)	R16596
AP-hu01	Е	8	E8	281043	Sapiens cDNA FLJ37676 fis, clone BRHIP2012627.	N50903
AP-hu01	Е	9	E9	129900	0	R19169
AP-hu01	Е	10	E10	281949	phosphoinositide-binding protein PIP3-E	N53325
AP-hu01	E	11	E11	130313	nitric oxide synthase 3 (endothelial cell)	R21248
AP-hu01	E	12	E12	284547	corticotropin releasing hormone binding protein	N76172
AP-hu01	Е	13	E13	132660	aldehyde dehydrogenase 3 family, member B2	R25587
AP-hu01	Е	14	E14	288654	general transcription factor IIE, polypeptide 2, beta 34kDa	N79351
AP-hu01	Е	15	E15	136549	Homo sapiens transcribed sequences	R34490
AP-hu01	Е	16	E16	288663	gap junction protein, beta 1, 32kDa (connexin 32, Charcot-Marie-Tooth neuropathy,	N79360
					X-linked)	
AP-hu01	E	17	E17	136919	chemokine (C-C motif) ligand 28	R38459
AP-hu01	E	18	E18	289268	S100 calcium binding protein A3	N68995
AP-hu01	E	19	E19	137297	LOC151201	R36598
AP-hu01	Е	20	E20	289459	pyruvate carboxylase	N76699
AP-hu01	E.	21	E21	137995	low density lipoprotein receptor-related protein 8, apolipoprotein e receptor	R63185
AP-hu01	Е	22	E22	289844	phosphatidylinositol glycan, class A (paroxysmal nocturnal hemoglobinuria)	N77115
AP-hu01	E	23	E23	139125	0	R62517
AP-hu01	E	24	E24	289981	0	N79966
AP-hu01	F	1	F1	415818	LOC253392	W84813
AP-hu01	F	2	F2	756596	dual specificity phosphatase 4	AA444068
AP-hu01	F	3	F3	415851	ras homolog gene family, member E	W86326
AP-hu01				756876	solute carrier family 35 (UDP-galactose transporter), member A2	AA428423
	F	4	F4			
AP-hu01	F	4 5	F4 F5	416803	Homo sapiens transcribed sequences	W86776
AP-hu01 AP-hu01	F F F	4 5 6	F4 F5 F6	416803 756965	Homo sapiens transcribed sequences regulator of G-protein signalling 14	W86776 AA428897
AP-hu01 AP-hu01 AP-hu01	F F F	4 5 6 7	F4 F5 F6 F7	416803 756965 417258	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2	W86776 AA428897 W87943
AP-hu01 AP-hu01 AP-hu01 AP-hu01	FFFF	4 5 6 7 8	F4 F5 F6 F7 F8	416803 756965 417258 757718	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma	W86776 AA428897 W87943 AA442448
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	F F F F F	4 5 6 7 8 9	F4 F5 F6 F7 F8 F9	416803 756965 417258 757718 417361	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain	W86776 AA428897 W87943 AA442448 W89211
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	FFFFFF	4 5 6 7 8 9 10	F4 F5 F6 F7 F8 F9 F10	416803 756965 417258 757718 417361 757875	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35)	W86776 AA428897 W87943 AA442448 W89211 AA442854
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	F F F F F F F	4 5 6 7 8 9 10 11	F4 F5 F6 F7 F8 F9 F10 F11	416803 756965 417258 757718 417361 757875 417435	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995	W86776 AA428897 W87943 AA442448 W89211 AA442854 W89047
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	F F F F F F F F F	4 5 6 7 8 9 10 11 12	F4 F5 F6 F7 F8 F9 F10 F11 F12	416803 756965 417258 757718 417361 757875 417435 757968	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter	W86776 AA428897 W87943 AA442448 W89211 AA442854 W89047 AA442633
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	F F F F F F F F F F	4 5 6 7 8 9 10 11 12 13	F4 F5 F6 F7 F8 F9 F10 F11 F12 F13	416803 756965 417258 757718 417361 757875 417435 757968 417801	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27	W86776 AA428897 W87943 AA442448 W89211 AA442854 W89047 AA442633 W88848
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	F F F F F F F F F F -	4 5 6 7 8 9 10 11 12 13 14	F4 F5 F6 F7 F9 F10 F11 F12 F13 F14	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein	W86776 AA428897 W87943 AA442488 W89211 AA442854 W89047 AA442633 W88848 AA403237
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	F F F F F F F F F F F F F	4 5 6 7 8 9 10 11 12 13 14 15	F4 F5 F6 F7 F8 F10 F11 F12 F13 F14 F15	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic)	W86776 AA428897 W87943 AA442448 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90535
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	F F F F F F F F F F F	4 5 6 7 8 9 10 11 12 13 14 15 16	F4 F5 F6 F7 F8 F10 F11 F12 F13 F14 F15 F16	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation cross	W86776 AA428897 W87943 AA442448 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90535 AA496054
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	FFFFFFFFFFFFFFFFFFFFFFFFFFFFFFFFFFFFFF	4 5 6 7 8 9 10 11 12 13 14 15 16 17	F4 F5 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17	416803 756965 417258 757718 417361 757875 417435 757968 417435 757968 417801 758534 418013 759207 418040	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGEB-induced factor (TALE family homeobox)	W86776 AA428897 W87943 AA442448 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90535 AA496054 W90532
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	F F F F F F F F F F F F F F F F F F F	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18	F4 F5 F6 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F18	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome)	W86776 AA428897 W87943 AA442448 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90535 AA496054 W90532 AA496054
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	F F F F F F F F F F F F F F	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19	F4 F5 F6 F7 F8 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 422750	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrilin 1 (Marfan syndrome) spectrin, albha, erythrocytic 1 (elliptocytosis 2)	W86776 AA428897 W87943 AA442848 W89211 AA442848 W89047 AA442633 W88848 AA403237 W90535 AA496054 W90532 AA423948 AA01897
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	F F F F F F F F F F F F F F F F F F F	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20	F4 F5 F6 F7 F8 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinie cell protein 4	W86776 AA428897 W87943 AA442488 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90535 AA496054 W90532 AA429948 AA001897 AA424561
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	÷ + + + + + + + + + + + + + + + + + + +	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20	F4 F5 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21	416803 756965 417258 757718 417361 757875 417435 757968 417801 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 oranzyme K (serine protease, granzyme 3; troptase II)	W86776 AA428897 W87943 AA442488 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90535 AA496054 W90532 AA423948 AA001897 AA424561
AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01 AP-hu01	卡卡卡卡卡卡卡卡卡卡卡卡卡卡卡卡	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22	F4 F5 F6 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22	416803 756965 417258 757718 417361 757875 417435 757968 417301 758534 418033 759207 418040 759881 427750 767174 428412 767955	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (hilateral acoustic neuroma)	W86776 AA428897 W87943 AA4428448 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90535 AA496054 W90532 AA423948 AA005382 AA424561 AA005382
AP-hu01 AP-hu01		4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23	F4 F5 F6 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FL10420	W86776 AA428897 W87943 AA442848 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90532 AA496054 W90532 AA423948 AA005382 AA424561 AA005382 AA418463
AP-hu01 AP-hu01		4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	F4 F5 F6 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23 F24	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibriliin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein	W86776 AA428897 W87943 AA442488 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90535 AA496054 W90532 AA429661 AA005382 AA418463 AA033649 AA418465
AP-hu01 AP-hu01	בבבבבבבבבבב בבבבבב	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1	F4 F5 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23 F24 G1	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein	W86776 AA428897 W87943 AA442448 W89047 AA442633 W8848 AA403237 W90535 AA496054 W90532 AA423948 AA403948 AA403849 AA424561 AA03649 AA418463 AA03649 AA418463
AP-hu01 AP-hu01	בבבבבבבבבב בבבבבבב	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2	F4 F5 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23 F24 G1 G2	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neuritin 1	W86776 AA428897 W87943 AA4428448 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90532 AA42948 AA405382 AA42948 AA001897 AA424561 AA005382 AA418463 AA418463 AA418465 R67704 N64489
AP-hu01 AP-hu01	רבבבבבבבבבבבבבבבבבבבבבבבבבבבבבבבבבבבב	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3	F4 F5 F6 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23 F24 G1 G2 G2 G2 G3	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neuritin 1 Homo sapiens transcribed sequences neuropilin 2	W86776 AA428897 W87943 AA4428448 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90532 AA496054 W90532 AA496054 W90532 AA423948 AA005382 AA424561 AA005382 AA4184653 R67704 N64489 R67834
AP-hu01 AP-hu01	בבבבבבבב בבבבב בבבבבבב	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3 4	F4 F5 F6 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23 F24 G1 G2 G3 G4	416803 756965 417258 757718 417351 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536 290517	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine proteases, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neuritin 1 Homo sapiens transcribed sequences neuropilin 2	W86776 AA428897 W87943 AA442848 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90532 AA496054 W90532 AA496054 W90532 AA423948 AA005382 AA424561 AA005382 AA418463 AA033649 AA418455 R67704 N64489 R67834 N79133
AP-hu01 AP-hu01	בבבבבבבבב בבבבבב בבבבבב	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 9 20 21 22 23 4 1 2 3 4 5	F4 F5 F6 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23 F24 G1 G2 G3 G4 G5	416803 756965 417258 757718 417358 757718 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536 290517 141815	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrilin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neuritin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome)	W86776 AA428897 W87943 AA442448 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90535 AA496054 W90532 AA423948 AA001897 AA424561 AA005382 AA418463 AA005382 AA418463 AA033649 AA418455 R67704 N64489 R67834 N79133 R70685
AP-hu01 AP-hu01	רברברברברברב ביביבים ממסמס מס	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 22 23 4 1 2 3 4 5 6	F4 F5 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23 F24 G1 G2 G3 G4 G5 G6	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536 290517 141815 290549	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein Nutritin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome)	W86776 AA428897 W87943 AA442488 W89211 AA442854 W89047 AA442633 W98848 AA403237 W90535 AA496054 W90532 AA429615 AA429615 AA42561 AA001897 AA424561 AA003849 AA418455 R67704 N64489 R67834 N79133 R70685 AF088104
AP-hu01 AP-hu01	רברברברברברב בבבנים מסממס מס	4 5 6 7 8 9 10 11 12 13 14 5 6 7 22 23 4 1 2 3 4 5 6 7	F4 F5 F6 F7 F8 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23 F24 G1 G2 G3 G4 G5 G6 G7	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536 290517 141815 290549 142283	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neuritin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome) Gem-interacting protein FLJ3036	W86776 AA428897 W87943 AA4428448 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90532 AA42633 W90532 AA423948 AA4005382 AA423948 AA005382 AA424561 AA005382 AA418465 R67704 N64489 R67704 N6489 R67834 N79133 R70685 AF086104 R69925
AP-hu01 AP-hu01	בבבבבבים מממממ משוון הביב בבבב בבים מממממ ממ	4 5 6 7 8 9 10 11 12 13 14 5 6 7 8	F4 F5 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F23 F24 G1 G2 G4 G5 G6 G6 G8	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536 290517 141815 290549 142383 291426	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) exclsion repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine proteases, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neurtitin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome) Gem-interacting protein hypothetical protein FLJ35036 acetyl-Coenzyme A synthetase 2 (AMP forming)-like	W86776 AA428897 W87943 AA442848 W89211 AA442854 W89047 AA442633 W98532 AA496054 W90532 AA496054 W90532 AA496054 W90532 AA423948 AA005382 AA418465 R67704 N64489 R67834 N79133 R70685 AF086104 R6925 W03611
AP-hu01 AP-hu01	רברברברברברב וברברברברססססססס	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 9 21 22 3 4 5 6 7 8 9	F4 F5 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23 F21 F22 G3 G2 G3 G2 G3 G5 G6 G7 G9 G9	416803 756965 417258 757718 417351 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536 290517 141815 290549 142383 291426 142244	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrilin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neuritin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome) Gem-interacting protein hypothetical protein FLJ35036 acetyl-Coenzyme A synthetase 2 (AMP forming)-like chromosome 6 open reading frame 37	W86776 AA428897 W87943 AA442848 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90535 AA496054 W90532 AA423948 AA001897 AA424561 AA005382 AA418463 R67704 N64489 R67834 N79133 R70685 AF086104 R69925 W03611 R71124
AP-hu01 AP-hu0	רברברברברברב ברברברברם0000000	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 21 22 34 5 6 7 8 9 10	F4 F5 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23 F24 G1 G2 G3 G4 G5 G6 G7 G8 G9 G10	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536 290517 141815 290549 142833 291426 142944 292212	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein neuritin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome) Gem-interacting protein hypothetical protein FLJ35036 acetyl-Coenzyme A synthetase 2 (AMP forming)-like chromosome 6 open reading frame 37 ATP-binding cassette, sub-family C (CFTR/MRP), member 5	W86776 AA428897 W87943 AA442488 W89211 AA442854 W89047 AA442633 W98848 AA403237 W90535 AA496054 W90532 AA423948 AA403237 W90532 AA423948 AA40303649 AA418455 R67704 N64489 R67834 N79133 R70685 AF086104 R69925 W03611 R71124 N80617
AP-hu01 AP-hu01	רברברברברברבר ברברברברססססססססס	4 5 6 7 8 9 10 11 12 13 4 15 6 7 8 9 10 11 22 23 4 5 6 7 8 9 10 11	F4 F5 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23 F24 G2 G3 G4 G5 G6 G7 G8 G90 G11	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536 290517 141815 290549 142383 291426 142944 292212 143887	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neuritin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome) Gem-interacting protein FLJ35036 acetyl-Coenzyme A synthetase 2 (AMP forming)-like chromosome 6 open reading frame 37 ATP-binding cassette, sub-family C (CFTR/MRP), member 5 protease, serine, 23	W86776 AA428897 W87943 AA442848 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90532 AA42633 W90532 AA423948 AA4005382 AA424561 AA005382 AA424561 AA005382 AA418465 R67704 N64489 R67834 N79133 R70865 AF086104 R69925 W03611 R71124 N80617 R76394
AP-hu01 AP-hu01		4 5 6 7 8 9 10 11 12 13 14 5 6 7 8 9 10 11 22 23 4 5 6 7 8 9 10 11 22 23 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 13 4 15 6 7 8 9 10 11 12 13 4 15 6 7 8 9 10 11 12 13 4 15 6 7 8 9 10 11 12 13 4 15 6 7 8 9 10 11 12 13 4 15 6 7 8 9 10 11 12 13 4 15 16 17 18 19 10 11 12 13 4 15 16 17 18 19 10 11 12 13 14 15 16 17 18 19 10 11 12 13 14 15 16 17 18 19 20 1 22 23 24 11 2 23 24 12 23 24 12 2 3 24 12 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	F4 F5 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F23 F24 G1 G2 G3 G4 G5 G6 G7 G8 G9 G11 G12	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418031 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536 290517 141815 290549 142283 291426 142944 292212 143887 292699	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLU10420 KIAA0251 protein neuritin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome) Gem-interacting protein hypothetical protein FLJ35036 acetyl-Coenzyme A synthetase 2 (AMP forming)-like chromosome 6 open reading frame 37 ATP-binding cassette, sub-family C (CFTR/MRP), member 5 protease, serine, 23 ankyrin 1, erythrocytic	W86776 AA428897 W87943 AA442848 W89211 AA442854 W89047 AA442633 W98532 AA496054 W90532 AA496054 W90532 AA496054 W90532 AA423948 AA005382 AA424561 AA005382 AA418465 R67704 N64489 R67834 N79133 R70685 AF086104 R69925 W03611 R71124 N80617 R76394 N80475
AP-hu01 AP-hu01		4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 9 21 22 3 4 5 6 7 8 9 10 11 23 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 14 15 16 7 8 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 11 12 3 12 2 12 2 3 12 3 12 3 12 3 1	F4 F5 F6 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F23 F24 G12 G34 G5 G6 G9 G10 G11 G12 G13	416803 756965 417258 757718 417351 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767174 428412 767295 470958 767302 140197 290340 140536 290517 141815 290549 142833 291426 142944 292212 143887 292299 144797	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neuritin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome) Gem-interacting protein hypothetical protein FLJ35036 acetyl-Coenzyme A synthetase 2 (AMP forming)-like chromosome 6 open reading frame 37 ATP-binding cassette, sub-family C (CFTR/MRP), member 5 protease, serine, 23 ankyrin 1, erythrocytic	W86776 AA428897 W87943 AA442848 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90532 AA496054 W90532 AA496054 W90532 AA423948 AA005382 AA424561 AA005382 AA418463 AA033649 AA418465 R67704 N64489 R67834 N79133 R70685 AF086104 R6925 W03611 R71124 N80617 R76394 N80475 R76553
AP-hu01 AP-hu0	רברברברברברבר ברברברברססססססססססס מ	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 21 22 23 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 7 8 9 10 11 12 3 4 15 6 7 7 8 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 11 12 3 14 15 16 17 11 2 2 3 4 5 16 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 11 12 3 11 12 3 11 12 3 12 2 12 2 3 4 5 6 7 8 9 10 11 12 3 12 2 12 2 3 4 12 3 4 5 6 7 8 9 10 11 12 3 2 12 2 3 4 5 6 7 8 9 10 11 12 3 3 4 5 6 7 8 9 10 11 12 3 11 2 3 4 5 6 7 8 9 10 11 12 3 1 12 3 1 12 3 11 12 3 12 1 12 1 12 1 12 1 12 1 12 1 12 1 12 1 12 1 12 1 12 1 12 1 1 1 1 1 1 1 11 1	F4 F5 F6 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F22 F23 F24 G12 G3 G4 G5 G6 G7 G8 G10 G11 G12 G13 G13	416803 756965 417258 757718 417358 757718 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536 290517 141815 290549 142833 291426 142844 292212 143887 292699 144797	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrilin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neuritin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome) Gem-interacting protein hypothetical protein FLJ35036 acetyl-Coenzyme A synthetase 2 (AMP forming)-like chromosome 6 open reading frame 37 ATP-binding cassette, sub-family C (CFTR/MRP), member 5 protease, serine, 23 ankyrin 1, erythrocytic a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 1	W86776 AA428897 W87943 AA442848 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90535 AA496054 W90532 AA423948 AA001897 AA424561 AA005382 AA418463 AA005382 AA418463 AA033649 AA418455 R67704 N64489 R67834 N79133 R70685 AF086104 R69925 W03611 R71124 N80617 R76394 N80617 R76394 N806753
AP-hu01 AP-hu01	בבבבבים ממממממממממ מישר בבבב בבב הבישר בבבב בבבב בבבב בבבב בבבב בבבב בבבב	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 21 22 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 13 4 15 16 7 8 9 10 11 12 13 4 15 16 7 8 9 10 11 12 13 4 15 16 17 8 9 10 11 12 13 4 15 16 17 18 19 10 11 12 13 4 15 16 17 18 19 10 11 12 13 4 15 16 17 18 19 10 11 12 13 4 15 16 17 18 19 10 11 12 13 4 15 16 17 18 19 10 11 12 13 14 15 16 17 18 19 10 11 12 13 14 15 16 17 18 19 10 11 12 13 14 15 16 17 18 19 10 11 12 13 14 15 16 17 18 19 10 11 12 13 14 15 16 17 18 19 10 11 12 13 14 15 16 17 18 19 10 11 12 13 14 15 16 11 12 13 14 15 16 17 18 19 20 12 23 24 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 1 12 3 4 5 6 7 8 9 10 11 11 12 3 1 11 12 3 1 11 11 12 11 11 11 11 11 11 11 11 11 1	F4 F5 F6 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F22 F23 F24 G2 G3 G4 G6 G7 G8 G90 G11 G12 G13 G14 G14	416803 756965 417258 757718 417358 757718 417351 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536 290517 141815 290549 142383 291426 142383 29157 290597 142383 291426 142383 291426 142383 291426 142383 291426 142383 291426 142383 291426 142383 291426 142383 291426 142383 291426 142383 291426 142383 291426 142383 291426 142383 291426 142383 291426 142383 291426 142383 291426 144797 292340 144797 292340 144797 292340 144797 292340 144797 292340 144797 292340 144797 292340 144797 292367 292567 292577 292577 292577 2925777 29257777777777	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neuritin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome) Gem-interacting protein hypothetical protein FLJ35036 acetyl-Coenzyme A synthetase 2 (AMP forming)-like chromosome 6 open reading frame 37 ATP-binding cassette, sub-family C (CFTR/MRP), member 5 protease, serine, 23 ankyrin 1, erythrocytic a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 1 chromodomain helicase DNA binding protein 2	W86776 AA428897 W87943 AA442844 W89211 AA442854 W89047 AA442633 W88848 AA403237 W90532 AA42633 W90532 AA42948 AA403237 W90532 AA423948 AA403237 W90532 AA423948 AA005382 AA424561 AA005382 AA4124561 AA005382 AA4124561 AA005382 AA4124561 AA033649 AA418455 R67704 N6489 R67834 N79133 R70685 AF086104 R69925 W03611 R71124 N80617 R76553 N90649
AP-hu01 AP-hu01		4 5 6 7 8 9 10 11 12 13 14 5 6 7 8 9 10 11 12 13 14 5 6 7 8 9 10 11 12 13 14 5 6 7 8 9 10 11 12 13 4 15 7 8 9 10 11 12 13 11 12 13 11 12 11 11 12 13 11 12 11 11 12 11 11 12 13 11 11 11 11 11 11 11 11 11 11 11 11	F4 F5 F6 F7 F8 F9 F10 F11 F12 F13 F14 F15 F16 F17 F18 F19 F20 F21 F23 F24 G12 G3 G4 G5 G6 G7 G8 G9 G11 G12 G13 G14 G15	416803 756965 417258 757718 417361 757875 417435 757968 417801 758534 418033 759207 418040 759881 427750 767174 428412 767295 470958 767302 140197 290340 140536 290517 141815 290549 142833 291426 142944 292212 143887 292699 144797 293017 145284	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) excision repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neuritin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome) Gem-interacting protein hypothetical protein FLJ35036 acetyl-Coenzyme A synthetase 2 (AMP forming)-like chromosome 6 open reading frame 37 ATP-binding cassette, sub-family C (CFTR/MRP), member 5 protease, serine, 23 ankyrin 1, erythrocytic a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 1 chromodomain helicase DNA binding protein 2 nuclear receptor coactivator 2	W86776 AA428897 W87943 AA442848 W89211 AA442854 W89047 AA442633 W38848 AA403237 W90532 AA496054 W90532 AA496054 W90532 AA496054 W90532 AA423948 AA005382 AA418465 R67704 N6489 R67834 N79133 R70685 AF086104 R69925 W03611 R71124 N80617 R76394 N80475 R76553 N90649 R77864
AP-hu01 AP-hu0	בבבבבים מממממממממממ שוויא שלים ממממי שוויא שלים ממממ	4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 9 20 22 23 4 1 2 3 4 5 6 7 8 9 10 11 23 14 15 16 17 18 9 20 12 23 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 18 9 10 11 12 3 14 15 16 17 11 12 23 24 12 3 4 5 6 7 8 9 10 11 12 3 14 15 16 11 12 3 14 15 16 17 18 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 11 12 12 3 12 12 3 12 12 12 3 12 12 12 12 12 12 12 12 12 12 12 12 12	F4 F5 F6 F7 F8 F9 F10 F112 F13 F14 F15 F16 F17 F18 F19 F20 F21 F23 F24 G12 G3 G4 G6 G6 G6 G6 G1112 G13 G112 G13 G112 G13 G15 G17 G1112 G13 G112 G13 G1112 G13 G112 G13 G112 G13 G112 G13 G112 G13 G112 G13 G112 G13 G112 G13 G112 G13 G112 G13 G112 G13 G112 G13 G112 G13 G112 G13 G112 G112	416803 756965 417258 757718 417351 757875 417435 757968 417801 758534 418031 758534 418033 759207 418040 759881 427750 767174 428412 767295 4709588 767302 140197 290340 140536 290517 141815 290549 142383 291426 142944 292212 143887 292699 144797 293017 145284 293914	Homo sapiens transcribed sequences regulator of G-protein signalling 14 fibrinogen-like 2 nuclear transcription factor Y, gamma cell growth regulatory with ring finger domain cyclin-dependent kinase 5, regulatory subunit 1 (p35) hypothetical protein BC013995 transcriptional activator of the c-fos promoter mitochondrial ribosomal protein L27 density-regulated protein carboxypeptidase A1 (pancreatic) exclsion repair cross-complementing rodent repair deficiency, complementation group 4 TGFB-induced factor (TALE family homeobox) fibrillin 1 (Marfan syndrome) spectrin, alpha, erythrocytic 1 (elliptocytosis 2) Purkinje cell protein 4 granzyme K (serine protease, granzyme 3; tryptase II) neurofibromin 2 (bilateral acoustic neuroma) hypothetical protein FLJ10420 KIAA0251 protein neurtitin 1 Homo sapiens transcribed sequences neuropilin 2 integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51) jagged 1 (Alagille syndrome) Gem-interacting protein hypothetical protein FLJ35036 acetyl-Coenzyme A synthetase 2 (AMP forming)-like chromosome 6 open reading frame 37 ATP-binding cassette, sub-family C (CFTR/MRP), member 5 protease, serine, 23 ankyrin 1, erythrocytic a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 1 chromodomain helicase DNA binding protein 2 nuclear receptor coactivator 2 sperm associated antigen 9	W86776 AA428897 W87943 AA442848 W89211 AA442854 W89047 AA442633 W98535 AA496054 W90532 AA496054 W90532 AA496054 W90532 AA496054 W90532 AA423948 AA005382 AA424561 AA005382 AA418463 AA033649 R67834 N59133 R70685 AF086104 R67834 N59133 R70685 AF086104 R67834 N59133 R70685 AF086104 R67553 N90649 R77864 N66027

AD 5.04	~	40	C19	208140	fatty acid binding protein 3, muscle and heart (mammary-derived growth inhibitor)	W00709
AP-0001	Ğ	10	G10	151942	handbatical protein El 112684	H03070
AP-nuui	č	19	619	209601	likely ortholog of mouse phosphorulated adaptor for RNA export	W04927
AP-huu1	G	20	G20	290001	tumor accrecic forter (ligand) superfamily, member 12	D55370
AP-nu01	G	21	G21	104/42	tumor necrosis racion (liganu) superianily, memoer 12	N01550
AP-hu01	G	22	G22	300071	Invited transcription factor i	N91550
AP-hu01	G	23	G23	155/66	Homo sapiens transcribed sequences	R/2149
AP-hu01	G	24	G24	300411	chemokine (C-C motir) ligand 19	W07401
AP-hu01	н	1	H1	471580	BCL2-associated atnanogene 3	AAU34946
AP-hu01	н	2	H2	767746	similar to ICEBERG caspase-1 inhibitor	AA418193
AP-hu01	н	3	H3	471639	chromosome 2 open reading frame 3	AA035031
AP-hu01	н	4	H4	767808	CD47 antigen (Rh-related antigen, integrin-associated signal transducer)	AA418709
AP-hu01	н	5	H5	485294	programmed death ligand 2	AA039608
AP-hu01	н	6	H6	768064	cytochrome P450, family 1, subfamily A, polypeptide 1	AA426553
AP-hu01	н	7	H7	485674	isocitrate dehydrogenase 1 (NADP+), soluble	AA039874
AP-hu01	н	8	H8	768067	ribonuclease, RNase A family, 4	AA426582
AP-hu01	н	9	H9	485778	Sapiens mRNA; cDNA DKFZp564O1016 (from clone DKFZp564O1016)	AA040319
AP-hu01	н	10	H10	768617	diphtheria toxin receptor (heparin-binding epidermal growth factor-like growth	AA430318
					factor)	
AP-hu01	н	11	H11	485857	delta sleep inducing peptide, immunoreactor	AA040063
AP-hu01	н	12	H12	770879	8D6 antigen	AA434501
AP-hu01	н	13	H13	485886	carnitine acetyltransferase	AA040121
AP-hu01	н	14	H14	770989	reticulon 4 receptor	AA428418
AP-hu01	н	15	H15	486233	RAB7, member RAS oncogene family	AA043680
AP-hu01	н	16	H16	771258	CD8 antigen, alpha polypeptide (p32)	AA443649
AP-bu01	н	17	H17	486660	myeloid cell leukemia sequence 1 (BCI 2-related)	AA044316
	н	18	H18	771312	testis zinc finger protein	AA476210
AD 5001	п.	10	L10	496606	oberokine (C.Y.C. matif) recentor 6	AA044440
AP-1001		19	113	772200		AAA29522
AP-hu01		20	H20	113200	Dee 1/(Jee 40) hemeles, subfamily A. member 4	AA420022
AP-nu01	н	21	HZ1	480/2/	Dnaj (risp40) nomolog, subramily A, member 4	AA044432
AP-nu01	н	22	H22	//3399	dystrophin (muscular dystrophy, Duchenne and Becker types)	AA420049
AP-hu01	н	23	H23	487396	Sapiens mRNA; cDNA DKFZp564F053 (from clone DKFZp564F053)	AA046660
AP-hu01	н	24	H24	773494	cathepsin K (pycnodysostosis)	AA428008
AP-hu01	I	1	11	160263	hypothetical protein FLJ22635	H21954
AP-hu01	I	2	12	301740	hypothetical protein LOC139202	W17094
AP-hu01	1	3	13	160946	c-fos induced growth factor (vascular endothelial growth factor D)	H24828
AP-hu01	1	4	14	301851	platelet factor 4 (chemokine (C-X-C motif) ligand 4)	W17158
AP-hu01	1	5	15	161323	autism susceptibility candidate 2	H25394
AP-hu01	1	6	16	302052	adaptor-related protein complex 3, mu 2 subunit	N89763
AP-hu01	1	7	17	162077	pleckstrin homology-like domain, family A, member 1	H26271
AP-hu01	1	8	18	302127	lipocalin 2 (oncogene 24p3)	W38398
AP-hu01	1	9	19	163838	protein kinase C. gamma	H14047
AP-hu01	1	10	110	302188	nebulette	W16723
AP-hu01	1	11	111	174835	microtubule-associated protein 1B	AF085863
AP-hu01	i	12	112	305284	hydroxysteroid (17-beta) dehydrogenase 12	N95053
AP-hu01	i	13	113	176198	ataxin 2-binding protein 1	H41778
AP-hu01	1	14	114	306813	v-ros LIR2 sarcoma virus oncogene homolog 1 (avian)	W24201
		15	115	177241	ankurin reneat and SOCS how containing 3	440730
	1	16	116	307506	Meis1 musicial acotronic viral integration site 1 homolog (mouro)	W21073
AD hu01	1	10	110	179257		4421073
AP-1001	-	40	117	1/025/	U nhaanhandasa kinaan annon 1 (musala)	H40030
AP-huu I	-	18	118	306766	phosphorylase kinase, gamma T (muscle)	N93274
AP-nu01		19	119	191708	protein tyrosine phosphatase, non-receptor type 3	H40246
AP-hu01	1	20	120	310070	DEAD/H (Asp-Giu-Ala-Asp/His) box polypeptide 16	N98554
AP-hu01	1	21	121	196005	aquaporin 3	R91904
AP-hu01	1	22	122	322939	forkhead box D1	W45110
AP-hu01	I	23	123	196387	ATP-binding cassette, sub-family C (CFTR/MRP), member 2	R91503
AP-hu01	1	24	124	323078	heme oxygenase (decycling) 1	W42558
AP-hu01	J	1	J1	487836	nuclear phosphoprotein similar to S. cerevisiae PWP1	AA045396
AP-hu01	J	2	J2	773568	POU domain, class 4, transcription factor 1	AA428894
AP-hu01	J	3	J3	487965	ubiquitin-conjugating enzyme E2A (RAD6 homolog)	AA045751
AP-hu01	J	4	J4	781738	GATA binding protein 4	AA431631
AP-hu01	J	5	J5	488033	DnaJ (Hsp40) homolog, subfamily B, member 9	AA045793
AP-hu01	J	6	JG	782288	glutathione S-transferase A2	AA432244
AP-hu01	J	7	J7	488629	hypothetical protein FLJ10154	AA045908
AP-hu01	J	8	.18	782758	CD209 antigen-like	AA448002
AP-hu01	Ĵ	9	.19	488842	osteoblast specific factor 2 (fasciclin Hike)	AA046245
AP-hu01	,Î	10	.110	784283		AA447502
	ĭ	11	111	480042	tumor pecrosis factor recentor superfamily, member 8	AA047162
	ĩ	12	112	794595	but macholineeterase	AA442200
		12	J12 140	480212	Duty yici ioni iosiorase	AA443299
AP-11001		13	J13	409212	DCV0 like enter the subusit of DNA palareness anter (used)	AA045659
AP-nu01	3	14	J14	/6495/	REV3-like, catalytic subunit of DNA polymerase zeta (yeast)	AA448067
AP-hu01	1	15	J15	490004	cytokine receptor-like factor 1	AA121532
AP-nu01	J	16	J16	786125	TAP12 RNA polymerase II, TATA box binding protein (TBP)-associated factor,	AA447903
AD 5001		47	14.7	400000	20KDa viazlase hme MiCD/ interaction site family, member 50	
AP-NUU1	J	17	J1/	490023	wingless-type MM I v integration site ramily, member 58	AA114966
AP-0001	J	18	J18	189049	Septens full length insert cUNA clone YW26E10	AA453077
AP-nu01	J	19	J19	490111	U-o-methylguanine-UNA methyltransferase	AA137218
AP-hu01	J	20	J20	795562	SMC4 structural maintenance of chromosomes 4-like 1 (yeast)	AA459795
AP-hu01	J	-21	J21	490642	cullin 3	AA115733
AP-hu01	J	22	J22	795871	muscleblind-like 3 (Drosophila)	AA460747
AP-hu01	J	23	J23	501994	interleukin 1 receptor-like 1	AA128153
AP-hu01	J	24	J24	796199	hypothetical protein FLJ20069	AA461119
AP-hu01	к	1	K1	197105	KIAA0265 protein	R93904
AP-hu01	κ	2	K2	323151	nuclear receptor subfamily 4, group A, member 3	W42610
AP-hu01	к	3	КЗ	201268	ephrin-A5	R99405
AP-hu01	к	4	K4	323181	fibroblast activation protein, alpha	W45237
AP-hu01	к	5	K5	202112	nuclear factor (erythroid-derived 2), 45kDa	H48815
AP-hu01	ĸ	6	K6	323238	chemokine (C-X-C motif) ligand 1 (melanoma growth stimulating activity alpha)	W42812
AP-hu01	ĸ	7	K7	202603	BCL2-associated athanogene 4	H53280
	••	•				100203

			-		001051	and a second fraction is the second	MACEOO
AP-hu	01	ĸ	8	K8	324054	herve growth factor, beta polypeptide	VV 40022
AP-bu	01	к	9	K9	203514	hypothetical protein FLJ00225	H55966
AD 6	~	2	10	KAO	205265	HIV 1 row binding protein 2	W62273
AP-nu	01	ĸ	10	KIU	320300	HIV-1 fev binding protein 2	W52273
AP-hu	01	ĸ	11	K11	205898	p8 protein (candidate of metastasis 1)	H58465
AP_hu	01	ĸ	12	K12	327060	dutathione S-transferase M1	W25634
			12		021000		000070
AP-hu	01	ĸ	13	K13	206779	Homo sapiens transcribed sequences	R98072
AP-hu	01	ĸ	14	K14	340588	uridine monophosphate synthetase (orotate phosphoribosyl transferase and	W56655
						orotidine-5'-decarboxylase)	
40 h.	~	~	45	MAR	000047		D09700
AP-nu	01	ĸ	15	K15	206917	adenyiate kinase 3	R96700
AP-hu	01	ĸ	16	K16	340644	integrin, beta 8	W56754
	04	v	17	147	207446	and the membrane protein hand 4.2	H59950
AP-nu	01	r.	17	K17	207410	eryurocyte memorane protein band 4.2	100009
AP-hu	01	ĸ	18	K18	341295	inducible T-cell co-stimulator	W58028
AD-bui	01	×	10	K10	207665	Homo series transcribed sequence with weak similarity to protein	H62267
AFHQ		ĸ	19	K19	20/003	rolling applets tarbondou soquence wat work shintarty to protein	1102207
						rer:NP_000434.1 (H.sapiens) hypothetical protein FL020037 [Homo sapiens]	
AP-hu	01	ĸ	20	K20	341763	caspase 5, apoptosis-related cystelle protease	W60764
AP-hu	01	к	21	K21	207838	hypothetical protein LOC129607	H60298
AD hus		2	20	100	340054	susing surjectide bioding protein (Constein), sinks inhibiting activity solvestide	W61100
AP-nu	01	n i	22	N22	342234	guarine nucleolide binding protein (G protein), alpha minibiting activity polypeptide	4401133
						1	
AP-hu	01	ĸ	23	K23	209731	pim-1 oncogene	H52219
AD but	01	ĸ	24	K24	342557	complement C1r-like proteinase	W68477
AF-10		n.	24	1.2.4	342337		******
AP-hu	01	L	1	L1	502055	aryisulfatase B	AA128826
AP-hu	01	L	2	12	796702	FKBP-associated protein	AA443131
10 10	~ .	-	-		500004		A 4 5 4 700
AP-nu	01	L	3	L3	503281		AA151736
AP-hu	01	L	4	L4	809464	fibroblast growth factor receptor 2 (bacteria-expressed kinase, keratinocyte growth	AA456160
						factor receptor craniofacial dysostosis 1. Crouzon syndrome. Pfeiffer syndrome	
	• •		-			Jackson-weiss syncrome)	
AP-hu	U1	L	5	L5	503313	interreron-induced protein with tetratricopeptide repeats 2	AA130252
AP-hu	01	L	6	L6	810391	hvaluronoolucosaminidase 1	AA464196
AD			÷		500040		
AP-nu	01	L	1	L/	503343	collagen, type IV, alpha 1	AA130288
AP-hu	01	L	8	L8	811942	general transcription factor IIH, polypeptide 1, 62kDa	AA455004
AD hus	0 4	ī	õ	10	EDAEDE	SEC24 related game family member (2/8 consulting)	AA450040
AP-00		-	э	La.	004030	SEC24 related gene lamily, member C (S. Cerevisiae)	AA 100049
AP-hu	01	L	10	L10	814287	X-ray repair complementing defective repair in Chinese hamster cells 3	AA459237
AD bu	01	1	11	1 1 1	505053	lentin recentor gene-related protein	44140805
APHIC				E11	303033	iepuiriecepioi generielateu protein	AA143000
AP-hu	01	L	12	L12	814578	matrix metalloproteinase 19	AA480939
AP-hu	01	1	13	1 13	665121	aldebyde dehydrogenase 3 family, member B1	AA194963
					045750		
AP-hu	01	L	14	L14	815/58	epidermal growth factor receptor pathway substrate 15	AA485200
AP-hu	01	1	15	L15	665424	histidyl-tRNA synthetase-like	AA195056
AD 6	~		40	140	004000		44400050
AP-nu	01	L	10	LIO	824803	0	AA488909
AP-hu	01	L	17	L17	666445	BRF1 homolog, subunit of RNA polymerase III transcription initiation factor IIIB (S.	AA232937
						cerevisiae)	
	01		10	110	925297	tumor percesis factor (ligand) superfamily member 11	A A E O 4 4 E O
AFHIU	01	-	10	LIO	020207	tunior necrosis factor (ligand) superfamily, member 11	MA004400
AP-hu	01	L	19	L19	667290	tyrosinase-related protein 1	AA227492
			20	1 20	825352	histone deacetvlase 9	AA504568
AP-hu	01		211				
AP-hu	01	L.	20		007407		
AP-hu AP-hu	01 01	L	21	L21	667407	sperm associated antigen 9	AA227958
AP-hu AP-hu AP-hu	01 01 01	L L	20 21 22	L21 L22	667407 878499	sperm associated antigen 9 hemoglobin, beta	AA227958 AA775838
AP-hu AP-hu AP-hu	01 01 01		20 21 22	L21 L22	667407 878499	sperm associated antigen 9 hemoglobin, beta	AA227958 AA775838
AP-hu AP-hu AP-hu AP-hu	01 01 01 01	L L L	20 21 22 23	L21 L22 L23	667407 878499 667440	spern associated antigen 9 hemoglobin, beta B aggressive lymphoma gene	AA227958 AA775838 AA228088
AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01	L L L	21 22 23 24	L21 L22 L23 L24	667407 878499 667440 1030527	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB	AA227958 AA775838 AA228088 AA555198
AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01		21 22 23 24	L21 L22 L23 L24	667407 878499 667440 1030527 210587	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB same domain immunoclobulin domain (In), transmembrane domain (TMI) and short	AA227958 AA775838 AA228088 AA5555198
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01	L L L M	20 21 22 23 24 1	L21 L22 L23 L24 M1	667407 878499 667440 1030527 210587	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short	AA227958 AA775838 AA228088 AA555198 H65936
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01	L L L L M	20 21 22 23 24 1	L21 L22 L23 L24 M1	667407 878499 667440 1030527 210587	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D	AA227958 AA775838 AA228088 AA555198 H65936
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01	L L L M M	20 21 22 23 24 1 2	L21 L22 L23 L24 M1	667407 878499 667440 1030527 210587 342614	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein)	AA227958 AA775838 AA228088 AA555198 H65936 W68761
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01	L L L M M	20 21 22 23 24 1 2	L21 L22 L23 L24 M1 M2 M3	667407 878499 667440 1030527 210587 342614 210658	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01	LLLMM	20 21 22 23 24 1 2 3	L21 L22 L23 L24 M1 M2 M3	667407 878499 667440 1030527 210587 342614 210658	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cytsteine and glycine-rich protein 3 (cardiac LIM protein) 0	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01	LLLMMMM	20 21 22 23 24 1 2 3 4	L21 L22 L23 L24 M1 M2 M3 M4	667407 878499 667440 1030527 210587 342614 210658 342633	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01	LLMMMM	20 21 22 23 24 1 2 3 4 5	L21 L22 L23 L24 M1 M2 M3 M4 M5	667407 878499 667440 1030527 210587 342614 210658 342633 212088	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01	LLLMMMM	20 21 22 23 24 1 2 3 4 5	L21 L22 L23 L24 M1 M2 M3 M4 M5	667407 878499 667440 1030527 210587 342614 210658 342633 212088 242144	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 case claticle projection (chemical cardiac LIM)	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W427202
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01	L L L M M M M M	20 21 22 23 24 1 2 3 4 5 6	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7)	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01	L L L M M M M M M	20 21 22 23 24 1 2 3 4 5 6 7	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor. ciade D (heparin cofactor), member 1	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68884 H68960 W67392 H72143
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0		20 21 22 23 24 1 2 3 4 5 6 7	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M6 M7 M8	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 213414	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 perce gravit factor recentor (TMEB superfinitum member 16)	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	L L L M M M M M M M M	20 21 22 23 24 1 2 3 4 5 6 7 8	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16)	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	L L L L L M M M M M M M M M M M M	20 21 22 23 24 1 2 3 4 5 6 7 8 9	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M6 M7 M8 M9	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cystelne) proteinase inhibitor, ciade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68884 H68960 W67392 H72143 W67446 H70847
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01		20 21 22 23 24 1 2 3 4 5 6 7 8 9 10	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 34388	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, ciade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-denendent torvien kinase 1 sipha	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01		20 21 22 23 24 1 2 3 4 5 6 7 8 9 10	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343898	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68884 H68960 W67392 H72143 W67446 H70847 W69176
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	LLLLM MMMMMMMMMM	21 21 22 23 24 1 2 3 4 5 6 7 8 9 10 11	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 343698 214427	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, ciade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0		21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M5 M6 M6 M6 M7 M8 M9 M10 M10 M12	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343898 214427 344107	spern associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	LLLLM MMMMMMMMMMMMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M6 M7 M8 M9 M10 M11 M12	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343698 214427 344107	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitymsin), member 1	AA227958 AA775838 AA228088 AA228088 AA555198 H65936 W68761 H66936 W68761 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	LLLLM MMMMMMMMMM ;	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343311 214144 343898 214427 344107	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	LLLLM MMMMMMMMMM M	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M6 M7 M8 M9 M10 M11 M12 M13	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343698 214427 344107 214469	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, ciade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, ciade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68884 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	LLLLM MMMMMMMMM MM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 14	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 3433698 214427 344107 214469 344384	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784 H73833 W73447
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	LLLLM MMMMMMMMMM MMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 14 5	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 5	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343698 214427 344107 214469 214469 214426	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, ciade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, ciade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echipodem microthukia easociated protein like 1	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68884 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81649
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	L L L L L M M M M M M M M M M M M M M M	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 343698 214427 344107 214469 344384 219766	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase in apha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618
AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu AP-hu	01 01 01 01 01 01 01 01 01 001 001 001	L L L L L M M M M M M M M M M M M M M M	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343311 214144 343698 214427 344107 214469 344384 219766 344443	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68884 H68960 W67392 H72143 W67346 H77955 W73784 H73833 W73784 H73833 W73447 H81618
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	L L L L L L M M M M M M M M M M M M M M	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 13 14 15 6 7 8 9	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M17	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343311 213414 343311 213414 343311 214144 343698 214427 344107 214469 344384 219766 344443 220826	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 001 001 001	L L L L L M M M M M M M M M M M M M M M	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 77 9	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M17	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343311 214144 343598 214427 344107 214469 344384 219766 344443 220826	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	L L L L L L M M M M M M M M M M M M M M	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 343698 214427 344107 214469 344384 219766 344443 220826 344889	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 00 00 00 0	L L L L L L M M M M M M M M M M M M M M	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 13 4 15 16 7 18 19	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343311 214144 343898 214427 344107 214469 344384 219766 344443 220826 344889 221191	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 00 00 00 0	L L L L L L M M M M M M M M M M M M M M	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 13 14 15 6 17 18 9 10	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M14 M15 M17 M18 M19 M19	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 343698 214427 344107 214469 344384 219766 344443 220826 344889 221191	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0	AA227958 AA775838 AA228088 AA5551936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 001 001 0	L L L L M MMMMMMMMMMMMMMMMMMMMMMMMMMMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M17 M18 M19 M20	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343311 214144 343698 214427 344107 214469 344384 219766 344443 220826 344889 221191 344953	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086339
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 00	L _ L L M _ M M M M M M M M M M M M M M	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 13 14 5 16 7 18 9 20 21	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M14 M15 M17 M18 M17 M18 M17 M18 M17 M18 M17 M12 M12 M12 M12 M12 M12 M1 M1 M2 M2 M1 M1 M2 M2 M1 M2 M2 M1 M2 M2 M2 M2 M2 M2 M2 M2 M2 M2 M2 M2 M2	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 213414 343311 214144 343698 214427 344107 214469 344384 219766 344433 220826 344889 221191 344953 3221527	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, colled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 1 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086339 H92126
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 001 001 001	L L L L M MMMMMMMMMMMMMMMMMMMMMMMMMMMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 7 18 19 20 21	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21	667407 878499 667440 1030527 210587 342614 210658 343111 213414 343311 213414 343311 214144 343311 214144 343311 214144 3434107 214469 344484 219766 344443 220826 344889 221191 344953 221527	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref NP 115694.1 (H sapiens) humothatical protein MGC13045 Homo senional	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W73091 H91887 AF086339 H92126
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 001 001 001	L L L L L L M M M M M M M M M M M M M M	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 2 3 4 5 6 7 8 9 10 11 2 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 10 10 10 10 10 10 10 10 10 10 10 10 10	L21 L22 L23 L24 M1 M2 M3 M4 M5 M5 M5 M6 M17 M18 M14 M15 M14 M15 M14 M15 M14 M17 M18 M17 M18 M19 M21	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 3433311 214144 3433311 214144 343898 214427 344107 214469 344384 219766 344443 220826 344489 221191 344953 221527 345034	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, colled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 1 1/ymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref.NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens]	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086339 H92126
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 00 00	L L L L L L M M M M M M M M M M M M M M	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 7 18 19 20 21 22 23 24 2 4 5 6 7 8 9 10 21 22 23 24 24 22 23 24 24 22 23 24 24 22 23 24 24 22 23 24 24 22 23 24 24 22 23 24 24 24 20 24 24 20 24 24 20 24 20 24 20 24 20 24 20 20 24 20 20 20 20 20 20 20 20 20 20 20 20 20	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M14 M15 M16 M17 M18 M20 M21 M22	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 343311 214144 3434107 214469 344427 344107 214469 344443 219766 344443 220826 344889 221191 344953 221527 345034	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784 H73833 W73047 H81618 W73191 H95623 W76009 H91887 AF086339 H92126 AF086340
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	LLLLM MMMMMMMMMMMMMMMMMMMMMMMMMMMMMMMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 13 14 5 16 7 18 9 20 21 22 23 24 2 23 24 2 23 24 2 23 24 2 23 24 2 23 24 22 23 24 22 23 24 24 22 23 24 24 22 23 24 24 22 23 24 24 22 23 24 24 22 23 24 24 25 26 26 26 26 26 26 26 26 26 26 26 26 26	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M16 M17 M18 M19 M20 M21 M22 M23	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343311 214144 343698 214427 344107 214469 344384 219766 344433 220826 344489 221191 344953 221527 345034 222454	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 1 1/ymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H84077
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 00 00 00 00 0	L L L L L L M M M M M M M M M M M M M M	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 7 18 19 20 21 22 23 24 2 4 5 6 7 8 9 10 21 22 23 24 24 22 23 24 24 22 23 24 24 22 23 24 24 22 23 24 24 22 23 24 24 22 23 24 24 24 20 24 24 20 24 24 20 24 20 24 20 24 20 24 20 20 24 20 20 20 20 20 20 20 20 20 20 20 20 20	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21 M22 M23 M24	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 3433698 214427 344107 214469 344384 219766 344433 220826 344889 221191 344953 221527 345034 222454 345034	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (cherokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 0 0 1/mphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein reft.NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apontoris antagonizing transcribed factor r	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73047 H81618 W73191 H95623 W76009 H91887 AF086340 H84077 W72325
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 01 01 01 0	LLLLM MMMMMMMMMMMMMMMMMMMMMMMMMMMMMMMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 9 20 21 22 23 24 24 2 3 4 5 6 7 8 9 10 11 22 23 24 10 22 23 24 24 24 22 23 24 24 23 24 24 23 24 24 24 23 24 24 24 24 24 24 24 24 24 24 24 24 24	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M16 M17 M18 M18 M19 M20 M21 M22 M23 M24	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343311 214144 343311 214144 343898 214427 344107 214469 344384 219766 344433 220826 344889 221191 344953 221527 345034 222454 345214	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 1 1/ymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W737847 H81618 W73191 H95623 H92126 AF086340 H84077 W72325
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 00 01 00 01 00 00	LLLLM MMMMMMMMMMMMMMMMMMMMMMMMMMMMMMMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 22 23 24 1 22 23 24 1 22 23 24 1 22 23 24 1 22 23 24 24 23 24 24 23 24 24 24 24 24 24 24 24 24 24 24 24 24	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21 M22 M23 M24 N1	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 3433698 214427 344107 214469 3444384 219766 344443 220826 344889 221191 344953 221527 345034 222454 345214 667596	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, colied-coli containing protein kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 0 0 0 1/mphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein refr.NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin)	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W730447 H81618 W73191 H95623 W76009 H91887 AF086339 H92126 AF086340 H84077 W72325 AA228129
AP-hu AP-hu	01 01 01 01 01 00 01 00 00 00 00 00 00 0	LLLLM MMMMMMMMMMMMMMMMMMMMMMMMMMMMMMMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 2 3 4 5 6 7 8 9 10 11 2 2 2 3 4 5 6 7 7 8 9 10 11 12 2 2 3 4 5 6 7 7 8 9 10 11 12 2 2 3 4 5 6 7 7 8 9 10 11 11 12 2 3 4 5 6 7 7 8 9 10 11 11 12 2 10 11 11 12 10 11 11 11 11 11 11 11 11 11 11 11 11	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M17 M18 M17 M18 M19 M20 M21 M22 M23 M24 N2	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343311 214144 343311 214144 343898 214427 344107 214469 344384 219766 344433 220826 344489 221191 344953 221527 345034 222454 345214 667596 1046762	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin)	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W73784 H73833 W73447 H81618 W73191 H95623 H92126 AF086340 H84077 W72325 AA228129 Aa644350
AP-hu AP-hu	01 01 01 001 001 001 001 001 001 001 00	LLLLM MMMMMMMMMMMMMMMMMMMMMMMMMMMMMMMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 7 18 19 20 21 22 23 24 1 22 23 24 1 2 2 3 4 5 6 7 8 9 10 11 22 23 24 1 22 23 24 1 2 2 3 24 1 2 2 3 24 1 2 2 3 2 4 5 6 7 8 9 10 11 12 2 3 2 4 5 6 7 8 9 10 11 12 2 3 2 4 5 6 7 8 9 10 11 12 2 3 2 4 5 6 7 8 9 10 11 12 2 3 2 4 5 6 7 8 9 10 11 12 2 3 12 2 10 11 12 2 11 12 2 11 12 12 12 12 12 12 12	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M14 M15 M14 M15 M14 M15 M14 M15 M14 M12 M20 M21 M21 M22 M23 M24 M1 M22 M23 M24 M23 M24 M2 M22 M23 M24 M2 M2 M23 M24 M3 M3 M4 M3 M4 M5 M3 M4 M5 M6 M1 M3 M4 M5 M6 M1 M3 M4 M5 M6 M1 M3 M4 M5 M6 M1 M3 M4 M5 M6 M1 M3 M4 M5 M6 M1 M3 M4 M5 M6 M1 M3 M4 M5 M6 M1 M3 M4 M5 M6 M1 M1 M3 M4 M5 M6 M1 M1 M2 M3 M4 M5 M6 M1 M1 M2 M1 M1 M2 M1 M2 M1 M3 M4 M2 M1 M1 M1 M1 M3 M4 M5 M6 M1 M1 M1 M1 M2 M1 M1 M2 M1 M1 M2 M1 M1 M1 M1 M2 M10 M11 M12 M12 M12 M14 M12 M14 M12 M14 M12 M14 M14 M15 M14 M14 M15 M14 M14 M15 M14 M14 M15 M14 M15 M14 M15 M14 M15 M14 M15 M14 M15 M14 M15 M14 M15 M14 M15 M14 M15 M16 M17 M12 M12 M12 M12 M12 M12 M12 M14 M15 M14 M15 M14 M15 M14 M15 M12 M2 M14 M15 M2 M2 M14 M15 M2 M14 M15 M2 M2 M14 M2 M2 M14 M2 M2 M2 M2 M2 M2 M2 M2 M2 M2 M2 M2 M2	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 3433698 214427 344107 214469 344384 219766 344489 221191 344953 220826 344889 221191 344953 221527 345034 222454 345214 667596 1046762	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, colied-coli containing protein kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 0 0 0 0 1/mphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H84077 W72325 AA228129 AA644350
AP-hu AP-hu	01 01 01 001 001 001 001 001 001 001 00	LLLLM MMMMMMMMMMMMMMMMMMMMMMMMMMMMMMMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 7 8 9 10 11 12 2 3 4 5 6 7 7 8 9 10 11 12 2 3 4 5 6 7 7 8 9 10 11 12 2 3 4 5 6 7 7 8 9 10 11 11 12 2 10 11 11 12 2 10 11 11 12 2 10 11 11 12 11 11 12 11 11 11 11 11 11 11	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M17 M18 M17 M18 M20 M21 M21 M22 M23 M24 N1 N2 N3	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343311 214144 343311 214144 343598 214427 344107 214469 344384 219766 34443 220826 344889 221191 344953 221527 345034 222454 345214 667596 1046762 668487	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor,	AA227958 AA775838 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W739447 H81618 W73191 H95623 H72126 AF086340 H84077 W72325 AA544350 AA243818
AP-hu AP-hu	01 01 01 01 01 01 01 01 01 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 01 00 00	LLLLM MMMMMMMMMMMMMMMMMMMMMMMMMMMMMMMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 13 14 15 6 7 8 9 10 11 12 23 24 1 22 23 24 1 22 23 24 1 22 23 24 1 22 23 24 1 22 23 24 1 22 23 24 24 23 24 24 23 24 24 23 24 24 24 24 24 24 24 24 24 24 24 24 24	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M14 M15 M14 M15 M14 M15 M14 M15 M20 M21 M21 M21 M21 M23 M24 N1 N2 N3	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 213414 343311 214144 343698 214427 344107 214469 344384 219766 344433 220826 344889 221191 344953 221527 345034 222454 345214 667596 1046762 668487	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32KDa	AA227958 AA775838 AA228088 AA5251936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H84077 W72325 AA228129 AA644350 AA243818
AP-hu AP-hu	01 01 01 001 001 001 001 001 001 001 00	LLLLM MMMMMMMMMMMMMMMMMMMMMMMMMMMMMMMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 3 24 1 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 13 12 12 10 10 11 12 2 10 11 12 2 10 11 11 12 12 10 11 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 11	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M17 M18 M17 M18 M20 M21 M21 M22 M23 M24 N1 N2 N3 N4	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 343311 214144 343698 214427 344107 214469 344384 219766 344443 220826 344889 221191 344953 221527 345034 222454 345214 667596 1046762 668487 1075967	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 ymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32KDa	AA227958 AA775838 AA725838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W739447 H81618 W73191 H95623 W76009 H91887 AF086339 H92126 AF086340 H84077 W72325 AA243818 AA243818 AI821703
AP-hu AP-hu	01 01 001 001 001 001 001 001 001 001 0	LLLLM MAMMAMAMAMA MAMMAMAMA MAMAZZZZ ZZ	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 13 14 15 6 7 8 9 10 11 12 23 24 1 2 2 3 4 5 6 7 8 9 10 11 22 23 24 1 22 23 24 1 2 2 3 24 1 2 2 3 2 4 5 6 7 8 9 10 11 2 2 3 2 4 5 6 7 8 9 10 11 12 2 3 2 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 2 4 5 6 7 8 9 10 11 12 2 1 2 1 2 1 2 1 2 1 12 2 1 12 2 1 11 1	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M14 M15 M14 M15 M14 M15 M14 M15 M20 M21 M21 M21 M21 M21 M21 M21 M21 M21 M21	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 213414 343311 214144 343698 214427 344107 214469 344384 219766 344443 220826 344889 221191 344953 221527 345034 222454 345214 667596 1046762 668487 1075967	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32KDa BCL2 binding component 3 transloading antigen antigen antigen set antigen antigen set antigen	AA227958 AA775838 AA228088 AA5551936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H84077 W72325 AA228129 AA644350 AA243818 AI821703
AP-hu AP-hu	01 01 001 001 001 001 001 001 001 001 0	LLLLM MMMMMMMMMMMMMMMMMMMMMMMMMMMMMMMM	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 2 3 24 1 2 2 3 24 1 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 2 4 11 2 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 15 10 11 12 2 3 4 5 10 11 12 2 3 4 5 10 11 12 2 3 4 5 10 10 11 2 2 3 4 5 10 11 12 2 3 2 4 11 2 2 3 2 4 5 10 11 12 2 3 2 4 11 12 2 3 2 4 11 12 2 3 2 4 1 12 2 3 2 4 11 12 2 3 2 4 1 1 2 2 3 2 4 1 2 2 3 2 4 1 2 2 3 2 4 1 2 3 2 4 1 1 2 2 3 2 4 1 2 2 3 2 4 1 2 2 3 2 4 1 2 2 3 2 4 1 2 2 3 2 4 1 2 3 2 4 5 5 2 3 2 4 1 2 2 3 2 4 4 5 5 2 2 3 2 4 5 5 2 2 2 2 2 4 5 5 2 2 2 2 2 2 4 4 5 5 2 2 2 2	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21 M20 M21 M22 M23 M24 N1 N2 N3 N4 N5	667407 878499 667440 1030527 210587 342614 210658 343111 213414 343311 213414 343311 214144 343311 214144 343311 214144 343348 214427 344107 214469 344384 219766 344443 220826 344889 221191 344953 221527 345034 222454 345034 222454 345214 667596 1046762 668487 1075967 669163	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 Iymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32kDa BCL2 binding component 3 translocation protein 1	AA227958 AA775838 AA725838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W739447 H81618 W73191 H95623 W7309 H91887 AF086339 H92126 AF086340 H84077 W72325 AA228129 AA644350 AA2443818 AI821703 AA234058
AP-hu AP-hu	01 01 001 001 001 001 001 001 001 001 0	LLLLX MAMAAAAAAAA MAAAAAAAAAAAAAAAAAAAAA	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 13 14 15 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 22 23 24 1 22 3 4 5 6 7 8 9 10 11 22 23 24 1 22 3 4 5 6 7 8 9 10 11 22 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 12 2 12 2 12 2 12 2 12 2 1	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M14 M15 M14 M15 M14 M15 M20 M21 M21 M22 M23 M24 N1 N2 N3 M24 N1 N2 N3 M4 N5 M2 M24 N1 M2 M2 M24 N1 M2 M2 M24 M1 M2 M2 M2 M2 M2 M2 M3 M4 M5 M6 M2 M3 M4 M5 M6 M3 M4 M5 M6 M6 M3 M4 M5 M6 M6 M7 M7 M7 M8 M9 M1 M7 M7 M8 M9 M1 M7 M7 M8 M9 M1 M7 M7 M8 M9 M1 M1 M1 M7 M7 M8 M9 M10 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 213414 343311 214144 343698 214427 344107 214469 344384 219766 344443 220826 344889 221191 344953 221527 345034 22254 345214 667596 1046762 668487 1075967 669163 1146395	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32KDa BCL2 binding component 3 translocation protein 1 SH2 domain protein 2A	AA227958 AA775838 AA228088 AA525198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H84077 W72325 AA228129 AA644350 AA243818 AI821703 AA24058 AA613070
АР-ћи АР-	01 01 001 001 001 001 001 001 001 001 0	LILLS SASANANANAN SANANAN SANANAN SANANAN	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 10 11 12 2 3 4 5 7 8 9 10 11 12 2 3 4 5 8 9 10 11 12 2 3 4 5 7 8 9 10 11 12 2 3 4 5 8 9 10 11 12 2 3 4 5 15 7 8 9 10 11 12 2 3 4 5 1 8 9 10 11 12 2 3 2 4 5 1 8 9 10 11 12 2 3 2 4 1 1 2 3 2 4 5 1 12 2 10 11 12 2 3 2 4 1 2 3 2 4 1 2 3 2 4 1 2 3 2 1 2 3 2 4 1 2 3 2 4 1 2 3 2 4 5 5 6 7 7 8 9 10 11 12 2 3 2 4 1 2 3 2 4 5 5 6 7 1 2 3 2 4 5 5 6 7 7 8 9 10 1 1 2 1 2 2 3 2 4 1 2 3 2 4 5 5 6 7 1 2 2 3 2 4 5 5 7 2 2 2 2 2 4 5 5 7 2 2 2 2 2 2 4 5 5 7 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M14 M15 M16 M17 M18 M20 M21 M23 M23 M24 N1 N2 N3 M24 N1 N2 N3 M2 N3 M2 N2 N3 M2 N3 M2 M2 N3 M2 M2 M2 M2 M2 M2 M2 M2 M2 M2 M2 M2 M2	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 343311 214144 343311 214144 3434107 214469 344489 214427 344107 214469 344384 219766 34489 221191 344953 221527 345034 222454 345214 667596 1046762 668487 1075967 669163 1146395	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 0 1 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sejiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32kDa BCL2 binding component 3 translocation protein 1 SH2 domain protein 2A zinc finger protein 2A	AA227958 AA775838 AA775838 AA228088 AA555198 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784 H73833 W73047 H73833 W73047 H73833 W73047 H31618 W73191 H95623 W76009 H91887 AF086339 H92126 AF086340 H84077 W72325 AA228129 AA644350 AA243818 AI821703 AA234058 AA613070
AP-hu AP-hu	01 01 001 001 001 001 001 001 001 001 0	LILLA MAMAAAAAAAA MAAAAAAAAAAAAAAAAAAAAA	21 22 23 24 1 2 3 4 5 6 7 8 9 101 12 13 14 15 6 7 8 9 101 12 13 14 15 6 7 2 2 2 4 1 2 3 4 5 6 7 8 9 101 12 2 3 4 5 6 7 8 9 101 12 2 3 4 5 6 7 8 9 101 12 2 3 4 5 6 7 8 9 101 11 2 2 3 4 5 6 7 8 9 101 11 2 2 3 4 5 6 7 8 9 101 11 2 2 3 4 5 6 7 8 9 101 11 2 2 3 4 5 6 7 8 9 101 11 2 2 3 4 5 6 7 8 9 101 11 2 2 3 4 5 6 7 8 9 101 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 7 8 9 10 11 12 2 3 4 5 7 8 9 10 11 12 7 13 10 11 12 13 14 15 16 7 10 11 12 13 14 15 16 17 11 12 11 12 11 12 11 12 11 12 111 2 1 12 11 12 11 11	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M17 M18 M19 M20 M21 M22 M23 M24 N1 N2 N3 M24 N1 N2 N3 M24 N1 N2 N3 M24 N1 M2 M3 M4 M5 M1 M3 M4 M1 M3 M4 M5 M1 M3 M4 M5 M1 M3 M4 M5 M3 M4 M5 M3 M4 M5 M3 M4 M5 M3 M4 M5 M3 M4 M5 M3 M4 M5 M3 M4 M5 M3 M4 M5 M5 M6 M1 M1 M3 M4 M5 M5 M6 M1 M1 M3 M4 M5 M6 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 213414 343311 214144 343698 214427 344107 214469 344384 219766 344443 220826 344889 221191 344953 221527 345034 222454 345214 667596 1046762 668487 1075967 669163 1146395 669194	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, colled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KKAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32kDa BCL2 binding component 3 translocation protein 1 SH2 domain protein 1 SH2 domain protein 2A zinc finger protein 451	AA227958 AA775838 AA228088 AA525198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H84077 W72325 AA228129 AA644350 AA243818 AI821703 AA24058 AA613070 AA236470
AP-hu AP-hu	01 01 01 001 001 001 001 001 001 001 00	LILLS SASASASASAS SASASASAS SASASASAS SA	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 8 9 10 11 12 2 3 4 5 7 8 9 10 11 12 2 3 4 5 8 9 10 11 12 2 3 2 3 4 5 8 9 10 11 12 2 3 2 2 1 2 3 2 2 1 2 3 2 2 1 1 2 3 2 3	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M14 M15 M16 M17 M18 M12 M20 M21 M20 M21 M23 M23 M24 N1 N2 N3 M2 N3 M2 N3 M2 N3 M2 N3 M2 N3 M2 M3 M4 M3 M4 M5 M6 M7 M1 M3 M4 M5 M6 M7 M7 M8 M6 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M7 M7 M8 M9 M10 M1 M1 M1 M7 M8 M7 M10 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M7 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 343311 214144 3433698 214427 344107 214469 344443 219766 344443 220826 344849 221191 344953 221527 345034 222454 345214 667596 1046762 668487 1075967 669163 1146395	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 0 1 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein refr.NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32kDa BCL2 binding component 3 translocation protein 1 SH2 domain protein 2A zinc finger protein 451 homeo box A9	AA227958 AA775838 AA775838 AA228088 AA555198 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W69176 H77955 W73784 H73833 W73047 H81618 W73191 H95623 W76009 H91887 AF086340 H84077 W72325 AA228129 AA644350 AA243818 AI821703 AA234058 AA613070 AA236470 AA236470 AA236470 AA267519
AP-hu AP-hu	01 01 001 001 001 001 001 001 001 001 0	LILLS MANAXANANA MANANANANANANANANANANANANANANANANA	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 23 24 1 2 3 4 5 6 7 8 9 10 11 12 23 24 1 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 11 2 2 3 4 12 2 3 4 12 2 11 2 2 3 4 12 2 3 4 12 2 12 2	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21 M22 M23 M24 N1 N2 N3 M24 N1 N2 N3 M24 N1 N2 M3 M2 M1 M3 M1 M1 M3 M4 M5 M1 M1 M3 M4 M5 M1 M3 M4 M5 M1 M3 M4 M5 M6 N2 M3 M1 M1 M3 M4 M3 M4 M5 M3 M4 M5 M5 M6 M1 M1 M3 M4 M5 M3 M4 M5 M5 M6 M1 M1 M1 M3 M4 M5 M5 M6 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 213414 343311 213414 343311 214144 343698 214427 344107 214469 344384 219766 344433 220826 344889 221191 344953 221527 345034 222454 345214 667596 1046762 668487 1075967 669163 1146395 669194 1147276	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, colled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32kDa BCL2 binding component 3 translocation protein 1 SH2 domain protein 2A zinc finger protein 451 homeo box A9 KIAA07563 gene product	AA227958 AA775838 AA228088 AA228088 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H84077 W72325 AA228129 AA644350 AA243818 AI821703 AA2436470 AA236470 AA236470 AA236477
АР-ћи АР-		LILLS SASASASASAS SASASASAS SASASAS SASASAS	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 12 2 3 4 5 6 7 8 9 10 11 2 1 2 3 4 5 6 7 8 9 10 11 2 1 2 3 4 5 6 7 8 9 10 11 2 1 12 2 3 4 5 6 7 8 9 10 11 2 1 12 1 2 3 4 5 6 7 8 9 10 11 12 12 12 12 12 12 12 12 12 12 12 12	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21 M22 M23 M24 N1 N2 N2 N2 N3 N4 N5 N6 N7 N2 N3 N4 N5 N6 N7 N2 N2 N3 N2 N2 N3 N2 N2 N3 N2 N2 N3 N2 N2 N3 N2 N3 N4 N5 N5 N6 N5 N6 N7 N2 N3 N4 N5 N5 N6 N7 N7 N8 N7 N5 N6 N7 N7 N8 N7 N7 N7 N8 N7 N7 N7 N7 N7 N7 N7 N7 N7 N7 N7 N7 N7	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 343318 214427 344107 214469 344384 219766 344443 220826 34489 221191 344953 221527 345034 222454 345034 22454 34503 1147276 669463 1147276 669496 1147276 669496 1147276 669496	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	AA227958 AA775838 AA775838 AA228088 AA555198 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H92126 AF086340 H92126 AF086340 H92126 AF086340 H84077 W72325 AA228129 AA644350 AA243818 AI821703 AA234778 AA234778
AP-hu AP-hu	01 01 001 001 001 001 001 001 001 001 0	LILLS SAMANANANAN SAMANAN SANANANANANANANANANANANANANANANANAN	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 23 24 1 2 3 4 5 6 7 8 9 10 11 12 23 24 1 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 12 2 3 4 12 2 3 4 12 2 3 4 12 2 3 4 12 2 3 4 12 2 12 2	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M17 M18 M19 M10 M21 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21 M22 M23 M24 N1 N2 N3 N4 N5 N6 N7 N8 N9 M10 M21	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 213414 343311 214144 343698 214427 344107 214469 344384 219766 344433 220826 344889 221191 344953 221527 345034 222454 345214 667596 1046762 668487 1075967 669163 1146395 669194 1147276 669496 1184337	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein as inhibitor, ciade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, colled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, ciade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 0 1 1ymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32kDa BCL2 binding component 3 translocation protein 1 SH2 domain protein 1 SH2 domain protein 1 SH2 domain protein 1 SH2 domain protein 451 homeo box A9 KIAA0763 gene product	AA227958 AA775838 AA228088 AA228088 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H84077 W72325 AA228129 AA644350 AA236470 AA236470 AA236470 AA2364719 AA234778 AA648413
AP-hu AP-hu	01 01 01 001 001 001 001 001 001 001 00	LILLS SASASASASAS SASASASAS SASASAS SASASAS	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 8 9 10 11 12 2 3 4 5 7 8 9 10 11 12 2 3 4 5 8 9 10 11 12 2 3 2 4 5 1 8 9 10 11 12 2 3 2 4 5 1 8 9 10 11 12 2 3 2 4 5 1 8 9 10 11 12 2 3 2 4 4 5 8 9 10 11 12 2 3 2 4 5 1 2 3 2 1 2 3 2 1 2 3 2 1 2 3 2 1 2 3 2 1 2 3 2 1 2 3 2 1 2 3 2 1 2 3 2 1 2 3 2 1 2 3 2 3	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21 M22 M23 M24 N1 N2 N2 N3 M22 N2 N1 N2 N3 N2 N1 N2 N3 N2 N1 N2 N3 N2 N1 N2 N3 N2 N1 N2 N2 N1 N2 N3 N2 N1 N3 N4 N5 N5 N6 N1 N1 N5 N6 N1 N7 N1 N5 N6 N1 N7 N1 N5 N6 N1 N7 N5 N6 N1 N7 N5 N6 N5 N6 N7 N7 N8 N1 N7 N5 N6 N1 N7 N5 N6 N7 N7 N8 N1 N7 N7 N8 N1 N7 N7 N8 N1 N7 N1 N1 N1 N1 N1 N1 N1 N1 N1 N1 N1 N1 N1	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 3433698 214427 344107 214469 344384 219766 344433 220826 344889 221191 344953 221527 345034 2221527 345034 22254 345034 222557 345034 22257 345034 22257 345034 221527 345034 212727 345034 212727 345034 212727 345034 212727 345034 212727 345034 212727 345034 212727 345034 212727 345034 212727 315037 3146395 31146395 669194 1147276 669496 1184337	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 1/mphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32kDa BCL2 binding component 3 translocation protein 1 SH2 domain protein 2A zinc finger protein 451 homeo box A9 KIAA0763 gene product Homo sapiens transcribed sequence with weak similarity to protein ref:NP_060312.1 (H.sapiens) hypothetical protein FLJ20489 [Homo sapiens]	AA227958 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H92126 AF086340 H92127 AF086340 H92127 AF086340 H92126 AF086340 H92127 AF086340 H92126 AF086340 H92127 AF086340 H92127 AF086340 H92127 AF086340 H92127 AF086340 H92177 AF086340 H92177 AF086340 H92177 AF086340 H92177 AF086340 H92177 AF086340 H92177 AF086340 H921777 AF0863477 AF0863477 AF0863477 AF0863477 AF0863477 AF0863477 AF08634777 AF0863477 AF086347778 AF086347778 AF086347778 AF086347778 AF086347778 AF086347778 AF086347778 AF086347778 AF086347778 AF08634778 AF086378 AF086378778 AF0863787787877878778778778778778778778778778
АР-ћи АР-ћ	01 01 001 001 001 001 001 001 001 001 0	LILLS SASSASSASS SASSASSASS SASSASSASSASSASS	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 23 24 1 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 12 2 3 4 5 6 7 8 9 10 11 12 12 12 2 3 4 5 6 7 8 9 10 11 12 12 12 12 12 12 12 12 12 12 12 12	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M17 M18 M19 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21 M22 M23 M24 N1 N2 N2 N3 N4 N5 N6 N7 N8 N9 M10 N10 N11	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 213414 343311 213414 343311 214144 343898 214427 344107 214469 344384 219766 344433 220826 344889 221191 344953 221527 345034 22254 345214 667596 1046762 668487 1075967 669163 1146395 669194 1147276 669496 1184337 681948	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, ciade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, colled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 ymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32kDa BCL2 binding component 3 translocation protein 2A zinc finger protein 451 homeo box A9 KIAA0753 gene product Homo sapiens transcribed sequence with weak similarity to protein ref:NP_60312.1 (H.sapiens) hypothetical protein FLJ20489 [Homo sapiens] fracilie histidine triad eane	AA227958 AA775838 AA228088 AA228088 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H84077 W72325 AA228129 AA6834058 AA234058 AA613070 AA236470 AA234778 AA648413 AA256123
АР-ћи АР-ћи		LILLS SESENTERS SERVICE STREETS SECTION STREETS SERVICE SECTION SECTIO	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 10 11 12 2 3 4 5 6 7 8 9 10 11 12 12 3 4 5 8 9 10 11 12 12 3 4 5 8 9 10 11 12 2 3 4 5 8 9 10 11 12 2 3 2 4 1 2 3 4 5 6 7 8 9 10 11 12 8 9 10 11 12 1 12 1 12 1 12 1 11 12 1 11 12 1 12 1 11 1	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21 M22 M23 M24 N1 N2 N3 M22 N3 M24 N5 N6 N7 N8 N9 N10 N1 N2 N3 M20 M21 M21 M22 M23 M24 M1 M20 M20 M21 M20 M20 M21 M20 M20 M21 M20 M21 M20 M21 M20 M21 M20 M21 M20 M21 M20 M21 M20 M21 M20 M21 M20 M21 M20 M21 M20 M20 M21 M20 M20 M21 M20 M20 M21 M20 M20 M21 M20 M20 M20 M20 M20 M20 M20 M20 M20 M20	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 213414 213414 213414 213414 213414 213414 213414 213414 214144 214469 344438 2214469 344443 220826 344889 221191 344953 221527 345034 221527 345034 222454 345034 221527 345034 214727 345034 214727 345034 214727 345034 214727 345034 214727 345034 214727 345034 214727 345034 214727 345034 214727 345034 214727 345034 214727 345034 2147337 3146395 311465 3114	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 1 1/mphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32kDa BCL2 binding component 3 translocation protein 1 SH2 domain protein 2A zinc finger protein 451 homeo box A9 KIAA0763 gene product Homo sapiens transcribed sequence with weak similarity to protein ref:NP_060312:1 (H.sapiens) hypothetical protein FLJ20489 [Homo sapiens] fragile histidine triad gene fragile histidine triad gene	AA227958 AA775838 AA228088 AA555198 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H92126 AF086340 H92126 AF086340 H92126 AF086340 H92126 AF086340 H84077 W72325 AA228129 AA644350 AA243818 AI821703 AA234058 AA234778 AA648413 AA256123
AP-hu AP-hu	01 01 001 001 001 001 001 001 001 001 0	LILLS SEESESSESSESSESSESSESSESSESSESSESSESSE	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 12 2 3 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 13 4 15 16 7 8 9 10 11 12 13 14 15 16 7 8 9 10 11 12 13 14 15 16 17 18 19 21 22 3 4 1 2 3 4 12 12 12 12 12 12 12 12 12 12 12 12 12	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21 M23 M24 N1 N2 N2 N2 N3 N4 N5 N6 N7 N8 N9 N10 N11 N12	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 214144 343311 214144 343311 214144 343598 214427 344107 214469 344384 219766 34489 221191 344953 221527 345034 222454 345214 667596 1046762 668487 1075967 669163 1146395 669194 1147276 669496 1184337 681948 1185189	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, colied-coli containing protein kinase 2 calcium/calmodulin-dependent protein kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 lymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein refr.NP_115594.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32KDa BCL2 binding component 3 translocation protein 1 SH2 domain protein 2A zinc finger protein 451 homeo box A9 KIAA0763 gene product Homo sapiens transcribed sequence with weak similarity to protein refixP_060312.1 (H.sapiens) hypothetical protein FLJ20489 [Homo sapiens] fragile histidine triad gene fragile X mental retardation 1	AA227958 AA775838 AA228088 AA228088 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H92126 AF086340 H84077 W72325 AA28129 AA644350 AA234058 AA613070 AA236470 AA2367512 AA648413 AA256123 AA648969
АР-ћи АР-ћи	01 01 01 001 001 001 001 001 001 001 00	LILLS SESERTED SERVICE SERVICES	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 2 3 2 4 5 6 7 8 9 10 11 2 2 3 2 4 1 2 3 2 4 1 2 3 4 5 6 7 8 9 10 11 2 2 3 2 4 5 6 7 8 9 10 11 2 2 3 2 4 5 8 9 10 11 2 2 3 2 4 5 1 8 9 10 11 2 2 3 2 4 5 1 8 9 10 11 2 2 3 2 2 3 2 3 4 5 6 7 8 9 10 11 2 2 3 2 2 3 2 2 3 2 3 2 3 4 1 2 3 2 3 2 3 2 1 2 3 2 3 2 3 2 3 2 1 2 3 2 3	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21 M22 M23 M24 N1 N2 N3 M22 M23 M24 N1 N2 N3 M22 M21 M22 M23 M24 N1 M22 M23 M24 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1	667407 878499 667440 1030527 210587 342614 210658 342633 212088 343111 213414 343311 213414 343311 214144 3433698 214427 344107 214469 344484 219766 344443 220826 344899 221191 344953 221527 345034 222454 345034 222454 345034 222454 345034 221527 345034 222454 345034 222454 345034 221527 315667 668487 1075967 668193 114535 668194 1147276 668496 1184337 681948 1185189 683915	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-c motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFA-Ilke effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 ymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukocyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32kDa BCL2 binding component 3 translocation protein 1 SH2 domain protein 2A zinc finger protein 451 homeo box A9 KIAA0763 gene product Homo sapiens transcribed sequence with weak similarity to protein ref:NP_060312.1 (H.sapiens) hypothetical protein FLJ20489 [Homo sapiens] fragile histidine triad gene fragile istidine triad gene fragile istidine triad gene fragile istidine triad gene	AA227958 AA775838 AA228088 AA555198 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H887 AF086340 H92126 AF086340 H887 AF086340 H84077 W72325 AA228129 AA644350 AA243818 AA644350 AA234778 AA648413 AA256123 AA64896124 AA64896144
АР-ћи АР-ћи		LILLS SASSASSASSASSASSASSASSASSASSASSASSASSA	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 2 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 5 6 7 8 9 10 11 12 2 3 4 12 2 3 4 12 2 3 4 12 3 4 12 12 12 10 11 12 2 3 4 12 2 3 4 12 2 3 4 12 2 3 4 12 2 3 4 12 3 4 12 10 11 12 2 3 4 12 12 10 11 12 2 3 4 12 3 4 12 12 12 10 11 12 2 3 4 12 12 12 12 12 12 12 10 11 12 2 3 4 12 12 12 12 12 12 12 12 12 12 12 12 12	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M17 M18 M17 M18 M17 M12 M20 M21 M21 M22 M23 M24 N1 N2 N2 N3 N2 N4 N5 N6 N7 N8 N2 N2 N2 N3 N2 N4 N5 N6 N7 N8 N2 N2 N2 N3 N2 N2 N3 N2 N2 N3 N2 N2 N3 N2 N2 N3 N2 N2 N3 N2 N2 N3 N2 N3 N2 N3 N2 N3 N4 N5 N5 N5 N5 N5 N5 N5 N5 N5 N5 N5 N5 N5	667407 878499 667440 1030527 210587 342614 210658 343111 213414 213414 343311 214144 343311 214144 343311 214144 343311 214144 343311 214144 343848 219766 344433 220826 344889 221191 344953 221527 345034 222454 345034 222454 345214 667596 1046762 668487 1075967 669163 1146395 669194 1147276 669496 1184337 681948 1185189 683915 1185189 1185	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	AA227958 AA775838 AA725838 AA775838 AA228088 AA555198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67448 H70847 W67448 H70847 W73784 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W7391 H91887 AF086339 H92126 AF086340 H84077 W72325 AA228129 AA6484350 AA243818 AI821703 AA243818 AI821703 AA243818 AI821703 AA2436470 AA236470 AA234778 AA648413 AA256123 AA648969 AA237088
АР-ћи АР-ћ		LILLA SASASASASAS SESSESSASA SECTOR ZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZ	21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 2 13 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 12 23 24 1 2 3 4 5 6 7 8 9 10 11 2 12 23 24 1 2 3 4 5 6 7 8 9 10 11 2 1 2 3 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 13 14 5 6 7 8 9 10 11 12 13 14 5 6 7 8 9 10 11 12 13 14 15 16 7 18 19 10 11 12 11 11	L21 L22 L23 L24 M1 M2 M3 M4 M5 M6 M7 M8 M9 M10 M11 M12 M13 M14 M15 M16 M17 M18 M19 M20 M21 M21 M22 M23 M24 N1 N2 N2 M21 M22 M23 M24 N1 N2 M24 N1 N2 M24 N1 N2 M24 N1 N2 M24 N1 M2 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1 M1	667407 878499 667407 878499 667400 1030527 210587 342614 210658 342633 212088 343111 213414 343311 2141427 344609 214427 344107 214469 344433 220826 3448953 221527 345034 222454 345214 667596 1046762 668487 1075967 1145395 669194 1147276 668496 1184337 681915 1185311	sperm associated antigen 9 hemoglobin, beta B aggressive lymphoma gene general transcription factor IIB sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D cysteine and glycine-rich protein 3 (cardiac LIM protein) 0 caspase recruitment domain family, member 10 ATP-binding cassette, sub-family G (WHITE), member 1 pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1 nerve growth factor receptor (TNFR superfamily, member 16) Rho-associated, coiled-coil containing protein kinase 2 calcium/calmodulin-dependent protein kinase kinase 1, alpha hypothetical protein MGC4309 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1 formin binding protein 3 cell death-inducing DFA-like effector a echinoderm microtubule associated protein like 1 thioredoxin-like 2 0 0 0 1 ymphoid enhancer-binding factor 1 Homo sapiens transcribed sequence with moderate similarity to protein ref:NP_115694.1 (H.sapiens) hypothetical protein MGC12945 [Homo sapiens] chemokine (C-X-C motif) ligand 14 leukcyte membrane antigen apoptosis antagonizing transcription factor myogenic factor 6 (herculin) KIAA0562 gene product TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32kDa BCL2 binding component 3 translocation protein 1 SH2 domain protein 2A zinc finger protein 451 homeo box A9 KIAA0763 gene product Homo sapiens transcribed sequence with weak similarity to protein ref:NP_060312.1 (H.sapiens) hypothetical protein FLJ20489 [Homo sapiens] fragile histidine triad gene fragile X mental retardation 1 centaurin, beta 2 glia maturation factor, beta	AA227958 AA775838 AA228088 AA525198 H65936 W68761 H66101 W68684 H68960 W67392 H72143 W67446 H70847 W69176 H77955 W73784 H73833 W73447 H81618 W73191 H95623 W76009 H91887 AF086340 H84077 W72325 AA228129 AA644350 AA24058 AA6435176 AA648413 AA648413 AA256123 AA648969 AA237088 AA651767

					SEC14L mRNA.	
AP-hu01	N	16	N16	1251853	KIAA0753 gene product	AA731225
AP-hu01	Ν	17	N17	686199	KIAA0053 gene product	AA262297
AP-hu01	N	18	N18	1270940	Homo sapiens transcribed sequence with moderate similarity to protein	AA748610
					pir:TVHUVV (H.sapiens) TVHUVV transforming protein vav - human (fragments)	
AP-hu01	N	19	N19	686216	apolipoprotein L, 3	AA262166
AP-hu01	N	20	N20	1307938	T-cell acute lymphocytic leukemia 2	AA767775
AP-hu01	N	21	N21	686463	KIAA0582 protein	AA255947
AP-hu01	N	22	N22	1318267	S100 calcium binding protein A5	AA810909
AP-hu01	N	23	N23	687208	DNA cross-link repair 1C (PSO2 homolog, S. cerevisiae)	AA259032
AP-hu01	N	24	N24	1322986	thiopurine S-methyltransferase	AA746081
AD-bu01		1	01	230126	vitronectin (serum spreading factor, somatomedin B, complement S-protein)	H78801
AP-NUUT	0			230120	significant (sorum oproducing racion, contractinicant b), compromotine o proteinty	W77812
AP-nu01	0	2	02	345856	cispiaun resistance associated	W77012
AP-hu01	0	3	03	233514	Homo sapiens transcribed sequences	H//320
AP-hu01	0	4	04	346016	procollagen-proline, 2-oxoglutarate 4-dioxygenase (proline 4-hydroxylase), alpha	W72635
			_		polypeptide I	
AP-hu01	0	5	O5	233939	NPC-related protein NAG73	AF085991
AP-hu01	0	6	O6	346146	branched chain keto acid dehydrogenase E1, alpha polypeptide (maple syrup urine	W78018
					disease)	
AP-hu01	0	7	07	242643	macrophage expressed gene 1	H94982
AP-hu01	0	8	O8	346604	advanced glycosylation end product-specific receptor	W74536
AP-hu01	0	9	O9	242770	metal-regulatory transcription factor 1	H94113
AP-hu01	0	10	O10	346628	S100 calcium binding protein A8 (calgranulin A)	W74581
AP-hu01	0	11	011	243341	CCCTC-binding factor (zinc finger protein)	N48105
AP-bu01	õ	12	012	346653	chromosome 5 open reading frame 13	W74451
	š	12	012	242526	KIAA0241 protein	N49464
AP-1001	0	13	013	243520		14104497
AP-hu01	0	14	014	346661	enolase 3, (peta, muscle)	W94107
AP-hu01	0	15	015	243554	Homo sapiens transcribed sequence with strong similarity to protein public PDH	149/94
	~		~ ~ ~	946990	(n.sapiens) & Chain G, nemoglobin (Deoxy, Human Fetal Fil)	W/70040
AP-hu01	0	16	016	346832		W/0010
AP-hu01	0	17	017	243741	gamma-aminobutyric acid (GABA) B receptor, 1	N49629
AP-hu01	0	18	O18	347092	COX17 homolog, cytochrome c oxidase assembly protein (yeast)	W79621
AP-hu01	0	19	O19	244970	PHD finger protein 3	N76260
AP-hu01	ō	20	020	347456	laminin, alpha 4	W81662
	õ	21	021	245015	NICE-4 protein	N72374
AP-bu01	ŏ	22	022	347736	solute carrier family 35 member F2	W81638
AP-HUUT	š	22	022	347750	solute carrier family 22 (extraneuronal monoamine transporter) member 3	N77230
AP-nu01	0	23	023	240009	solute carrier raining 22 (exclanedronal monoanime transporter), memory o	WAAGOE
AP-hu01	0	24	024	347761	src tamily associated phosphoprotein 2	W01005
AP-hu01	Р	1	P1	687234	casein kinase 1, epsilon	AA25/959
AP-hu01	Р	2	P2	1327016	SP110 nuclear body protein	AA724854
AP-hu01	Р	3	P3	687592	oncostatin M receptor	AA234544
AP-hu01	Р	4	P4	1335756	Homo sapiens transcribed sequences	AA828293
AP-hu01	Р	5	P5	700393	solute carrier family 35 (CMP-sialic acid transporter), member A1	AA290699
AP-hu01	P	6	P6	1336836	RNA binding motif protein 15	AA809402
AD-bu01		7	P7	700668	alvoine recentor, alpha 3	AA283885
AD 5001	'n	· ·	D0	100000	B coll CLI //mohome 11A /zinc finger protein)	AA914477
AP-hu01	P	8	20	1339205	B-cen CLDhymphoma i rA (zinc miger protein)	AA007450
AP-hu01	Р	9	P9	/0112/	Dasic leucine zipper nuclear factor 1 (JEM-1)	AA28/408
AP-hu01	Р	10	P10	1339831	regulatory factor X, 5 (influences HLA class II expression)	AA815144
AP-hu01	Р	11	P11	701272	T-cell leukemia/lymphoma 6	AA287767
AP-hu01	Р	12	P12	1339933	apoptosis-related protein PNAS-1	AA815207
AP-hu01	Р	13	P13	704600	tuberous scierosis 1	AA279895
AP-hu01	P	14	P14	1352486	Homo saniens transcribed sequences	AA829994
	D	15	D15	705056	hutvronhilin subfamily 1 member A1	AA279987
AD bu01	6	16	D16	1270282	SMC5 structural maintenance of chromosomes 5-like 1 (veast)	AA847845
	5	47	P10	7407202	similar to S. sentricing SSM	A A 200560
AP-nuui	P	17	P1/	/12/39	Similar to 5. Cerevisiae 55044	AA200000
AP-hu01	Р	18	P18	1370624	hypothetical protein LOC339091	AA833686
AP-hu01	Р	19	P19	713862	ubiquitin-protein isopeptide ligase (E3)	AA284599
AP-hu01	Р	20	P20	1371906	SKI-like	AA827643
AP-hu01	Р	21	P21	725269	secreted frizzled-related protein 4	AA291847
AP-hu01	Р	22	P22	1371945	Glucocorticoid receptor alpha mRNA, variant 3' UTR	AA827786
AP-hu01	P	23	P23	726563	transcription elongation factor A (SII), 2	AA394221
	P	24	P24	1388373	0	AA844124
		4	A4	1401702	HUS1 checkpoint homolog (S. nombe)	AA854050
	^		A1	3450330	viete anthrohisetosis vinus 526 anacana homaina 1 (mina)	ALAE7020
AF-0002	A,	4	A2	2100232		A 40000/0
AP-hu02	A	3	A3	1406233	transmemorane 4 supernamily member 1	AA889018
AP-hu02	A	4	A4	2150502	chemokine (C-C motif) ligand 22	AI457797
AP-hu02	Α	5	A5	1411935	Sapiens, Similar to phospholipase A2, group IB (pancreas), clone MGC:12509	AA921407
		_			IMAGE:3950450, mRNA, complete cds	
AP-hu02	Α	6	A6	2157583	apoptosis regulator BCL-G	AI478889
AP-hu02	Α	7	A7	1466849	renal tumor antigen	AA885446
AP-hu02	Α	8	A8	2162740	Fas apoptotic inhibitory molecule 2	AI468058
AP-hu02	A	9	A9	1467763	0	AA883106
AP-hu02	A	10	A10	2167853	phosphoinositide-3-kinase adaptor protein	A1582746
AP_hu02		11	A11	1469157	host cell factor C1 (VP16-accessory protein)	AA865726
	~	40	A40	2227274	DEAD/H (Aen_Glu_Ala_Aen/His) hav nalupantida 1	AIR00470
	^	12	A40	1470004	complement component 1 e subcomponent	AA070074
AP-0002	A	13	A13	14/2204	Complement Component 1, 5 subcomponent	MAD/32/1
AP-hu02	Α	14	A14	2242064	Homo sapiens transcribed sequence with strong similarity to protein sp:P25391	AI633741
AD 1: 25				4 507400	(H.sapiens) LMA1_HUMAN Laminin alpha-1 chain precursor (Laminin A chain)	
AP-hu02	A	15	A15	1507192	A I -nook transcription factor AKNA	AA904466
AP-hu02	Α	16	A16	2244541	G protein-coupled receptor 86	AI656746
AP-hu02	Α	17	A17	1520827	HIV TAT specific factor 1	AA910247
AP-hu02	Α	18	A18	2248373	I-kappa-B-interacting Ras-like protein 1	AI659150
AP-hu02	A	19	A19	1521793	tumor necrosis factor (ligand) superfamily, member 13b	AA906714
AP-hu02		20	A20	2271560	insulinoma-associated 1	AIGROR27
AP_h-002	Ā	24	A 24	1522/02	nostmeintic segrenation increased 2-like 6	V 1000051
	<u>,</u>	21	A21	1022482		AIC04047
AP-NU02	A	22	AZZ	22/3055	(A musculus) POL2 MOUSE Betraviaus related POL ask protein sp:P11369	AI081945
					(m.musculus) FOL2_MOUSE Retrovirus-related FOL polyprotein [Contains: Reverse transcriptase : Endosuclesse]	
4P-h-02	۸	22	A00	1503511	core binding factor, punt domain, alpha subunit 2: translocated to 1: audio D	AA00//252
	~	20	n23	1020011	related	101004000

•

AP-hu02	A	24	A24	2285313	guanine nucleotide binding protein (G protein), alpha 14	AI629022
AP-hu02	в	1	B1	470185	mitogen-activated protein kinase kinase kinase 7 interacting protein 1	AA028980
AP-hu02	в	2	82	1144761	low density lipoprotein receptor (familial hypercholesterolemia)	AA621897
AP-hu02	в	3	B3	471252	retinoic acld receptor, gamma	AA034461
AP-hu02	в	4	B4	1183847	syndecan binding protein (syntenin)	AA687473
AP-hu02	в	5	B5	486128	fibrillin 1 (Marfan syndrome)	AA043236
AP-hu02	в	6	B6	1258499	interleukin 23, alpha subunit p19	AA729815
AP-hu02	в	7	87	487402	polymerase (DNA directed), mu	AA046663
AP-hu02	в	8	B8	1272059	granzyme K (serine protease, granzyme 3; tryptase II)	AA743283
AP-hu02	в	9	B9	488909	histone deacetylase 5	AA057030
AP-hu02	в	10	B10	1285738	Homo sapiens transcribed sequence with weak similarity to protein	AA743682
	_				ref:NP_060312.1 (H.sapiens) hypothetical protein FLJ20489 [Homo sapiens]	A A 102100
AP-hu02	в	11	B11 B40	489800	serologically defined coort cancer analyen o	AA102109
AP-hu02	в	12	B12	1280223	Sections close IMAGE:5525036 mPNA	AA114041
AP-NUU2	8	13	D13	409070	Sapielis, cone invisionance of chromosomes 5-like 1 (vesst)	AA800572
AP-NU02	8	14	D14 D15	1200031	integrin beta-like 1 (with EGE-like repeat domains)	AA150407
AP-NUUZ		10	B16	1301810	ovidese (cytochrome c) assembly 1-like	AA767234
	0	17	B17	401727	Homo saplens transcribed sequence with weak similarity to protein	AA156691
	U	.,	017		ref:NP 060265.1 (H.sapiens) hypothetical protein FLJ20378 [Homo sapiens]	
AP-hu02	в	18	B18	1305734	Glucocorticoid receptor alpha mRNA, variant 3' UTR	AA736674
AP-hu02	в	19	B19	501883	fibromodulin	AA129908
AP-hu02	в	20	B20	1316966	Homo sapiens transcribed sequence with strong similarity to protein pir:CRHU5	AA767841
					(H.sapiens) CRHU5 carbonate dehydratase (EC 4.2.1.1) V precursor - human	
AP-hu02	в	21	B21	502257	cathepsin K (pychodysostosis)	AA156664
AP-hu02	в	22	B22	1321372	caspase recruitment domain family, member 10	AA/59191
AP-hu02	в	23	B23	502463	zinc finger protein 24 (KOX 17)	AA134//1
AP-hu02	8	24	B24	1324864	0 S	AA740500
AP-hu02	C	1	C1	1524563	S-antigen; retina and pineal gland (arrestin)	AA904007
AP-hu02	C	2	C2	2285/18	cytochrome P450, tamily 3, subtamily A, polypeptide 4	AI/00920
AP-hu02	C	3	C3	1540470	ubiquitin carrier protein	AA926110
AP-hu02	C	4	C4	229/925	EpnA4	AI913004
AP-hu02	C	5	C5	1542260	pyruvate dehydrogenase (lipoamide) alpha 2	AA927400
AP-hu02	C	6	06	2303920	death associated transcription factor 1	A1000041
AP-hu02	C	1	67	1543844	19A24 protein	AA921/00
AP-hu02	C	8	08	2304135	aminoadipate-semiaidenyde syntnase	A1001132
AP-hu02	C	9	C9	1551/65	Inteneukin 1, alpha	AA922009
AP-NU02	C	10	010	2315207	chemokine (C-A-C motif) ligand 6 (granulocyte chemotactic protein 2)	A1000047
AP-NU02	C o	11	011	100/028	regulator of G-protein signalling 9	AN910901
AP-NUUZ	C	12	CIZ	2310430	member 12	A1100001
AP-hu02	С	13	C13	1558715	platelet-derived growth factor alpha polypeptide	AA977022
AP-hu02	č	14	C14	2325602	cadherin 17, Ll cadherin (liver-intestine)	AI688206
AP-hu02	č	15	C15	1561976	homeo box C11	AA953456
AP-hu02	č	16	C16	2325850	solute carrier family 6 (neurotransmitter transporter, taurine), member 6	AI688443
AP-hu02	č	17	C17	1566020	SCAN domain containing 1	AI073530
AP-hu02	č	18	C18	2330876	complement component 1, q subcomponent, receptor 1	AI693483
AP-hu02	Ċ	19	C19	1570420	chemokine (C-C motif) ligand 1	AA931884
AP-hu02	С	20	C20	2341256	aryl hydrocarbon receptor nuclear translocator	AI697689
AP-hu02	С	21	C21	1579639	interferon, gamma	AA969504
AP-hu02	С	22	C22	2348576	killer cell lectin-like receptor subfamily C, member 2	AI798774
AP-hu02	С	23	C23	1581406	CD80 antigen (CD28 antigen ligand 1, B7-1 antigen)	AA983817
AP-hu02	С	24	C24	2349996	TAF1 RNA polymerase II, TATA box binding protein (TBP)-associated factor,	AI806456
	_				250kDa	
AP-hu02	D	1	D1	503206	immunoglobulin mu binding protein 2	AA148925
AP-NU02	D	2	D2	1325272	cytochrome c oxidase subunit viic	AA/41399
AP-nu02	5	3	03	503280	Inteneukin 19 wirsless has MATE (integration site family, member 104	AA101733
AP-NUU2	5	4	D4	1320381	wingless-type MMI V integration site family, member 10A	AA131/62
AP-huuz	5	5	05	1003/40	chroniosome o open reading name 4	AA131403
AP-11002	5	7	07	F04926	transcription factor A mitochondrial	AA150777
	ň	é		1335806	CLIC triplet repeat RNA binding protein 1	AA828368
	ň	0	D0	505259	cochome P450 family 11 subfamily A notypentide 1	AA142870
	ň	10	D10	1330537	hutvronbilin subfamily 1 member A1	AA805147
	ň	11	D10	667009	SWI/SNE related matrix associated actin dependent regulator of chromatin	AA253109
-1002	5		511	00,000	subfamily a, member 1	
AP-hu02	D	12	D12	1340493	formin binding protein 3	AA938206
AP-hu02	D	13	D13	667614	histone deacetylase 6	AA228073
AP-hu02	D	14	D14	1350983	glia maturation factor, beta	AA806487
AP-hu02	D	15	D15	668465	ADP-ribosyltransferase (NAD+; poly (ADP-ribose) polymerase)-like 1	AA243734
AP-hu02	D	16	D16	1352169	0	AA808204
AP-hu02	D	17	D17	668977	suppressor of cytokine signaling 4	AA253351
AP-hu02	D	18	D18	1355977	SET domain, bifurcated 2	AA831594
AP-hu02	D	19	D19	669261	Sapiens, clone IMAGE:4214177, mRNA	AA236520
AP-hu02	D	20	D20	1357778	ubiquitin-protein isopeptide ligase (E3)	AA832340
AP-hu02	D	21	D21	669310	mitogen-activated protein kinase-activated protein kinase 5	AA236814
AP-hu02	D	22	D22	1371215	aiscs, iarge (Drosophila) nomolog 1	AA847183
AP-hu02	0	23	D23	683358	nomo sapiens transcribed sequences	AA215307
AP-hu02	5	24	U24	1372313	receptor-interacting senne-threonine kinase 2	AA826052
AP-hu02	E	1	E1	1584882	ankyrin repeat and SUCS box-containing 4	AI/91680
AP-hu02	Ë	2	E2	2354198	macrophage scavenger receptor 1	AI/34941
	5	3	E3	1000010	prosphonositide-o-kinase adaptor protein	A1007000
AP-hu02	E	4	E4	2357389	CUTD anugen, D polypepude	AI62/808
AP-002	E	5	E5	1089786	norociast growth factor 9 (gila-activating factor)	AA946776
AP-0002	E	0	E0	2309664	cas-cir-m (munne) ecoropic retroviral transforming sequence	A 4000000
AP-0002	5	/ 0	E/	100294/	Unonionine (U-U mour) liganu o	ANJ09000
AP-11002	Ē	0		2309300	iyinphotyto alphamilaso laukotriana A4 hydrolaea	A 4007000
AP_h-02	5	10	E3 E10	2270474	A kingse (PRKA) anchor omtein (untigo) 0	AI761707
	6	10	EIV	2010111	A Relieve (r rave) and lot proton (Jourde) a	AU V 11 21

	F	11	E11	1621044	ubiquitously transcribed tetratricopeptide repeat gene. Y chromosome	AI004861
AP-hu02	F	12	E12	2379167	vanin 1	AI762340
	F	13	E13	1625719	protein tyrosine phosphatase, receptor type, D	AI005089
AP-bu02	F	14	E14	2382863	cvtochrome P450, family 3, subfamily A, polypeptide 5	AI759985
AP-hu02	F	15	E15	1634832	class I cytokine receptor	AI088984
	F	16	E16	2383199	notassium inwardly-rectifying channel, subfamily J, member 1	AI795925
AP-hu02	F	17	E17	1638431	postmeiotic segregation increased 2-like 5	AI015925
AP-hu02	F	18	E18	2384363		AI830536
	Ē	10	E10	1641007	0	AI040940
	5	20	E20	2386044	Kninnel-like factor 2 (lund)	A1765962
	Ē	21	E21	1643606	protein transport protein SEC61 alpha subunit isoform 1	AI023590
	5	21	E22	2386478	NP220 nuclear protein	A1808939
	5	22	E22	1645371	autochome D450 family 17 subfamily & polynentide 1	AI028740
	5	20	E24	2202865	crantume M (jumphocute met-ace 1)	AIQ10778
	5	1	E1	683822	ATP-binding cassette sub-family E (GCN20) member 1	AA255836
AP-11002	г с	2	F1 F2	1373030	Seniere CDNA EL 133408 fie, clone BRACE2010550, highly similar to Human cell	AA825685
AF-11002	Г	4	12	13/3030	arowth regulator CGR19 mRNA.	74020000
AP-hu02	F	3	F3	684760	hypothetical protein MGC4175	AA251625
AP-hu02	F	4	F4	1387641	glutathione S-transferase A3	AA846513
AP-hu02	F	5	F5	684917	BUB1 budding uninhibited by benzimidazoles 1 homolog beta (yeast)	AA251909
AP-hu02	F	6	F6	1401913	histone deacetvlase 7A	AA854147
AP-bu02	F	7	F7	685470	tumor necrosis factor receptor superfamily, member 14 (herpesvirus entry	AA262421
	•	•	••		mediator)	
AP-hu02	F	8	F8	1405643	KIAA0562 gene product	AA890549
AP-hu02	F	9	F9	685554	macrophage stimulating, pseudogene 9	AA262037
AP-hu02	F	10	F10	1405890	I-kappa-B-interacting Ras-like protein 1	AA890397
AP-hu02	F	11	F11	687301	chemokine (C-X-C motif) ligand 16	AA235226
AP-hu02	F	12	F12	1407260	thymic stromal lymphopoietin	AA889581
AP-hu02	F	13	F13	687934	translocation protein 1	AA236805
AP-hu02	F	14	F14	1412597	carboxypeptidase A1 (pancreatic)	AA835150
AP-hu02	F	15	F15	700704	ankyrin repeat and SOCS box-containing 3	AA285075
AP-hu02	F	16	F16	1435368	PRKC, apoptosis, WT1, regulator	AA834474
AP-hu02	F	17	F17	703508	0	AA278673
AP-hu02	F	18	F18	1435643	death associated protein 3	AA858184
AP-hu02	F	19	F19	703643	tripartite motif-containing 14	AA278637
AP-hu02	F	20	F20	1455450	chemokine (C-X-C motif) ligand 14	AA865643
AP-hu02	F	21	F21	704456	deoxyribonuclease I-like 3	AA279835
AP-hu02	F	22	F22	1467195	cytochrome P450, family 11, subfamily B, polypeptide 1	AA884709
AP-hu02	F	23	F23	712161	histone 1. H2bk	AA280257
AP-hu02	F	24	F24	1492385	phosphoinositide-3-kinase adaptor protein	AA894359
AP-hu02	G	1	G1	1651122	ubiquitin specific protease 15	AI091570
AP-hu02	Ğ	2	G2	2394148	solute carrier family 1 (neuronal/enithelial high affinity glutamate transporter	AI763334
	τ.	-	-		system Xaq), member 1	
AP-hu02	G	3	G3	1662143	sphingomyelin phosphodiesterase 2, neutral membrane (neutral sphingomyelinase)	AI083508
AP-hu02	G	4	G4	2404737	T cell receptor alpha locus	AI816828
AP-hu02	G	5	G5	1662507	KIAA0169 protein	AI083541
AP-hu02	G	6	G6	2406104	interleukin 1 family, member 9	AI828999
AP-hu02	G	7	G7	1663139	interleukin 11	AI128942
AP-hu02	~	-		2406442	UL16 binding protein 1	AI830832
	G	8	G8			
AP-hu02	G	8 9	G8 G9	1663164	0	AI129113
AP-hu02 AP-hu02	G G G	8 9 10	G8 G9 G10	1663164 2409058	0 B cell RAG associated protein	Al129113 Al819384
AP-hu02 AP-hu02 AP-hu02	G G G G	8 9 10 11	G8 G9 G10 G11	1663164 2409058 1664589	0 B cell RAG associated protein collagen, type XI, alpha 1	AI129113 AI819384 AI041918
AP-hu02 AP-hu02 AP-hu02 AP-hu02	G G G G G G	8 9 10 11 12	G8 G9 G10 G11 G12	1663164 2409058 1664589 2429773	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila)	Al129113 Al819384 Al041918 Al858773
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	66666	8 9 10 11 12 13	G8 G9 G10 G11 G12 G13	1663164 2409058 1664589 2429773 1672339	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15	Al129113 Al819384 Al041918 Al858773 Al090427
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		8 9 10 11 12 13 14	68 G9 G10 G11 G12 G13 G14	1663164 2409058 1664589 2429773 1672339 2460033	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1	Al129113 Al819384 Al041918 Al858773 Al090427 Al936239
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		8 9 10 11 12 13 14 15	G8 G9 G10 G11 G12 G13 G14 G15	1663164 2409058 1664589 2429773 1672339 2460033 1673209	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IdG, low affinity Illa, receptor for (CD16)	Al129113 Al819384 Al041918 Al858773 Al090427 Al936239 Al141618
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		8 9 10 11 12 13 14 15 16	G8 G9 G10 G11 G12 G13 G14 G15 G16	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylolutaryI-coenzyme A reductase	Al129113 Al819384 Al041918 Al858773 Al090427 Al936239 Al141618 Al962842
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	66666666666	8 9 10 11 12 13 14 15 16 17	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsvnaotic protein CRIPT	AI129113 AI819384 AI041918 AI858773 AI090427 AI936239 AI141618 AI962842 AI086377
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		8 9 10 11 12 13 14 15 16 17 18	G8 G9 G10 G11 G12 G13 G14 G15 G16 G16 G17 G18	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, low affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryI-Coenzyme A reductase postsynaptic protein CRIPT T-cell Iymohoma invasion and metastasis 1	AI129113 AI819384 AI041918 AI858773 AI090427 AI936239 AI141618 AI962842 AI086377 AW003835
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		8 9 10 11 12 13 14 15 16 17 18 19	68 69 610 612 613 614 615 616 617 618 619	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryI-coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo saniens transcribed sequence with strong similarity to protein pir:B46012	AI129113 AI819384 AI041918 AI858773 AI090427 AI936239 AI141618 AI962842 AI086377 AW003835 AI142438
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	66666666666666	8 9 10 11 12 13 14 15 16 17 18 19	68 69 610 612 613 614 615 616 617 618 619	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryI-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment)	Al129113 Al619384 Al041918 Al658773 Al990427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		8 9 10 11 12 13 14 15 16 17 18 19 20	68 69 611 612 613 614 615 616 617 618 619 620	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryI-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6	Al129113 Al819384 Al041918 Al858773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		8 9 10 11 12 13 14 15 16 17 18 19 20 21	68 69 610 612 613 614 615 616 617 618 619 620 621	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein	Al129113 Al819384 Al041918 Al858773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		8 9 10 11 12 13 14 15 16 17 18 19 20 21 22	68 69 610 611 612 613 614 615 616 617 618 619 620 621 622	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5	Al129113 Al819384 Al041918 Al058773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23	68 69 610 611 612 613 614 615 616 617 618 619 620 621 622 623	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryI-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase	Al129113 Al819384 Al041918 Al058773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	68 69 610 611 612 613 614 615 616 617 618 619 620 621 622 622 623 624	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 1703954 2490644 1704556 2495108	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryI-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) low density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6	Al129113 Al819384 Al041918 Al658773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972744 Al962515 AW001035
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	тооооо ооооооооооо	8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G19 G20 G21 G22 G23 G24 H1	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein	Al129113 Al819384 Al041918 Al058773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AW001035 AA281913
AP-hu02 AP-hu02	ттоосоо осососоосоосо	8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2	68 69 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 H1 H2	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5	Al129113 Al819384 Al041918 Al658773 Al090427 Al936239 Al141618 Al96239 Al141618 Al966377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AW001035 AA281913 AA894914
AP-hu02 AP-hu02	тттооооо оооооооооооо	8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3	68 69 610 611 612 613 614 615 616 617 618 620 621 622 623 624 H1 H2 H3	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 712943	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryI-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit)	Al129113 Al819384 Al041918 Al041918 Al058773 Al090427 Al36239 Al141618 Al362842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AW001035 AA894914 AA282380
AP-hu02 AP-hu02	ттттосос сососососос	8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 23 24 1 2 3 4	68 69 610 611 612 613 614 615 616 617 618 620 621 622 623 624 H1 H2 H3 H4	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 2490644 1704556 2495108 712441 1493445 712943 1504500	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) low density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2	Al129113 Al819384 Al041918 Al058773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972744 Al962615 AW001035 AA284913 AA282380 AA904907
AP-hu02 AP-hu02	ттттеооооо оооооооооооо	8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3 4 5	68 69 610 611 612 613 614 615 616 617 618 619 621 622 621 622 623 624 H1 H2 H3 H3 H4	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 712943 1504500 713619	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryI-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) low density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein')	Al129113 Al819384 Al041918 Al658773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AW001035 AA281913 AA894914 AA282380 AA904907 AA283840
AP-hu02 AP-hu02	тттттооооо оооооооооттттт	8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3 4 5 6	68 69 610 611 612 613 614 615 616 617 618 621 622 623 624 H1 H2 H3 H4 H5 H6	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 712943 1504550	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein') protein phosphatase 2 (formerly 2A), regulatory subunit B", alpha	Al129113 Al819384 Al041918 Al058773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AW001035 AA281913 AA894914 AA282380 AA904907 AA283840 AA906152
AP-hu02 AP-hu02	ττττττοοοοο οοοοοοοοοιτ	8 9 10 11 12 13 14 15 16 17 18 20 21 22 23 24 1 2 3 4 5 6 7	68 69 610 611 612 613 614 615 616 617 618 620 621 622 623 624 H1 H2 H3 H4 H5 H6 H7	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712943 1504500 713619 1504851 714050	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, low affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) low density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein') protein phosphatese 2 (formerly 2A), regulatory subunit B", alpha uridine morophosphate synthetase (orotate phosphoribosyl transferase and	Al129113 Al819384 Al041918 Al058773 Al090427 Al36239 Al141618 Al362842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AA281913 AA894914 AA282380 AA904907 AA283840 AA904152 AA284909
AP-hu02 AP-hu0	: ттттттооооо оооооооооооо	8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3 4 5 6 7 2	68 69 611 612 613 614 615 616 617 618 620 622 623 624 H1 H2 H3 H4 H5 H6 H7 	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 1703954 1703954 1703954 2490644 1704556 2495108 712441 1493445 712943 1504500 713619 1504851 714050	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein') protein phosphatase 2 (formerly 2A), regulatory subunit B", alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and protein directed).	Al129113 Al819384 Al041918 Al058773 Al090427 Al306239 Al141618 Al362842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972744 Al962515 AW001035 AA281913 AA894914 AA282380 AA904907 AA283840 AA906152 AA284909
AP-hu02 AP-hu0	т ттттттооооо ооооооооооооо	8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 3 4 5 6 7 8	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G17 G18 G21 G21 G22 G23 G24 H1 H2 H3 H4 H5 H6 H7 H8	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 1504500 713619 1504851 714050 1508334	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) low density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein') protein phosphatase 2 (formerly 2A), regulatory subunit B", alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and orotidine-5'-decarboxylase) jagged 1 (Alagille syndrome)	Al129113 Al819384 Al041918 Al658773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972744 Al160261 Al972744 Al086515 AW001035 AA281913 AA394914 AA282380 AA904907 AA283840 AA904907 AA283840 AA904052 AA284909 AA947272
AP-hu02 AP-hu02	тт ттттттооооо осоосоосоосо	8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 4 5 6 7 8 9 2	68 69 610 611 612 613 614 615 616 617 618 620 621 622 623 624 H1 H2 H3 H4 H5 H6 H7 H8 H9	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 712943 15045500 713619 1504851 714050 1508334 723784	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor (docking protein') protein phosphates 2 (formerly 2A), regulatory subunit B", alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and orotidine-5'-decarboxylase) peroxisome proliferative activated receptor, delta	Al129113 Al819384 Al041918 Al658773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AW001035 AA281913 AA894914 AA282380 AA904907 AA283409 AA949075 AA284909 AA947272 AA292445
AP-hu02 AP-hu02	τττι τιτιτοοροο οσοσοσοσοσοσοσο	8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3 4 5 6 7 8 9 10	68 69 610 611 612 613 614 615 616 617 618 620 621 622 623 624 H1 H2 H3 H4 H5 H6 H7 H8 H9 H10	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 712943 1504851 713619 1504851 714050	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, low affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) low density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein') protein phosphatese 2 (formerly 2A), regulatory subunit B", alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and cortidine-5'-decarboxylase) jagged 1 (Alagille syndrome) peroxisome proliferative activated receptor, delta vacuolar protein sorting 4B (yeast)	Al129113 Al819384 Al041918 Al058773 Al090427 Al36239 Al141618 Al36242 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AW001035 AA281913 AA894914 AA282380 AA904907 AA282380 AA904907 AA282380 AA904907 AA282380 AA904552 AA284909 AA9472722 AA292445 AA916662
AP-hu02 AP-hu0	ττττ ττττοοοοο οοοοοοοοοοοοοοοο	8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3 4 5 6 7 8 9 10 11	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G20 G21 G22 G23 G24 H1 H2 H3 H4 H5 H6 H7 H8 H9 H10 H11	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 1703954 1703954 1703954 1703954 1704556 2495108 712441 1493445 712943 1504500 713619 1504851 714050 1508334 723784 1527121 724388	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphome invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor (docking protein') protein phosphatase 2 (formerly 2A), regulatory subunit B", alpha uridine monophosphate syntheses (orotate phosphoribosyl transferase and orotidine-5'-decarboxylase) jagged 1 (Alagille syndrome) peroxisome proliferative activated receptor, delta vacuolar protein sorting 4B (yeast) EphA2	AI129113 AI819384 AI041918 AI041918 AI090427 AI396239 AI141618 AI362842 AI086377 AW003835 AI142438 AI972744 AI160261 AI972744 AI9602615 AW001035 AA281913 AA894914 AA9282380 AA904907 AA283840 AA906152 AA284909 AA947272 AA292445 AA916562 AA911065
AP-hu02 AP-hu0	τττττ ττττοοοοο οοοοοοοοοοοοοοο	8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 3 4 5 6 7 8 9 10 11 22 23 24 1 2 3 4 5 6 7 8 9 10	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G17 G18 G20 G21 G22 G23 G24 H1 H2 H3 H4 H5 H6 H7 H8 H9 H10 H11 H12	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 1504500 713619 1504851 714050 1508334 723784 1527121 724368 1534940	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein') protein phosphatase 2 (formerly 2A), regulatory subunit B", alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and cortidine-5'-decarboxylase) jagged 1 (Alagille syndrome) peroxisome protein sorting 4B (yeast) EphA2 BcI-2-associated transcription factor	AI129113 AI819384 AI041918 AI858773 AI090427 AI936239 AI141618 AI962842 AI086377 AW003835 AI142438 AI972744 AI160261 AI972744 AI160261 AI972925 AI096515 AW001035 AA284901 AA282380 AA904907 AA283840 AA904907 AA283840 AA904907 AA283840 AA904907 AA283840 AA904907 AA284909 AA947272 AA292445 AA916562 AA411065 AA918867
AP-hu02 AP-hu0		8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 4 5 6 7 8 9 10 11 22 3 4 5 6 7 8 9 10 11	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G20 G21 G22 G23 G24 H1 H2 H3 H4 H5 H6 H7 H8 H9 H10 H11 H12 H13	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 712943 1504851 714050 1508334 723784 1527121 1508334 723784 1524940 726388	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein') protein phosphates 2 (formerly 2A), regulatory subunit B", alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and orotidine-5'-decarboxylase) peroxisome proliferative activated receptor, delta vacuolar protein sorting 4B (yeast) EphA2 BcI-2-associated transcription factor tripartite motif-containing 29	Al129113 Al819384 Al041918 Al658773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AW001035 AA281913 AA894914 AA282380 AA904907 AA283840 AA904907 AA283840 AA904907 AA283840 AA904907 AA283840 AA904907 AA283840 AA904152 AA284909 AA947272 AA292445 AA916562 AA916562 AA916567 AA918667 AA293775
AP-hu02 AP-hu0		8 9 10 11 12 13 14 15 16 17 18 19 20 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G20 G21 G22 G23 G24 H1 H2 H3 H4 H5 H6 H7 H8 H9 H10 H11 H112 H13 H14	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 712943 1504851 714050 1508334 723784 1527121 724368 1534940 726388 1534940	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, low affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) low density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein') protein phosphates 2 (formerly 2A), regulatory subunit B", alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and cortidine-5'-decarboxylase) jagged 1 (Alagille syndrome) peroxisome proliferative activated receptor, delta vacuolar protein sorting 4B (yeast) EphA2 Bcl-2-associated transcription factor tripartite motif-containing 29 S-antigen; retina and pineal gland (arrestin)	Al129113 Al819384 Al041918 Al058773 Al090427 Al36239 Al141618 Al36242 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AW001035 AA284914 AA282380 AA904907 AA282380 AA904907 AA282380 AA904907 AA282380 AA904907 AA282380 AA904552 AA284909 AA9472722 AA292445 AA916562 AA491867 AA91867 AA293775 AA918179
AP-hu02 AP-hu0		8 9 10 11 12 13 15 16 17 18 9 20 22 23 24 1 2 3 4 5 6 7 8 9 10 11 23 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 13 4 15 16 17 18 19 10 11 12 13 4 15 16 17 18 19 10 11 12 13 4 15 16 17 18 19 20 12 22 23 24 1 2 3 4 5 6 7 7 8 9 10 11 12 13 4 15 16 17 18 19 10 11 12 13 14 15 16 17 18 19 20 12 21 22 23 24 12 23 24 2 23 24 12 23 24 2 23 24 2 23 24 2 23 24 2 2 2 2	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G20 G21 G22 G23 G24 H1 H2 H3 H4 H5 H6 H7 H8 H9 H10 H11 H112 H113 H14 H15	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 1703954 1703954 1703954 1703954 1703954 1704556 2495108 712441 1493445 712943 1504500 713619 1504500 713619 1504851 714050 1508334 723784 1527121 724368 1534940 726388 1543210 728133	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor (docking protein') protein phosphatase 2 (formerly 2A), regulatory subunit B", alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and orotidine-5-decarboxylase) jagged 1 (Alagille syndrome) peroxisome proliferative activated receptor, delta vacuolar protein sorting 4B (yeast) EphA2 BcI-2-associated transcription factor tripartite motif-containing 29 S-antigen; retina and pineal gland (arrestin) chromosome 12 open reading frame 22	AI129113 AI819384 AI041918 AI058773 AI090427 AI36239 AI141618 AI962842 AI086377 AW003835 AI142438 AI972744 AI160261 AI972744 AI9602615 AW001035 AA281913 AA894914 AA282380 AA906152 AA283840 AA906152 AA283840 AA906152 AA284909 AA947272 AA292445 AA916562 AA918667 AA293775 AA293775 AA91877 AA435633
AP-hu02 AP-hu0		8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 3 4 5 6 7 8 9 10 11 22 3 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 13 4 15 16 7 18 19 10 11 12 13 14 15 16 7 18 19 10 11 12 13 14 15 16 7 18 19 10 11 12 13 14 15 16 7 18 19 20 21 22 3 4 5 6 7 8 9 10 11 12 13 14 15 16 7 18 19 20 11 12 11 12 11 14 15 16 7 18 19 20 11 12 23 24 1 2 23 24 1 2 2 3 4 5 6 7 8 9 10 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 11	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G17 G18 G20 G21 G22 G22 G22 G22 H1 H2 H3 H5 H6 H7 H8 H9 H11 H12 H11 H12 H11 H15 H16	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 712943 1504500 713619 1504851 714050 1508334 723784 1527121 724368 1534940 726388 1534940 728133 1543859	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein') protein phosphates 2 (formerly 2A), regulatory subunit B*, alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and orotidine-5'-decarboxylase) jagged 1 (Alagille syndrome) peroxisome protein sorting 4B (yeast) EphA2 BcI-2-associated transcription factor tripartite motif-containing 29 S-antigen; retina and pineal gland (arrestin) chromosome 12 open reading frame 22 signal recognition particle 54kDa	Al129113 Al819384 Al041918 Al658773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972744 Al160261 Al972744 Al160261 Al972744 Al160261 Al96515 AA284909 AA945152 AA284909 AA947272 AA292445 AA918662 AA918667 AA918867 AA91879 AA93179 AA921832
AP-hu02 AP-hu0	τιτττττττι τιτττικοοορο ουοοοοοοοοοοοοοοο	8 9 10 11 12 13 14 15 16 17 18 9 21 22 23 4 5 6 7 8 9 10 11 23 4 5 6 7 8 9 10 11 23 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 14 15 16 7 7 8 9 10 11 12 3 14 15 16 7 7 8 9 10 11 12 3 14 15 16 7 7 8 9 10 11 12 3 14 15 16 7 7 8 9 10 11 12 3 11 12 3 14 15 16 7 7 8 9 10 11 12 3 11 12 11 12 11 12 11 12 11 11 12 11 11	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G20 G21 G22 G23 H1 H2 H3 H4 H5 H6 H7 H8 H9 H10 H11 H12 H13 H14 H15 H16 H17	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 712943 1504851 714050 1508334 723784 1527121 1508334 723784 152488 1534940 726388 1534940 726388 1543859 729886	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukernia 2 signal recognition particle receptor ('docking protein') protein phosphates 2 (formerly 2A), regulatory subunit B*, alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and orotidine-5*-decarboxylase) peroxisome proliferative activated receptor, delta vacuolar protein sorting 4B (yeast) EphA2 BcI-2-associated transcription factor tripartite motif-containing 29 S-antigen; retina and pineal gland (arrestin) chromosome 12 open reading frame 22 signal recognition particle for KADa ATP-binding cassette, sub-family C (CFTR/MRP), member 1	Al129113 Al819384 Al041918 Al058773 Al090427 Al336239 Al141618 Al362842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AA281913 AA394914 AA282380 AA904907 AA283840 AA904907 AA283840 AA9049052 AA284909 AA947272 AA29445 AA916562 AA411065 AA91867 AA91867 AA931877 AA435633 AA9412168
AP-hu02 AP-hu0		8 9 10 11 12 13 14 15 16 17 18 9 20 12 22 23 4 5 6 7 8 9 10 11 23 4 5 6 7 8 9 10 11 12 34 5 6 7 8 9 10 11 12 34 5 6 7 8 9 10 11 12 34 15 16 17 18 19 10 11 12 34 15 16 17 18 19 20 12 22 23 24 12 34 5 6 7 8 9 10 11 12 34 15 16 17 18 19 20 12 21 22 23 24 12 34 5 6 7 8 9 10 11 12 34 15 16 17 18 19 20 12 21 22 23 24 12 34 5 6 7 8 9 10 11 12 34 15 16 17 18 19 20 11 12 34 15 16 17 18 19 20 12 20 12 20 12 20 10 11 12 34 11 12 11 12 11 11 11 12 11 11 11 11 11	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G21 G22 G23 G24 H1 H2 H3 H45 H6 H7 H10 H11 H112 H13 H14 H15 H16 H17 H18	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 1703954 1703954 1703954 1703954 1703954 1703954 1703954 1703954 1703954 1703954 1703954 1504500 713619 1504851 714050 1508334 723784 1527121 724368 1534940 726388 1543210 728133 1543859 729886 1555042	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, low affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) low density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein') protein phosphatese 2 (formerly 2A), regulatory subunit B", alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and cortidine-5'-decarboxylase) jagged 1 (Alagille syndrome) peroxisome proliferative activated receptor, delta vacuolar protein sorting 4B (yeast) EphA2 Bcl-2-associated transcription factor tripartite motif-containing 29 S-antigen; retina and pineal gland (arrestin) chromosome 12 open reading frame 22 signal recognition particle 54kDa ATP-binding cassette, sub-family C (CFTR/MRP), member 1 par-6 partitioning defective 6 homolog alpha (C.elegans)	Al129113 Al819384 Al041918 Al058773 Al090427 Al36239 Al141618 Al36242 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AW001035 AA284901 AA904515 AA284909 AA947272 AA292445 AA916562 AA411065 AA918867 AA291775 AA91887 AA91837 AA91837 AA91837 AA91837 AA91837 AA91837 AA91837 AA91837 AA91837 AA91837 AA91837 AA91837 AA91837 AA91837 AA91837 AA91832 AA9182 AA9182 AA9182 AA9182 AA9182 A9182 AA9182 A918 A9184 A918 A918 A918 A918 A918 A918 A918 A918
AP-hu02 AP-hu0		8 9 10 11 12 13 4 15 16 17 18 9 20 12 23 24 1 2 3 4 5 6 7 8 9 10 11 21 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 11 12 13 4 15 16 17 18 19 20 12 22 23 24 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 10 11 12 10 11 12 10 11 12 10 11 12 11 11 12 11 11 12 11 11 12 11 11	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G20 G22 G22 G23 G24 H1 H2 H3 H4 H5 H6 H7 H18 H10 H11 H112 H14 H15 H16 H17 H18 H19 H10	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 1703954 1703954 1703954 1703954 1704556 2495108 712441 1493445 712943 1504500 713619 1504851 714050 1508334 723784 1527121 724368 1543210 726388 1543210 728133 1543859 729886 1555042 730149	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein receptor-related protein 6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor (docking protein') protein phosphatase 2 (formerly 2A), regulatory subunit B", alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and crotidine-5-decarboxylase) jagged 1 (Alagille syndrome) peroxisome proliferative activated receptor, delta vacuolar protein sorting 4B (yeast) EphA2 Bcl-2-associated transcription factor tripartite motif-containing 29 S-antigen; retina and pineal gland (arrestin) chromosome 12 open reading frame 22 signal recognition particle 54kDa ATP-binding cassette, sub-family C (CFTR/MRP), member 1 par-6 partitioning defective 6 homolog alpha (C.elegans) transcription elongation factor A (SII), 2	AI129113 AI819384 AI041918 AI041918 AI058773 AI090427 AI336239 AI141618 AI362842 AI086377 AW003835 AI142438 AI972744 AI160261 AI972744 AI160261 AI972744 AI066515 AW001035 AA281913 AA894914 AA282380 AA904907 AA283840 AA906152 AA284909 AA947272 AA282445 AA916562 AA411065 AA918867 AA293775 AA292445 AA918867 AA293775 AA918179 AA435633 AA921832 AA91283 AA91283 AA914 AA914 AA914 AA914 AA914 AA914 AA914 AA914
AP-hu02 AP-hu0		8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 3 4 5 6 7 8 9 10 11 23 14 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 10 11 12 3 4 5 6 7 11 12 3 4 5 6 7 11 12 3 4 5 6 7 11 12 3 4 5 6 7 11 12 3 4 5 6 7 11 12 3 14 15 16 7 17 18 19 20 11 12 3 14 15 16 7 18 19 10 11 12 3 14 15 16 7 18 19 20 11 12 20 20 11 12 20 20 11 12 20 11 12 20 11 12 20 20 11 12 20 20 11 12 20 20 20 11 12 20 20 11 12 20 20 11 12 20 20 11 12 20 20 11 12 20 20 11 12 20 20 11 12 20 20 11 12 20 20 11 12 20 20 11 12 20 20 11 12 20 20 11 12 20 20 20 11 12 20 20 11 12 20 20 11 12 20 20 11 12 20 20 20 20 20 20 20 20 20 20 20 20 20	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G17 G18 G21 G22 G22 G22 G22 G22 H1 H2 H3 H5 H6 H7 H8 H9 H11 H12 H13 H15 H15 H16 H17 H15 H16 H17 H17 H17 H17 H17 H17 H17 H17 H17 H17	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 16787 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 1504500 713619 1504851 714050 1508334 723784 1534940 726388 1543859 729886 1553042 730149 1562070	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity Illa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein') protein phosphatase 2 (formerly 2A), regulatory subunit B", alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and orotidine-5'-decarboxytase) jagged 1 (Alagille syndrome) peroxisome protiferative activated receptor, delta vacuolar protein sorting 4B (yeast) EphA2 BcI-2-associated transcription factor tripartite motif-containing 29 S-antigen; retina and pineal gland (arrestin) chormosome 12 open reading frame 22 signal recognition particle 54kDa ATTP-binding cassette, sub-family C (CFTR/MRP), member 1 par-6 partitioning defective 6 homolog alpha (C.elegans) transcription elongation factor A (SII), 2 pyruvate dehydrogenase (lipoamide) alpha 1	Al129113 Al819384 Al041918 Al058773 Al090427 Al936239 Al141618 Al962842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972744 Al962515 AW001035 AA281913 AA894914 AA282380 AA904907 AA283840 AA906152 AA284909 AA947272 AA292445 AA918662 AA918667 AA918667 AA918667 AA918667 AA918667 AA91867 AA91867 AA91867 AA91879 AA92445 AA91879 AA924518 AA921832 AA412613 AA933938
AP-hu02 AP-hu0	οοοοοοοοοοοοοοοοοοοοοοοοοοοοοοοοοοοοοο	8 9 10 11 12 13 14 15 16 17 18 9 10 11 12 13 14 15 16 7 8 9 10 11 12 13 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 14 15 16 7 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 11 12 11 14 15 16 7 18 19 10 11 12 2 2 3 4 5 6 7 11 12 2 3 4 5 6 7 10 11 12 3 4 5 16 7 11 12 10 11 12 11 11 11 11 11 11 11 11 11 11 11	G8 G9 G10 G11 G12 G13 G14 G15 G16 G17 G18 G20 G21 G22 G22 G22 H1 H2 H3 H4 H5 H6 H7 H11 H12 H113 H14 H15 H16 H17 H18 H19 H111 H15 H16 H17 H18 H19 H112 H112 H112 H112 H112 H112 H112	1663164 2409058 1664589 2429773 1672339 2460033 1673209 2474074 1678175 2478677 1697461 2490395 1703954 2490644 1704556 2495108 712441 1493445 712943 1504851 714050 713619 1504851 714050 1508334 723784 1527121 724388 1534940 726388 1534940 726388 1543210 728133 1543859 729886 1555042 730149 1562070 730845	0 B cell RAG associated protein collagen, type XI, alpha 1 frizzled homolog 2 (Drosophila) caspase recruitment domain family, member 15 transcription elongation factor A (SII), 1 Fc fragment of IgG, Iow affinity IIIa, receptor for (CD16) 3-hydroxy-3-methylglutaryl-Coenzyme A reductase postsynaptic protein CRIPT T-cell lymphoma invasion and metastasis 1 Homo sapiens transcribed sequence with strong similarity to protein pir:B46012 (H.sapiens) B46012 pp52/LSP1 homolog 2 - human (fragment) Iow density lipoprotein receptor-related protein 6 adhesion glycoprotein apoptosis inhibitor 5 lipoprotein lipase homeo box B6 succinate dehydrogenase complex, subunit D, integral membrane protein homeo box C5 polymerase (DNA directed), epsilon 2 (p59 subunit) T-cell acute lymphocytic leukemia 2 signal recognition particle receptor ('docking protein') protein phosphates 2 (formerly 2A), regulatory subunit B*, alpha uridine monophosphate synthetase (orotate phosphoribosyl transferase and orotidine-5'-decarboxytase) peroxisome proliferative activated receptor, delta vacuolar protein sorting 4B (yeast) EphA2 BcI-2-associated transcription factor tripartite motif-containing 29 S-antigen; retina and pineal gland (arrestin) chromosome 12 open reading frame 22 signal recognition particle for 6 AkDa ATP-binding cassette, sub-family C (CFTR/MRP), member 1 par-6 partilloning defective 6 homolog alpha 1 NADH dehydrogenase (ubiquinone) 1, alpha/beta subcomplex, 1, 8kDa	Al129113 Al819384 Al041918 Al058773 Al090427 Al36239 Al141618 Al362842 Al086377 AW003835 Al142438 Al972744 Al160261 Al972925 Al096515 AA281913 AA394914 AA282380 AA904907 AA283840 AA904907 AA283840 AA904907 AA283840 AA904907 AA283840 AA904907 AA283840 AA904907 AA283840 AA904907 AA283840 AA91652 AA218057 AA218657 AA218657 AA218657 AA218657 AA218657 AA218657 AA218657 AA218657 AA218657 AA21867 AA218275 AA31867 AA31875 AA318775 AA33840 AA31867 AA31875 AA318775 AA31775 AA318775 AA33375 AA318775 AA33375 AA318775 AA318775 AA33375 AA318775 AA33375 AA318775 AA318775 AA33375 AA3177575 AA3337575 AA31775757575757757757757777777777

AP-hu02	н	23	H23	730902	Mdm4, transformed 313 cell double minute 4, p53 binding protein (mouse)	AA416645
AP-hu02	н	24	H24	1562173	RNA binding motif protein 15	AA933971
AP_hu02	1	1	11	1716220	0	AI160585
AP 1002	-		10	0506522	ting finger protein subfamily 1A 3 (Ajolos)	AW071111
AP-NUU2	1	2	12	2536535	Zinc iniger protein, subtaining i.e. s (violas)	ANG / 1 / 1
AP-hu02	1	3	13	1721224	inositol polyphosphate-5-phosphatase, 40kDa	AI151146
AP-hu02	1	4	14	2544446	angiopoletin 1	AW044503
	i	5	15	1731840	myoglobin	AI184297
		5	10	0507507	a distance and matellanetainana domain 15 (materaidin)	AN072264
AP-hu02		6	16	2567597	a disintegrin and metalloproteinase domain 15 (metalgidin)	AVV0/2304
AP-hu02	1	7	17	1742445	chemokine (C-C motif) ligand 26	AI186047
AP_hu02	1	8	18	2605431	protein phosphatase 1, regulatory (inhibitor) subunit 3A (glycogen and	AW118744
74 11402	•	v			sarconlasmic reticulum binding subunit, skeletal muscle)	
AD 5002		0	10	1751372	home how A10	AI187278
AP-HUUZ	1		19	1751572		ANN/404004
AP-hu02	1	10	110	2662365	B7 nomolog 3	AW 181901
AP-hu02	1	11	111	1752018	fms-related tyrosine kinase 3	AI150354
AP-hu02	1	12	112	2663077	recombination activating gene 1	AW183306
	÷ .	42	14.2	1755900	regulatory factor V 3 (influences HI & class II expression)	A1203838
AP-nuuz	1	13	113	1100090	regulatory factor X, 3 (initiances FILA class in expression)	A1203030
AP-hu02	1	14	114	2690595	ret finger protein 2	AW237758
AP-hu02	1	15	115	1847208	adrenomedullin receptor	AI243295
	1	16	116	2011855	III 16 binding protein 2	AW510737
AD 500	-	10	147	4040400	ciello said hisdina like lostin 7	A1247571
AP-nuuz	1	17	117	1849100	static acto binding ignike recurr /	A1247071
AP-hu02	1	18	118	2945121	matrix metalloproteinase 20 (enamelysin)	AW593141
AP-hu02	1	19	119	1854028	Down syndrome cell adhesion molecule	AI243628
AD 5002	i i	20	120	2045274	Ken37 notein	AW503206
AP-11002	!	20	120	2940274		1040404
AP-hu02	1	21	121	1857875	alpha-1-microglobulin/bikunin precursor	AI246161
AP-hu02	1	22	122	2975903	interleukin 21 receptor	AW629191
AP-hu02	1	23	123	1870748	0	AI245910
	:	20	12.0	0743030	Annual and the second	A14/407000
AP-nu02	1	24	124	2/1/0/0	transforming growth factor, beta receptor I (activity A receptor type n-like kinase,	AW 13/000
					53kDa)	
AP-hu02	J	1	J1	730933	protein phosphatase 2, regulatory subunit B (B56), beta isoform	AA421193
AP-hu02	J	2	J2	1565637	HUS1 checkpoint homolog (S. pombe)	AI167268
AD hu02	ĩ	-	12	740286	DP1-associated protein 1 (negative cofactor 2 alpha)	AA476942
APHILUZ	3	3	33	140200	Divise Social de Protein - (Indgalive Colacior 2 alpha)	AA070342
AP-hu02	J	4	J4	1585823	alpha-fetoprotein	AA9/6/55
AP-hu02	J	5	J5	741974	0	AA405401
AP-hu02	.1	6	.16	1587251	paired-like homeodomain transcription factor 2	AA977103
AD 5000	Ť.	7	17	744120		AA63/317
AP-huuz	J		37	/44120		AA034317
AP-hu02	J	8	J8	1587390	interleukin 12 receptor, beta 2	AA977194
AP-hu02	J	9	J9	744130	wingless-type MMTV integration site family, member 3	AA634318
AP-hu02	.i.	10	.110	1588397	annexin A6	AA975225
	š.			746490		A A 44 7022
AP-nu02	J	11	J11	740109	BCL0 co-repressor	MM41/033
AP-hu02	J	12	J12	1609989	ankyrin repeat and SOCS box-containing 1	AI001012
AP-hu02	J	13	J13	753215	guanine nucleotide binding protein (G protein), alpha inhibiting activity polypeptide	AA406420
	•					
AD 5000		14	14.4	1620057	, core binding factor, sunt domain, sinhs subunit 2: translocated to 1: ovelin D-	44002855
AFMIUUZ	J	14	314	1020007		77332000
					related	
AP-hu02	J	15	J15	753734	TATA box binding protein (TBP)-associated factor, RNA polymerase I, C, 110kDa	AA411726
AP-hu02	J	16	J16	1626874	cysteine and glycine-rich protein 3 (cardiac LIM protein)	AI018334
	Ĩ.	17	117	752045	disce Jarge (Drosophile) homolog 5	AA470265
AP-1002	3	17	317	100040		AVA47 3200
AP-hu02	J	18	J18	1627091	retinoic acid early transcript 1K pseudogene	AI017460
AP-hu02	J	19	J19	754351	hyaluronoglucosaminidase 1	AA436276
AP-hu02	.1	20	.120	1627998	leukemia inhibitory factor receptor	AI014497
AD hu02	ň	24	104	756500	audia (nucleosido disherabata lisked meiab) V) time metit 1	A A 4 4 4 0 2 0
AP-nuuz	J	21	JZI	/ 50502	nudix (nucleoside diprosphate linked molety X)-type motil 1	MM444020
AP-hu02	J	22	J22	1628761	interleukin 1 family, member 7 (zeta)	Al014548
AP-hu02	J	23	J23	758266	thrombospondin 4	AA437064
AP-hu02	i.	24	124	1631832	Ren1 quaning nucleotide exchange factor directly activated by cAMP	A1004302
AF41002	3	24	J24	1031032	rap I guainne incledute dataining factor in edga duvated by chair	A1004302
AP-nuu2	ĸ	1	K1	18/4500	caimodulin 1 (phosphorylase kinase, delta)	AI281330
AP-hu02	κ	2	K2	2719958	interleukin 21 receptor	AW205118
AP-hu02	к	3	K3	1875345	transmembrane 4 superfamily member 3	AI274190
AD hu02	~	-	KA	2720244		A14/129002
AP-nu02	<u> </u>	4	N4	2720344	Notal homolog 5 (brosphila)	AW 136903
AP-hu02	ĸ	5	K5	1880926	nuclear RNA export factor 1	AI268362
AP-hu02	к	6	K6	2721127	matrix metalloproteinase 28	AW207107
AP-hu02	к	7	K7	1882591	SWI/SNE related matrix associated actin dependent regulator of chromatin	A1279006
AFHIOZ	N.	'	N	1002031	subfemily a member A	AI2/ 3000
AD	ĸ		20	2724452	nratein turneine nhaenhatase, recentor time. L	AW201524
	n.	0	NO	2129902	protein tyruaine phuaphataaa, teteptui type, n	ATT281034
AP-hu02	ĸ	9	K9	1899192	chemokine (C-C motif) receptor-like 2	AI288845
AP-hu02	к	10	K10	3032869	chemokine-like receptor 1	AW771926
AP-hu02	к	11	K11	1912380	glutathione peroxidase 2 (gastrointestinal)	AI304515
AD 5002			K40	0005000		ANUT70004
AP-NUU2	N.	12	K12	3035088	KOUNOAOKINASE (ITUCIOKINASE)	AVV//9891
AP-hu02	κ	13	K13	1913168	Homo sapiens transcribed sequence with moderate similarity to protein	AI304610
					ref:NP_060265.1 (H.sapiens) hypothetical protein FLJ20378 [Homo sapiens]	
AP-hu02	к	14	K14	2734295	0	AW449304
AP-hu02	ĸ	15	K15	1917820	cytidine deaminase	A1343863
				0700000	avelagenie segentes DOV like 4	A14/45/000
AP-nu02	ĸ	16	K16	2/36931	purmergic receptor M2X-like 1, orpnan receptor	AW451665
AP-hu02	κ	17	K17	1926309	death associated protein 3	Al346132
AP-hu02	к	18	K18	3053174	postmeiotic segregation increased 2-like 6	AW401404
AP-hung	ĸ	10	K10	1031876	verh-h2 enthrohlastic leukemia viral oncorene homeles 2 (orien)	A1322040
AF-11002	<u> </u>	19	118	19310/0	verb-oz eryukoulasuc leukenika viral oncogene nomolog 3 (avian)	A1333219
AP-hu02	ĸ	20	K20	3054559	pranched chain alpha-ketoacid dehydrogenase kinase	AW575117
AP-hu02	κ	21	K21	1934125	cytochrome P450, family 11, subfamily B, polypeptide 1	AI334352
AP-hu02	ĸ	22	KOO	3059780	nolymerase (DNA directed) lembde	AW675005
AD Luco	Ň		1400	4020054	perjana anavidana	AI044546
AP-NU02	ĸ	23	K23	1939054	squalene epoxidase	AI341546
AP-hu02	ĸ	24	K24	3061074	ankyrin repeat and SOCS box-containing 1	AW576120
–		1	11	773190	KIAA0763 gene product	44429474
AP-hu02	È	•	- · ·	1636043	nuclear chambonatoin similar to Successful a DM/D4	AI047070
AP-hu02	L	•	10	10.30613	nuclear prosproprotein similar to 5. cerevisiae PWP1	AIU1/373
AP-hu02 AP-hu02	L	2	L2			
AP-hu02 AP-hu02 AP-hu02	L L L	2 3	L2 L3	773274	frizzled-related protein	AA425521
AP-hu02 AP-hu02 AP-hu02 AP-hu02		2 3 4	L2 L3 L4	773274 1639806	frizzled-related protein postmeiotic segregation increased 2-like 5	AA425521 AI025190
AP-hu02 AP-hu02 AP-hu02 AP-hu02		2 3 4 5	L2 L3 L4	773274 1639806 781222	frizzied-related protein postmeiotic segregation increased 2-like 5 TGERI-induced anti-apontalic fonter 1	AA425521 Al025190
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		2 3 4 5	L2 L3 L4 L5	773274 1639806 781222	frizzled-related protein postmeiotic segregation increased 2-like 5 TGFB1-induced anti-apoptotic factor 1	AA425521 Al025190 AA446223
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		2 3 4 5 6	L2 L3 L4 L5 L6	773274 1639806 781222 1643585	frizzied-related protein postmeiotic segregation increased 2-like 5 TGFB1-induced anti-apoptotic factor 1 guanine nucleotide binding protein (G protein), alpha activating activity polypeptide,	AA425521 Al025190 AA446223 Al024119
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		2 3 4 5 6	L2 L3 L4 L5 L6	773274 1639806 781222 1643585	frizzied-related protein postmeiotic segregation increased 2-like 5 TGFB1-induced anti-apoptotic factor 1 guanine nucleotide binding protein (G protein), alpha activating activity polypeptide, olfactory type	AA425521 Al025190 AA446223 Al024119
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		2 3 4 5 6 7	L2 L3 L4 L5 L6 L7	773274 1639806 781222 1643585 781575	frizzled-related protein postmeiotic segregation increased 2-like 5 TGFB1-induced anti-apoptotic factor 1 guanine nucleotide binding protein (G protein), alpha activating activity polypeptide, olfactory type B aggressive lymphoma gene	AA425521 Al025190 AA446223 Al024119 AA431459
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		2 3 4 5 6 7 8	L2 L3 L4 L5 L6 L7 L8	773274 1639806 781222 1643585 781575 1644971	frizzied-related protein postmeiotic segregation increased 2-like 5 TGFB1-induced anti-apoptotic factor 1 guanine nucleotide binding protein (G protein), alpha activating activity polypeptide, offactory type B aggressive lymphoma gene SKI-like	AA425521 Al025190 AA446223 Al024119 AA431459 Al025801
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02		2 3 4 5 6 7 8 9	L2 L3 L4 L5 L6 L7 L8 L8	773274 1639806 781222 1643585 781575 1644971 784532	frizzied-related protein postmeiotic segregation increased 2-like 5 TGFB1-induced anti-apoptotic factor 1 guanine nucleotide binding protein (G protein), alpha activating activity polypeptide, olfactory type B aggressive lymphoma gene SKI-like phosphofurdokinase, muscle	AA425521 Al025190 AA446223 Al024119 AA431459 Al025801 AA437412

AP-hu02 AP-hu02	1	40				
AP-hu02		10	L10	1647990	pyridoxal (pyridoxine, vitamin B6) kinase	AI033704
	L	11	L11	784726	C-type (calcium dependent, carbohydrate-recognition domain) lectin, superfamily	AA478515
	-				member 12	
AP-hu02	1	12	L12	1649917	N-acetvitransferase 1 (arviamine N-acetvitransferase)	AI022585
AP-hu02	ī	13	113	785141	serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 2	AA476471
	-	13	144	4050630	CDC6 coll division guals 6 homolog (6, corovision)	AI052065
AP-nu02	L	14	L14	1002032		AI032003
AP-hu02	L	15	L15	785345	optic atrophy 1 (autosomai dominant)	AA476505
AP-hu02	L	16	L16	1653944	chemokine (C-C motif) ligand 8	AI085069
AP-hu02	L	17	L17	788566	Purkinje cell protein 4	AA452966
AP-hu02	-	18	1 1 8	1662047	mitogen-activated protein kinase kinase kinase 7 interacting protein 2	AI082790
AP	-	10	140	705075	Auros protein p52 indusible protein 3	AA464040
AP-huuz	L	19	LIA	/952/5	tumor protein poo inducidie protein o	AV4451610
AP-hu02	L	20	L20	1662532	0	AI083552
AP-hu02	L	21	L21	796085	biglycan	AA460811
AP-hu02	L	22	L22	1662927	Homo sapiens transcribed sequence with strong similarity to protein sp:Q9Y2Y9	AI074707
	-				(H.sapiens) KLFD_HUMAN Krueppel-like factor 13 (Transcription factor BTEB3) (Basic transcription element binding protein 3) (BTE-binding protein 3) (RANTES factor of late activated T lymphocytes-1) (RFLAT-1) (Transcription factor NSLP1) (Novel Sp1-like zinc finger transcription factor 1) (Transcription factor NSLP1)	
AP-hu02	L	23	L23	796446	laminin, beta 2 (laminin S)	AA460076
AP-hu02	L	24	L24	1663533	protein phosphatase 2 (formerly 2A), regulatory subunit A (PR 65), beta isoform	AI034287
AP-hu02	м	1	M1	1941548	0	AI206322
AP-hu02	M	2	M2	3068359	transient recentor potential cation channel, subfamily V, member 6	AW452394
AD bu02	M	-	142	1042624	programmed cell death 1 ligand 1	A1202006
AP-huuz	M	3	MO	1942034	programmed ceil death i ligand i	AI202990
AP-hu02	м	4	M4	3070861	a disintegrin and metalloproteinase domain 8	BF513621
AP-hu02	м	5	M5	1943890	feline sarcoma oncogene	AI199245
AP-hu02	м	6	M6	3071092	adrenergic, beta, receptor kinase 2	BF513491
AP-hu02	M	7	M7	1944395	thrombovane A synthese 1 (platelet cytochrome P450 family 5 subfamily A)	AI202694
AD hu02		,	149	2721709	firming homolog 0 (Desception)	AM/207262
AP-NU02	м	•	MO	2/31/96	mzzied nomolog a (Drosophila)	AVV29/203
AP-hu02	м	9	M9	1946205	jun D proto-oncogene	AI339181
AP-hu02	м	10	M10	3172977	zinc finger protein 145 (Kruppel-like, expressed in promyelocytic leukemia)	BE222963
AP-hu02	м	11	M11	1948222	PPAR binding protein	AI352382
AD 5002	14	12	1412	2175742	chemokine (C motif) ligand 2	BE219461
AP-11002	M	12	M12	31/5/45		DE2 10401
AP-hu02	м	13	M13	2012241	leukocyte membrane antigen	AI356/22
AP-hu02	M	14	M14	3197927	fer (fps/fes related) tyrosine kinase (phosphoprotein NCP94)	BE464122
AP-hu02	м	15	M15	2029551	stanniocalcin 1	AI367280
AP-hu02	м	16	M16	3212988	activating transcription factor 5	BE467506
AD bu02	M	17	M17	2032630	leukon te immunoglobulin-like recentor, subfamily A (with TM domain), member 1	A1703206
AP-1002	N1	17	M117	2032039	disadely and the second s	AIT 93200
AP-0002	м	18	M18	3213997	diacyigiycerol kinase, gamma 90kDa	BE468027
AP-hu02	м	19	M19	2055309	B aggressive lymphoma gene	Al307751
AP-hu02	M	20	M20	3219673	alkaline phosphatase, intestinal	BE550404
AP-hu02	м	21	M21	2055594	interleukin 18 recentor accessory protein	AI307682
	N4	20	1400	2000004	interfection 10 receptor accessory protein	DE 435474
AP-huuz	M	22	MZZ	3200021		DF4331/4
AP-hu02	M	23	M23	2061969	interleukin 1 family, member 7 (zeta)	AI343258
AP-hu02	м	24	M24	3323544	tumor necrosis factor receptor superfamily, member 9	BF064069
AP-hu02	N	1	N1	796471	secretory leukocyte protease inhibitor (antileukoproteinase)	AA460434
AP-hu02	N	2	N2	1669661	chromosome 1 open reading frame 16	AI057273
AD hu02	N	2	AI2	900597	Bhe anesific quantino pusicatido evaluando fostar e114	A A 456630
AP-nuuz	N	3	NJ	609567	Rho-specific guarine nucleotide exchange factor p114	AA456630
AP-hu02	N	4	N4	1670127	HIV TAT specific factor 1	AI074508
AP-hu02	N	5	N5	810137	fructose-1,6-bisphosphatase 1	AA464360
AP-hu02	N	6	N6	1671227	general transcription factor IIIA	AI086037
AP-hu02	N	7	N7	810255	G protein_coupled recentor kinase_interactor 1	AA464724
				4674700	b protein coupled receptor kinds and a receptor in the actor in	AI075774
AD hung			NIO	10/4/00	nypolielical protein MGC4043	AU/5//1
AP-hu02	N	0	N8			
AP-hu02 AP-hu02	N N	9	N8 N9	810672	SD6 antigen	AA463987
AP-hu02 AP-hu02 AP-hu02	N N N	9 10	N8 N9 N10	810672 1674873	8D6 antigen T-cell acute lymphocytic leukemia 1	AA463987 Al076543
AP-hu02 AP-hu02 AP-hu02 AP-hu02	N N N N	9 10 11	N8 N9 N10 N11	810672 1674873 810777	SDS antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4	AA463987 Al076543 AA481753
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	2 2 2 2	9 10 11 12	N8 N9 N10 N11 N12	810672 1674873 810777 1675856	8D6 antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein	AA463987 AI076543 AA481753 AI078341
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	~ ~ ~ ~ ~ ~	9 10 11 12 12	N8 N9 N10 N11 N12	810672 1674873 810777 1675856	8D6 antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAEE illia BNA columerase II, p200/CBB associated factor (PCAE) associated	AA463987 AI076543 AA481753 AI078341
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	N N N N N	9 10 11 12 13	N8 N9 N10 N11 N12 N13	810672 1674873 810777 1675856 811810	SDS antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor (PCAF)-associated	AA463987 Al076543 AA481753 Al078341 AA463470
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	N N N N N	9 10 11 12 13	N8 N9 N10 N11 N12 N13	810672 1674873 810777 1675856 811810	SDS antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa	AA463987 AI076543 AA481753 AI078341 AA463470
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	N N N N N N N N	9 10 11 12 13 14	N8 N9 N10 N11 N12 N13 N14	810672 1674873 810777 1675856 811810 1676286	8D6 antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB	AA463987 Al076543 AA481753 Al078341 AA463470 Al075694
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	N N N N N N N	9 10 11 12 13 14 15	N8 N9 N10 N11 N12 N13 N14 N15	810672 1674873 810777 1675856 811810 1676286 812139	SDS antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde	AA463987 Al076543 AA481753 Al078341 AA463470 Al075694 AA456021
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	9 10 11 12 13 14 15	N8 N9 N10 N11 N12 N13 N14 N15	810672 1674873 810777 1675856 811810 1676286 812139	SDS antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase)	AA463987 AI076543 AA481753 AI078341 AA463470 AI075694 AA456021
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	9 10 11 12 13 14 15 16	N8 N9 N10 N11 N12 N13 N14 N15 N16	810672 1674873 810777 1675856 811810 1676286 812139 1677559	SDS antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111	AA463987 AI076543 AA481753 AI078341 AA463470 AI075694 AA456021 AI078579
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	~~~~~	9 10 11 12 13 14 15 16 17	N8 N9 N10 N12 N13 N14 N15 N16 N17	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991	SDS antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter	AA463987 AI076543 AA481753 AI078341 AA463470 AI075694 AA456021 AI078579 AA455702
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	~~~~~	9 10 11 12 13 14 15 16 17 18	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249	SDS antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0	AA463987 AI076543 AA481753 AI078341 AA463470 AI075694 AA456021 AI078579 AA455702 A142896
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	~~~~~~	9 10 11 12 13 14 15 16 17 18 19	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234	SDS antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit	AA463987 AI076543 AA481753 AI078341 AA463470 AI075694 AA456021 AI078579 AA455702 AI42896 AA465233
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	9 10 11 12 13 14 15 16 17 18 19 20	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1670255	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antinen	AA463987 AI076543 AA481753 AI078341 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA455233 AI068455
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	~~~~~	9 10 11 12 13 14 15 16 17 18 19 20 21	N8 N9 N10 N12 N13 N14 N15 N16 N17 N18 N19 N20	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aidehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	~~~~~	9 10 11 12 13 14 15 16 17 18 19 20 21	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0	AA463987 AI076543 AA481753 AI078341 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456 AA465022
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	~~~~~~	9 10 11 12 13 14 15 16 17 18 19 20 21 22	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456 AA459022 AI079412
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	~~~~~~	9 10 11 12 13 14 15 16 17 18 19 20 21 22 23	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N20 N21 N22 N23	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651	ADS antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aidehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456 AA465223 AI066456 AA459022 AI079412 AA481094
AP-hu02 AP-hu02	~~~~~~	9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1678256 814358 1680119 814651 1682318	SD6 antigen T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4	AA463987 AI076543 AA481753 AI078341 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456 AA459022 AI079412 AA481094 AI066675
AP-hu02 AP-hu02	222222 22 222222 0	9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2065463	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collageen, type VI. aloha 2	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA455702 AI142896 AA455022 AI079412 AA481094 AI066675 AI375553
AP-hu02 AP-AP-AP-AP-AP-AP-AP-AP-AP-AP-AP-AP-AP-A	222222 22 2222222200	5 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 22 23 24 1 2	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2065463 3072354	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 tips finger protein 147 (setrogen responsive finger protein)	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456 AA465233 AI066456 AA459022 AI079412 AA481094 AI066675 AI375553 BC221641
AP-hu02 AP-hu02	222222 22 2222222200	9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 22 23 24 1 2	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2065463 3072364	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein)	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA455022 AI078452 AI066456 AA459022 AI079412 AA481094 AI066675 BG231641
AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02 AP-hu02	OOOZZZZZZ ZZ ZZZZZZZ	5 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 23 24 1 2 3	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814338 1680119 814651 1682318 2065463 3072364 2067693	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI075699 AA455702 AI142896 AA465233 AI066456 AA465233 AI066456 AA459022 AI079412 AA481094 AI0666755 AI375553 BG231641 AI383639
AP-hu02 AP-hu02	0000ZZZZZZZZZZZZZZZZZ	5 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 22 23 24 1 2 3 4	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N18 N20 N21 N21 N22 N23 N24 O1 O2 O2 O2 O2 O2	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2065463 3072364 2067693 3073056	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456 AA465233 AI066456 AA459022 AI079412 AA481094 AI066675 AI375553 BG231641
AP-hu02 AP-hu0	00000222222222 22 222222	9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 22 3 4 5	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3 O4 O5	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2065463 3072364 2067693 3073056 2090446	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA459022 AI079412 AA481094 AI066675 AI375553 BG231641 AI383639 BG231641 AI554887
AP-hu02 AP-hu0	0000022222222222 22 222222	9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 23 4 5 6	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3 O4 O5 O6	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2065463 3072364 2067693 3073056 2090446	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apontotic factor 1	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456 AA465233 AI066456 AA459022 AI079412 AA481094 AI066675 AI375553 BG231641 AI383639 BG231641 AI383639
AP-hu02 AP-hu0	000000222222222222222222222222222222222	9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 23 4 5 6 7	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3 O4 O5 O6 O5	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2065463 3072364 2067693 3073056 2090446 3078407	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456 AA465233 AI066456 AA459022 AI079412 AA481094 AI066675 AI375553 BG231641 AI383639 BG231641 AI554887 BG231641
AP-hu02 AP-hu0	000000022222222222222222222222222222222	9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3 4 5 6 7 2	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N21 N22 O3 O4 O1 O2 O3 O4 O5 O6 O6 O7	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814234 1680119 814651 1682318 2065463 3072364 2067693 3073056 2090446 3078407 2094946	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 olymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA455022 AI142896 AA459022 AI079412 AA481094 AI079412 AA481094 AI079412 AA481094 AI079553 BG231641 AI383639 BG231641 AI383639 BG231641 AI554887 BG231641 AI424264
AP-hu02 AP-hu02	000000022222222222222222222222222222222	9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 23 4 5 6 7 8	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3 O4 O5 O6 O7 O8	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814324 1679256 814358 1680119 814651 1682318 2065463 3073056 2090446 3078407 2094946 3083083	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0	AA463987 AI076543 AA481753 AI078341 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456 AA465233 AI066456 AI375553 BG231641 AI383639 BG231641 AI454887 BG231641
AP-hu02 AP-hu02	000000002222222222222222222222222222222	9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3 4 5 6 7 8 9	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3 O3 O3 O4 O5 O6 O7 O8 O9	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1678256 814358 1680119 814651 1682318 2065463 307364 2067693 3073056 2090446 3078407 2094946 3083083 2108826	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor)	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA455022 AI142896 AA455022 AI079412 AA481094 AI066675 AI375553 BG231641 AI383639 BG231641 AI554887 BG231641 AI424264 BG231641 AI394321
AP-hu02 AP-hu02	22222222222222222222222222222222222222	9 10 11 12 13 14 15 16 17 18 19 20 21 22 24 1 23 4 5 6 7 8 9 10	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3 O4 O2 O3 O4 O5 O6 O7 O8 O9 O10	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814328 1680119 814651 1682318 2065463 3072364 2067693 3073056 2090446 3078407 2094946 3083083 2108826	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor)	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI075694 AA456021 AI075694 AA456702 AI142896 AA465233 AI066456 AA465233 AI066456 AA459022 AI079412 AA481094 AI0666755 AI375553 BG231641 AI383639 BG231641 AI32321641 AI3231641 AI3231641
AP-hu02 AP-hu0	000000000222222222222222222222222222222	9 10 11 12 13 14 15 16 17 18 19 20 21 22 24 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 7 18 19 20 21 21 21 21 21 21 21 21 21 21	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3 N24 O1 O2 O3 O4 O5 O6 O7 O8 O9 O10	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2065463 3072364 2067693 3073056 2090446 3078407 2094946 3083083 2108826 3085549	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor) 0	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456 AA465233 AI066456 AA465233 AI066456 AI375553 BG231641 AI383639 BG231641 AI424264 BG231641 AI394321 BG231641 AI394321 BG231641
AP-hu02 AP-hu02	000000000022222222222222222222222222222	5 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 23 24 10 11 12 13 14 15 16 17 18 19 20 21 23 24 15 16 17 18 19 20 21 23 24 12 13 14 15 16 17 18 19 20 21 22 23 24 12 13 14 15 16 17 18 19 20 21 22 23 24 12 13 12 13 14 15 16 17 18 19 20 12 12 13 14 15 16 17 18 19 20 12 12 12 12 12 12 12 12 12 12	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 O1 O2 O3 O4 O5 O6 O7 O8 O9 O10 O11	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2067693 3073056 2090446 3078407 2094946 3083083 2108826 3085549 2109792	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 chackpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor) 0 guanine nucleotide binding protein (G protein), alpha transducing activity	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA459022 AI079412 AA481094 AI079412 AA481094 AI079412 AA481094 AI079412 AA481094 AI079553 BG231641 AI383639 BG231641 AI424264 BG231641 AI394321 BG231641 AI394321 BG231641 AI493909
AP-hu02 AP-hu0	000000000052222222 22 22222	9 10 11 12 13 14 15 16 17 18 19 20 21 23 24 1 23 4 5 6 7 8 9 10 11 12 13 14 15 16 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 23 24 1 23 24 1 20 21 21 20 21 21 20 21 21 20 21 21 20 21 21 20 21 21 20 21 21 20 21 21 21 21 21 21 21 21 21 21	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3 O4 O2 O3 O4 O2 O3 O4 O5 O6 O7 O8 O9 O11	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2065463 3073056 2090446 3073056 2090446 3078407 2094946 3083083 2108826 3085549 2109792	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor) 0 guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456 AA465233 AI066456 AA459022 AI079412 AA481094 AI066675 AI375553 BG231641 AI383639 BG231641 AI383639 BG231641 AI394321 BG231641 AI394321 BG231641 AI394321 BG231641 AI3943909
AP-hu02 AP-hu02	0 0000000000022222222222222222222222222	9 10 11 12 13 14 15 16 17 18 19 20 21 22 24 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 7 18 19 20 21 21 22 24 1 2 2 2 2 2 2 2 2 2 2 2 2 2	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3 O4 O5 O6 O7 O8 O9 O10 O111 O12	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1678256 814358 1680119 814651 1682318 2065463 3072364 2065463 3073056 2090446 3078407 2094946 3083083 2108826 3085549 2109792 3085866	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor) 0 guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA455022 AI142896 AA459022 AI079412 AA481094 AI066675 AI37553 BG231641 AI354887 BG231641 AI394321 BG231641 AF493909 BG231641
AP-hu02 AP-hu0	00 000000000022222222222222222222222222	5 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 23 24 12 13 14 15 16 17 18 19 20 21 23 24 12 13 14 15 16 17 18 19 20 21 23 24 12 13 14 15 16 17 18 19 20 21 22 23 24 12 13 14 15 16 17 18 19 20 21 22 23 24 12 13 12 13 12 13 14 15 16 17 18 19 20 12 12 13 14 15 16 17 18 19 20 12 23 24 12 13 12 13 12 13 14 15 16 17 18 19 20 12 23 24 12 3 4 5 6 7 8 9 10 11 12 23 24 12 23 24 12 23 24 12 23 24 12 23 24 12 23 24 12 23 24 12 23 24 12 23 24 12 23 24 12 23 24 11 12 23 24 11 12 13 11 11 12 13 13 11 11 12 13	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N21 N23 N21 02 03 04 05 06 07 08 09 010 011 012 013	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814851 1682318 2067693 3073056 2090446 3078407 2094946 30783083 2108826 3085849 2109792	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor) 0 guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2 0 H2.0-like homeo box 1 (Drosophila)	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA456022 AI079412 AA481094 AI066456 AA459022 AI079412 AA481094 AI066675 BG231641 AI383639 BG231641 AI454887 BG231641 AI494321 AF493909 BG231641 AF493909
AP-hu02 AP-hu02	000 000000000002222222 22 22222	9 9 10 11 12 13 14 15 16 17 18 19 20 21 22 3 4 5 6 7 8 9 10 11 12 34 5 6 7 8 9 10	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3 O4 O5 O6 O7 O8 O9 O10 O11 O12 O13 O14	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2067693 3073056 2090446 3078407 2094946 3083083 2108826 3085549 2109792	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor) 0 guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2 0 H2.0-like homeo box 1 (Drosophila) early growth response 2 (Krox-20 homolog, Drosophila)	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI075694 AA455702 AI142896 AA455702 AI142896 AA465233 AI066456 AA459022 AI079412 AA481094 AI066675 AI375553 BG231641 AI383639 BG231641 AI394321 BG231641 AI394321 BG231641 AI493909 BG231641 AI468772 BG231641
AP-hu02 AP-hu02	0000 0000000000002222222222222222222222	6 9 10 11 12 13 14 15 16 17 18 19 20 21 22 24 1 2 3 4 5 6 7 8 9 10 11 12 13 4 5 6 7 8 9 10	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3 O4 O5 O6 O7 O8 O9 O10 O11 O12 O13 O145	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2065463 3073056 2090446 3078407 2094946 3083083 2108266 3085549 2109792 3085866 2125556 3087768 2130480	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor) 0 guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2 0 H2.0-like homeo box 1 (Drosophila) early growth response 2 (Krox-20 homolog, Drosophila) phosphoenoloyruvate carboxytimese 1 (soluble)	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA455022 AI142896 AA459022 AI079412 AA481094 AI066675 AI37553 BG231641 AI384321 BG231641 AI394321 BG231641 AI448772 BG231641 AI468772 BG231641 AI468772 BG231641
AP-hu02 AP-hu0	22222 22 222222222222222222222222222222	5 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 23 4 5 6 7 8 9 10 11 12 13 14 15 16 7 8 9 10 11 20 21 22 23 24 12 34 56 7 8 9 10 11 12 13	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N21 N23 N21 02 03 04 05 06 07 08 09 010 011 012 013 014 015	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814851 1682318 2065463 3072364 2067693 3073056 2090446 3078407 2094946 30783083 2108826 3085849 2109792 3085866 2125556 3087768 2130480	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor) 0 guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2 0 H2.0-like homeo box 1 (Drosophila) early growth response 2 (Krox-20 homolog, Drosophila) phosphoenolpyruvate carboxykinase 1 (soluble)	AA463987 AI076543 AA481753 AI076543 AA481753 AI078341 AA463470 AI075694 AA456021 AI075694 AA456021 AI075694 AA456702 AI142896 AA459022 AI079412 AA481094 AI066455 AI375553 BG231641 AI383639 BG231641 AI424264 BG231641 AI4493090 BG231641 AI468772 BG231641 AI464177 BG231641 AI464177
AP-hu02 AP-hu02	00000 000000000000000000000000000000000	8 9 10 11 12 13 14 15 16 7 18 19 20 21 22 24 1 23 4 5 6 7 8 9 10 11 12 34 5 6 7 8 9 10 11 12 13	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N3 N24 O1 O2 O3 O4 O5 O6 O7 O8 O9 O10 O11 O12 O13 O14 O15 O16	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1678249 814234 1679256 814358 1680119 814651 1682318 2065463 3072364 2067693 3073056 2090446 3078407 2094946 3083083 2108826 3085549 2109792 3085866 2125556 2085768 2130480 3643475	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor) 0 guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2 0 H2.0-like homeo box 1 (Drosophila) early growth response 2 (Krox-20 homolog, Drosophila) phosphoenolpyruvate carboxykinase 1 (soluble) v-abi Abelson murine leukemia viral oncogene homolog 2 (arg, Abelson-related cene)	AA463987 AI076543 AA481753 AI078541 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA465233 AI066456 AA459022 AI079412 AA481094 AI066675 AI375553 BG231641 AI383639 BG231641 AI384321 BG231641 AI394321 BG231641 AI468772 BG231641 AI494177 BG231641
AP-hu02 AP-hu0	222222 22 22222222222222222222222222222	5 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 12 3 4 5 6 7 8 9 10 11 12 13	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N20 N21 N22 N23 N24 O1 O2 O3 O4 O5 O6 O7 O8 O9 O10 O11 O12 O13 O14 O15 O16 O17	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814651 1682318 2065463 3072364 2067693 3073056 3073056 3073056 3078407 2094946 3083083 2108266 3085549 2109792 3085866 2125556 3087768 2130480 3643475 2124222	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor) 0 guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2 0 H2.0-like homeo box 1 (Drosophila) early growth response 2 (Krox-20 homolog, Drosophila) phosphoenolpyruvate carboxykinase 1 (soluble) v-abi Abelson murtine leukemia viral oncogene homolog 2 (arg, Abelson-related gene)	AA463987 AI076543 AA481753 AI076543 AA481753 AI078341 AA463470 AI075694 AA456021 AI078579 AA455702 AI142896 AA459022 AI079412 AA481094 AI079412 AA481094 AI079412 AA481094 AI079412 AA481094 AI079412 AA481094 AI079412 AA481094 AI079553 BG231641 AI394321 BG231641 AI468772 BG231641 AI468772 BG231641 AI468772
AP-hu02 AP-hu0	00 00000 000000000002222222222222222222	5 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 1 23 4 5 6 7 8 9 10 11 12 13 14 15 16 7 8 9 10 11 20 21 22 23 24 12 23 4 5 6 7 8 9 10 11 12 13	N8 N9 N10 N11 N12 N13 N14 N15 N16 N17 N18 N19 N21 N23 N24 O1 O2 O3 O4 O5 O6 O7 O8 O9 O10 O11 O12 O13 O14 O15 O16 O17	810672 1674873 810777 1675856 811810 1676286 812139 1677559 813991 1678249 814234 1679256 814358 1680119 814851 1682318 2065463 3072364 2067693 3073056 2090446 3078407 2094946 30783083 2108826 3085849 2109792 3085866 2125556 3087768 2130480 3643475 2131329	T-cell acute lymphocytic leukemia 1 dual specificity phosphatase 4 KIAA0217 protein TAF5-like RNA polymerase II, p300/CBP-associated factor (PCAF)-associated factor, 65kDa general transcription factor IIB aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase) chromosome 14 open reading frame 111 transcriptional activator of the c-fos promoter 0 succinate-CoA ligase, GDP-forming, beta subunit renal tumor antigen 0 interleukin 11 checkpoint suppressor 1 reticulon 4 collagen, type VI, alpha 2 zinc finger protein 147 (estrogen-responsive finger protein) p53-regulated apoptosis-inducing protein 1 0 polymerase (DNA directed), eta TGFB1-induced anti-apoptotic factor 1 RNA binding motif protein 5 0 interleukin 18 (interferon-gamma-inducing factor) 0 guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2 0 H2.0-like homeo box 1 (Drosophila) early growth response 2 (Krox-20 homolog, Drosophila) phosphoenolpyruvate carboxykinase 1 (solubie) v-abi Abelson murine leukemia viral oncogene homolog 2 (arg, Abelson-related gene) programmed cell death 8 (apoptosis-inducing factor)	AA463987 AI076543 AA481753 AI076543 AA481753 AI078341 AA463470 AI075694 AA456021 AI075694 AA455702 AI142896 AA455702 AI142896 AA459022 AI079412 AA481094 AI066455 AI375553 BG231641 AI383639 BG231641 AI424264 BG231641 AI494321 BG231641 AI494177 BG231641 AI494177 BG231641 AI494177

AP-hu02	0	19	O19	2139463	choline/ethanolaminephosphotransferase	AI470615
AP-hu02	0	20	O20	4142901	0	BG231641
AP-hu02	0	21	O21	2144779	WNT1 inducible signaling pathway protein 3	AI457678
AP-hu02	0	22	022	469577	ribonuclease, RNase A family, 4	AA027189
AP-hu02	0	23	023	2148030	Bcl-2-associated transcription factor	AI476761
AP-hu02	0	24	024	469931	32kDa	AAU29/00
AP-hu02	Р	1	P1	815040	signal peptidase complex (18kD)	AA465269
AP-hu02	Р	2	P2	1686798	sushi-repeat protein	AI089690
AP-hu02	Р	3	P3	815049	pyruvate dehydrogenase (lipoamide) beta	AA465285
AP-hu02	Р	4	P4	1688907	laminin, alpha 4	Al140443
AP-hu02	Р	5	P5	898925	inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase beta	AA480228
AP-hu02	Р	6	P6	1693375	homeo box A2	AI140882
AP-hu02	P	7	P7	926309	neurofibromin 1 (neurofibromatosis, von Recklinghausen disease, Watson disease)	AA534609
AP-hu02	P	8	P8	1698339	collagen, type XI, alpha 1	AIU966/1
AP-NUU2	2	9	P9 010	90008/	Zinc tinger protein, subtamily (A, 5 (Pegasus)	AA303960
AP-nuuz	P	10	PIU	1/02/44	1. 37kDa)	X1030304
AP-hu02	Р	11	P11	1030258	Gem-interacting protein	AA554534
AP-hu02	Р	12	P12	1702957	NICE-4 protein	AI097419
AP-hu02	Р	13	P13	1030484	translocated promoter region (to activated MET oncogene)	AA555031
AP-hu02	Р	14	P14	1706935	prefoldin 5	AI126765
AP-hu02	Р	15	P15	1031248	solute carrier family 30 (zinc transporter), member 4	AA609034
AP-hu02	Р	16	P16	1709428	Fc fragment of IgG, low affinity Illa, receptor for (CD16)	AI130777
AP-hu02	Р	17	P17	1032632	POU domain, class 4, transcription factor 1	AA779554
AP-hu02	P	18	P18	1710655	neuropilin 2	AI139365
AP-hu02	P	19	P19	1032647	endothelial differentiation, lysophosphatidic acid G-protein-coupled receptor, 4	AA//95/0
AP-nu02	Р	20	P20	1/12192	TAP1 KNA polymerase II, TATA box binding protein (TBP)-associated factor,	AI125912
AP-hu02	Р	21	P21	1056449	Notch homolog 3 (Drosophila)	AA557201
AP-hu02	P	22	P22	1715851	hemoglobin, gamma G	AI151352
AP-hu02	P	23	P23	1057103	human immunodeficiency virus type I enhancer binding protein 2	AA557381
AP-hu02	Ρ	24	P24	1725451	leukocyte-associated Ig-like receptor 2	AI189606
AP-hu03	A	1	A1	PK_A1	Human 90Kda heat shock protein	#N/A
AP-hu03	Α	2	A2	1737502	trinucleotide repeat containing 11 (THR-associated protein, 230kDa subunit)	AI138726
AP-hu03	Α	3	A3	PK_A2	60s Acidic Ribosomal Protein PO	#N/A
AP-hu03	Α	4	A4	1740083	thyroid transcription factor 1	AI187322
AP-hu03	Α	5	A5	PK_A4	nucleophosmin	#N/A
AP-hu03	A	6	A6	1740578	host cell factor C1 (VP16-accessory protein)	AI185261
AP-hu03	A	7	A7	PK_A6	heat shock cognate 70kDa protein	#N/A
AP-hu03	Â.	8	A8	1756042	modulator of apoptosis 1	AI203801
AP-NU03	Â	9	A9	PK_A10	NADH-Ubiquinone oxidoreductase SGDH subunit precusor	#NVA
AP-NUU3	A:	10	A10	1/09/10 DK A11	Linean HenC2 3' region cDNA	#N/A
	2	12	A12	1840595	homen hav 45	#IVA
AP-hu03	Â	13	A13	PK B1	Human triosephosphate isomerase	#N/A
AP-hu03	Â	14	A14	1841164	low density lipoprotein receptor-related protein 8, apolipoprotein e receptor	AI219647
AP-hu03	A	15	A15	PK B2	Human 3-hydroxy-3-methylglutavi coenzyme A reductase	#N/A
AP-hu03	Α	16	A16	1846264	myoglobin	AI239419
AP-hu03	Α	17	A17	PK_B3	Elongation factor 1 alpha subunit	#N/A
AP-hu03	Α	18	A18	1855723	ret proto-oncogene (multiple endocrine neoplasia and medullary thyroid carcinoma	AI263439
		40		DK 04	1, Hirschsprung disease)	461/4
AP-NUU3	Å	19	A19 A20	PK_B4	Stratagene tetal retina Willims tumor 1	#N/A
AP-1003	2	20	A20 A21	DK B5	nre mDNA enlicing factor SDD20	#N/A
AP-hu03	2	22	A22	1858266	pre linker spilling lactor SKrzu	#IN/A
AP-hu03	Â	23	A23	PK B6	Human acidic ribosomal phosphorotein PO (60s)	#N/A
AP-hu03	Ā	24	A24	1868332	cell growth regulatory with EF-hand domain	AI261384
AP-hu03	в	1	B1	0	0	#N/A
AP-hu03	в	2	B2	Ō	0	#N/A
AP-hu03	в	3	B3	0	0	#N/A
AP-hu03	в	4	B4	0	0	#N/A
AP-hu03	в	5	85	0	0	#N/A
AP-hu03	в	6	B6	0	0	#N/A
AP-hu03	в	7	B7	0	0	#N/A
AP-NU03	в	8	88	0	0	#N/A
AP-NU03	в	9	89	0	0	#N/A
AP-11003	B	11	B10 B11	0	0	#N/A #N/A
AP-hu03	B	12	812	0	0	#N/A #N/A
AP-hu03	B	13	B13	õ	0	#N/A
AP-hu03	в	14	P14	ō.	0	#N/A
AP-hu03			D14			
AP-hu03	В	15	B15	õ	0	#N/A
AP-hu03	BB	15 16	B15 B16	0	0	#N/A #N/A
	B B B	15 16 17	B14 B15 B16 B17	0 0 0	0 0 0	#N/A #N/A #N/A
AP-hu03	B B B B	15 16 17 18	B14 B15 B16 B17 B18	0 0 0 0	0 0 0 0	#N/A #N/A #N/A #N/A
AP-hu03 AP-hu03	B B B B	15 16 17 18 19	B14 B15 B16 B17 B18 B19	0 0 0 0 0		#N/A #N/A #N/A #N/A
AP-hu03 AP-hu03 AP-hu03	B B B B B B B B B B B B B B B B B B B	15 16 17 18 19 20	B14 B15 B16 B17 B18 B19 B20			#N/A #N/A #N/A #N/A #N/A
AP-hu03 AP-hu03 AP-hu03 AP-hu03	B B B B B B B B B B B B B B B B B B B	15 16 17 18 19 20 21	B14 B15 B16 B17 B18 B19 B20 B21		0 0 0 0 0 0	#N/A #N/A #N/A #N/A #N/A #N/A
AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03	B B B B B B B B B B B B B B B B B B B	15 16 17 18 19 20 21 22 22	B14 B15 B16 B17 B18 B19 B20 B21 B22 B22			#N/A #N/A #N/A #N/A #N/A #N/A #N/A
AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03	8 8 8 8 8 8 8 8 8 8 8 8 8 8	15 16 17 18 19 20 21 22 23 24	B14 B15 B16 B17 B18 B19 B20 B21 B22 B23 B24	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		#N/A #N/A #N/A #N/A #N/A #N/A #N/A #N/A
AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03	B B B B B B B B B B B C	15 16 17 18 19 20 21 22 23 24 1	B15 B15 B17 B18 B19 B20 B21 B22 B23 B24 C1	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	#N/A #N/A #N/A #N/A #N/A #N/A #N/A #N/A
AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03	B B B B B B B B B B B C C	15 16 17 18 20 21 22 23 24 1 2	B15 B15 B17 B18 B19 B20 B21 B22 B23 B24 C1 C2	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 EST 27474 (Brain-human) and alpha-fetoprotein enhancer binding protein microtubule-associated protein 4	#N/A #N/A #N/A #N/A #N/A #N/A #N/A #N/A
AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03	B B B B B B B B B B B C C C	15 16 17 18 20 21 22 23 24 1 2 3	B15 B16 B17 B18 B19 B20 B21 B22 B23 B24 C1 C2 C3	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 EST 27474 (Brain-human) and alpha-fetoprotein enhancer binding protein 0 EST 27474 (Brain-human) and alpha-fetoprotein enhancer binding protein Muman ribosomel protein 4 Human ribosomel protein 130	#N/A #N/A #N/A #N/A #N/A #N/A #N/A #N/A
AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03	B B B B B B B B B B B C C C C	15 16 17 18 19 20 21 22 23 24 1 2 3 4	B15 B16 B17 B18 B19 B20 B21 B22 B23 B24 C1 C2 C3 C4	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 EST 27474 (Brain-human) and alpha-fetoprotein enhancer binding protein 0 0 EST 27474 (Brain-human) and alpha-fetoprotein enhancer binding protein microtubule-associated protein 4 Human ribosomal protein L30 GS2 gene	#N/A #N/A #N/A #N/A #N/A #N/A #N/A #N/A
AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03	BBBBBBBBCCCCC	15 16 17 18 19 20 21 22 23 24 1 2 3 4 5	B15 B15 B17 B18 B19 B20 B21 B22 B23 B23 B24 C1 C2 C3 C4 C5	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	#N/A #N/A #N/A #N/A #N/A #N/A #N/A #N/A
AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03 AP-hu03	BBBBBBBBCCCCC	15 16 17 18 19 20 21 22 23 24 1 2 3 4 5 6	B15 B15 B17 B18 B19 B20 B21 B22 B23 B23 B24 C1 C2 C3 C4 C5 C6	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	#N/A #N/A #N/A #N/A #N/A #N/A #N/A #N/A

AP-hu03	с	8	C8	1929994	fatty acid binding protein 3, muscle and heart (mammary-derived growth inhibitor)	Al313221
AP-hu03	С	9	C9	PK_B11	3 LINE 1 elements	#N/A
AP-hu03	С	10	C10	1949396	thromboxane A synthase 1 (platelet, cytochrome P450, family 5, subfamily A)	AI342602
AP-hu03	c	11	C11	PK_B12	Trans-acting transcriptional protein ICPO	#N/A
AP-hu03	c	12	C12	2014456	prominin 1	AI362100
AP-NUU3	č	13	C13	2015313	adhesion alvooprotein	AI361991
AP-hu03	č	15	C15	PK C3	peptidyl-prolyl:cis-trans isomerase A	#N/A
AP-hu03	č	16	C16	2015920	natural killer cell receptor 2B4	AI363168
AP-hu03	С	17	C17	PK_C4	PK_C4	#N/A
AP-hu03	С	18	C18	2029834	general transcription factor IIH, polypeptide 3, 34kDa	AI373329
AP-hu03	C	19	C19	PK_C5	Eukaryotic initiation factor 4B (EIF-4B)	#N/A
AP-hu03	C	20	C20	2042756	laminin, alpha 3	AI3/5/8/
AP-hu03	6	21	C21 C22	PK_00	PK_C0	#INA
AP-hu03	č	22	C22	PK C8	Guanine nucleotide-binding protein beta subunit-like protein	#N/A
AP-hu03	č	24	C24	2066379	chloride channel 4	AI802379
AP-hu03	Ď	1	D1	0	0	#N/A
AP-hu03	D	2	D2	0	0	#N/A
AP-hu03	D	3	D3	0	0	#N/A
AP-hu03	D	4	D4	0	0	#N/A
AP-hu03	D	5	D5	0	0	#N/A
AP-NUU3	D	0 7	D6	0		#IN/A #NI/A
AP-hu03	D	8	D8	õ	0	#N/A
AP-hu03	D	9	D9	õ	0	#N/A
AP-hu03	Ð	10	D10	0	0	#N/A
AP-hu03	D	11	D11	0	0	#N/A
AP-hu03	D	12	D12	0	0	#N/A
AP-hu03	D	13	D13	0	0	#N/A
AP-hu03	D	14	D14	0	0	#N/A #N/A
AP-NUUS	D D	10	D15	0		#N/A #N/A
AP-hu03	D	17	D17	õ	0	#N/A
AP-hu03	D	18	D18	ō	0	#N/A
AP-hu03	D	19	D19	0	0	#N/A
AP-hu03	D	20	D20	0	0	#N/A
AP-hu03	D	21	D21	0	0	#N/A
AP-hu03	D	22	D22	0	0	#N/A
AP-hu03	D	23	D23	0	0	#N/A
	F	24	E1	PK C9	u elongation factor (vesst)	#Ν/Α #Ν/Δ
AP-hu03	E	2	E2	2067266	histone deacetvlase 8	AI803366
AP-hu03	Ē	3	E3	PK_C10	Guanine nucleotide-binding protein beta subunit-like protein	#N/A
AP-hu03	Ε	4	E4	2068299	adrenomedullin receptor	AI379917
AP-hu03	Е	5	E5	PK_C12	Soares pregant uterur NbHPU Homo sapien cDNA clone	#N/A
AP-hu03	Е	6	E6	2068829	joined to JAZF1	AI378171
AP-hu03	E	7	E7	PK_D2	Yeast Pep II protein	#N/A
	t c	8	E8	2072912	apoptosis antagonizing transcription factor	AI439571
AP-hu03	F	9 10	E9 E10	2091078	ras homolog gene family, member N	#INA
AP-hu03	E	11	E11	PK D5	PK D5	#N/A
AP-hu03	E	12	E12	2094592	activin A receptor type II-like 1	AI420822
AP-hu03	Е	13	E13	PK_D8	16s rRNA	#N/A
AP-hu03	Ε	14	E14	2095272	regulator of G-protein signalling 9	AI420112
AP-hu03	E	15	E15	PK_D10	Heat shock cognate 71kDa protein	#N/A
AP-hu03	E	16	E16	2096365	tumor protein p53 inducible nuclear protein 1	AI419377
AP-hu03	E E	1/	E1/	PK_D11	PK_D11 transmistion classocian factor A (SII) 1	#N/A
AP-11003	F	10	E10	2096303 PK F1	HIA class II histocompatibility antigen	#N/A
AP-hu03	Ē	20	E20	2108356	suppressor of Ty 6 homolog (S. cerevisiae)	AI392835
AP-hu03	E	21	E21	PK_E3	Soares parathyroid tumour NbHPA	#N/A
AP-hu03	Ε	22	E22	2108870	chemokine (C-X-C motif) ligand 6 (granulocyte chemotactic protein 2)	AI391519
AP-hu03	E	23	E23	PK_E6	Human ribosomal DNA	#N/A
AP-hu03	E	24	E24	2111581	mitogen-activated protein kinase 11	AI394426
AP-hu03	F	1	F1	0	0	#N/A
AP-NU03	F	2	F2 F2	0	0	#N/A
AP-hu03	F	4	F4	0	0	#Ν/Α #Ν/Δ
AP-hu03	F	5	F5	õ	0	#N/A
AP-hu03	F	6	F6	õ	0	#N/A
AP-hu03	F	7	F7	0	0	#N/A
AP-hu03	F	8	F8	0	0	#N/A
AP-hu03	F	9	F9	0	0	#N/A
AP-hu03	F	10	F10	0	0	#N/A
AP-0003	F	11	F11 F12	0	0	#NVA #N/A
AP-hu03	F	13	F12	0	0	#1¥Α #Ν/Δ
AP-hu03	F	14	F14	õ	0	#N/A
AP-hu03	F	15	F15	ō	0	#N/A
AP-hu03	F	16	F16	0	0	#N/A
AP-hu03	F	17	F17	0	0	#N/A
AP-hu03	F	18	F18	0	0	#N/A
AP-hu03	F	19	F19	0	0	#N/A
AP-0003	F	20	F20	0	0	#N/A #N/A
AP-hu03	F	22	F21	0	0	#IN/A #N/A
AP-hu03	F	23	F23	ō	0	#N/A

AP-hu03	F	24	F24	0	0	#N/A
AP-hu03	G	1	G1	PK_E9	Tubulin gaunachain	#N/A
AP-hu03	G	2	G2	2113938	amyloid beta (A4) precursor protein-binding, family B, member 2 (Fe65-like)	AI400246
AP-hu03	G	3	G3	PK_E11	alu subfamily	#N/A
AP-hu03	G	4	G4	2122503	leukotnene A4 hydrolase	AI521212
AP-NUU3	G	с С	GS	2137887	integrin, sinha 2 (CD49B, sinha 2 subunit of VI A-2 recentor)	#IWA AI538195
	G	7	G7	PK F1	Colon Carcinoma HCC	#N/A
AP-hu03	Ğ	8	G8	2138953	general transcription factor IIH, polypeptide 1, 62kDa	AI439100
AP-hu03	Ğ	9	G9	PK_F3	60s ribosomal protein L30	#N/A
AP-hu03	G	10	G10	2148566	0	AI469873
AP-hu03	G	11	G11	PK_F6	PK_F6	#N/A
AP-hu03	G	12	G12	2149893	Homo saplens transcribed sequence with moderate similarity to protein sp:Q92915 (H.sapiens) FGFE_HUMAN Fibroblast growth factor-14 (FGF-14) (Fibroblast growth factor homologous factor 4) (FHF-4)	AI887836
AP-hu03	G	13	G13	PK_F8	40s ribosomal protein s18	#N/A
AP-hu03	G	14	G14	2151139	programmed cell death 8 (apoptosis-inducing factor)	AI589001
AP-hu03	G	15	G15	PK_F10	CAMP-regulated phosphoprotein	#N/A
	G	10	G16 G17	2214009 PK F11	40e rihosomal protein s3A	#N/A
AP-hu03	Ğ	18	G18	2237098	low density lipoprotein receptor-related protein 6	AI633264
AP-hu03	Ğ	19	G19	empty	0	#N/A
AP-hu03	G	20	G20	2241908	thousand and one amino acid protein kinase	AI637830
AP-hu03	G	21	G21	empty	0	#N/A
AP-hu03	G	22	G22	2242601	pyruvate carboxylase	AI637857
AP-hu03	G	23	G23	empty		#N/A
AP-hu03	G	24	G24	2243192	transcriptional adaptor 2 (ADA2 homolog, yeast)-like	AI655771
AP-hu03		1	111	0		#IN/A #N/A
AP-hu03	н	3	H3	õ	0	#N/A
AP-hu03	н	4	H4	õ	0	#N/A
AP-hu03	Ĥ	5	H5	Ō	0	#N/A
AP-hu03	н	6	H6	0	0	#N/A
AP-hu03	н	7	H7	0	0	#N/A
AP-hu03	н	8	H8	0	0	#N/A
AP-hu03	н	9	H9	0	0	#N/A
AP-hu03	н	10	H10	0	0	#N/A #N/A
	н	12	H12	0	0	#Ν/Α #Ν/Δ
AP-hu03	н	13	H13	õ	0	#N/A
AP-hu03	н	14	H14	Ō	0	#N/A
AP-hu03	н	15	H15	0	0	#N/A
AP-hu03	H	16	H16	0	0	#N/A
AP-hu03	н	17	H17	0	0	#N/A
AP-hu03	н	18	H18	0	0	#N/A
AP-hu03	н	19	H19	0	0	#N/A
AP-NUU3	н	20	H20	0	0	#NVA #NI/A
AP-hu03	н	21	H22	0	0	#NVA #N/A
AP-hu03	н	23	H23	õ	0	#N/A
AP-hu03	н	24	H24	0	0	#N/A
AP-hu03	1	1	11	empty	0	#N/A
AP-hu03	I.	2	12	2243934	interleukin 21 receptor	AI656754
AP-hu03	1	3	13	empty	0	#N/A
AP-hu03	1	4	14	2244213	E3-binding protein	AI657202
AP-hu03	1	5	15	empty		#N/A
AP-NU03	1	5	10	2251/83	potassium voltage-gated channel, shaker-related subtamily, beta member 1	41628686
AP-hu03	÷	8	17	2252681	chemokine (C motif) ligand 1	#INA
AP-hu03	i	9	19	empty		#N/A
AP-hu03	1	10	110	2273173	wingless-type MMTV integration site family, member 4	AI634496
AP-hu03	I	11	111	empty	0	#N/A
AP-hu03	I	12	112	2284066	vanin 1	AI625852
AP-hu03	1	13	113	empty	0	#N/A
AP-hu03	1	14	114	2284081	stanniocalcin 1	AI625879
AP-0003		15	115	empty 2284472	U zina finaer protein 443	#N/A
	-	10	110	2204473	2inc inger protein 443	41013104 #NI/A
AP-hu03	i	18	118	2304672	v telomerase reverse transcriptase	#INA
AP-hu03	i	19	119	empty	0	#N/A
AP-hu03	1	20	120	2305123	Homo sapiens transcribed sequences	AI631732
AP-hu03	ł	21	121	empty	0	#N/A
AP-hu03	I	22	122	2316826	cytochrome P450, family 3, subfamily A, polypeptide 4	AI671993
AP-hu03	1	23	123	empty	0	#N/A
AP-hu03	-	24	124	2321108	pasic leucine zipper nuclear factor 1 (JEM-1)	AI675660
AP-0003	J	1	J] CI	0	0	#N/A #N/A
	J	2	.12	õ	ů.	#1%/A #N/A
AP-hu03	Ĵ	4	.14	õ	<u>0</u>	#N/A
AP-hu03	Ĵ	5	J5	0	0	#N/A
AP-hu03	J	6	J6	0	0	#N/A
AP-hu03	J	7	J7	0	0	#N/A
AP-hu03	J	8	3L	0	0	#N/A
AP-hu03	J	9	J9	0	0	#N/A
AP-hu03	J	10	J10	0	0	#N/A
AP-hu03	J	11	J11	0	U	#N/A
AP-0003	J	12	J12	0		#N/A
m -1003	~	13	113	v	v	#11//1

AP-hu03	J	14	J14	0	0	#N/A
AP-hu03	Ĵ	15	J15	ō	0	#N/A
AP-hu03	J	16	J16	0	0	#N/A
AP-hu03	J	17	J17	0	0	#N/A
AP-hu03	J	18	J18	0	0	#N/A
AP-hu03	J	19	J19	0	0	#N/A
AP-hu03	J	20	J20	0	0	#N/A
AP-hu03	J	21	J21	0	0	#N/A
AP-hu03	J	22	J22	0	0	#N/A
AP-hu03	J	23	J23	0	0	#N/A
AP-hu03	J	24	J24	0	0	#N/A #N/A
AP-hu03	ĸ	1	K1 K0	empty	U BAD54 like 2 (S. comulcies)	#19/4
AP-NU03	ĸ	2	K2	2330252	RADSTHIKE S (S. Cerevisiae)	#N/A
AP-0003	R K	3	KA	authors	U transient recentor potential cation channel, subfamily V, member 6	#WA52307
AP-11003	ĸ	4	K5	empty		#N/A
AP-11003	ĸ	6	KG	3070183	RNA binding motif protein 5	BE513550
	ĸ	7	K7	empty		#N/A
AP-hu03	ĸ	8	K8	3071696	SWI/SNF related, matrix associated, actin dependent regulator of chromatin.	BF513730
		•			subfamily a, member 3	
AP-hu03	к	9	K9	empty	0	#N/A
AP-hu03	к	10	K10	2731796	TAF4b RNA polymerase II, TATA box binding protein (TBP)-associated factor,	AW297262
	~		1444		105kDa	
AP-hu03	ĸ	11	K11	empty	U Name continue transactional and upperson	#INA
AP-NUUS	È.	12	K12	2/3/9/3		#N/A
AP-11003	ĸ	14	K13	3173557	v regulator of G-protein signalling 4	#INA BE222672
AP-1003	ĸ	15	K15	empty		#N/A
	ĸ	16	K16	3178227	programmed cell death 6	BE219939
	ĸ	17	K17	emoty		#N/A
AP-hu03	ĸ	18	K18	3183031	lipoprotein lipase	BE465407
AP-hu03	ĸ	19	K19	empty	0	#N/A
AP-hu03	ĸ	20	K20	3184548	epidermal growth factor (beta-urogastrone)	BE501422
AP-hu03	ĸ	21	K21	empty	0	#N/A
AP-hu03	к	22	K22	3196945	aminoadipate-semialdehyde synthase	BE466144
AP-hu03	к	23	K23	empty	0	#N/A
AP-hu03	κ	24	K24	3209795	adrenergic, beta, receptor kinase 2	BE504864
AP-hu03	L	1	L1	0	0	#N/A
AP-hu03	L	2	L2	0	0	#N/A
AP-hu03	L	3	L3	0	0	#N/A
AP-hu03	L	4	L4	0	0	#N/A
AP-hu03	L	5	L5	0	0	#N/A
AP-hu03	L	6	L6	0	0	#N/A
AP-hu03	L	7	L7	0	0	#N/A
AP-hu03	L	8	1.8	0	0	#N/A
AP-NU03	L	9	L9	0	0	#N/A
AP-NUU3	L	10	L10	0	0	#NVA
AP-11003		12	112	0	0	#N/A
AP-hu03	i i	13	113	0	0	#N/A
	1	14	1 14	ŏ	ů O	#N/A
AP-hu03	ĩ	15	1 15	õ	ů O	#N/A
AP-hu03	Ē	16	L16	õ	0	#N/A
AP-hu03	Ē	17	L17	õ	0	#N/A
AP-hu03	Ē	18	L18	ō	0	#N/A
AP-hu03	L	19	L19	0	0	#N/A
AP-hu03	L	20	L20	0	0	#N/A
AP-hu03	L	21	L21	0	0	#N/A
AP-hu03	L	22	L22	0	0	#N/A
AP-hu03	L	23	L23	0	0	#N/A
AP-hu03	L	24	L24	0	0	#N/A
AP-hu03	м	1	M1	empty	0	#N/A
AP-hu03	м	2	M2	3213061	19A24 protein	BE467670
AP-hu03	M	3	M3	empty		#N/A
AP-hu03	M	4	M4	3218773	nypotnetical protein FLJ20793	BE503286
AP-hu03	M	5	M5	empty	0 One De M (musice) e estes de la tracta de la t	#N/A
AP-NUU3	M	0	MO	32201/3	cas-pr-m (munne) ecotropic retroviral transforming sequence	BE050582
AP-11003	MI MA		M/	SUBLE	U Homo canions transcribed eccuences	#INA
AP-NUU3	M	8	MB	3220758		BE6/3055
	M	10	M9	3220042	u insulinoma associated 1	#INVA
	M	11	M10	empty	0	#N/A
AP-hu03	M	12	M12	3286412	complement component 1 a subcomponent recentor 1	RE671226
AP-bu03	M	13	M13	empty		#N/A
AP-hu03	м	14	M14	3316202	- protein tyrosine phosphatase, receptor type, H	BF000072
AP-hu03	M	15	M15	empty	0	#N/A
AP-hu03	M	16	M16	3316686	glycogen synthase 1 (muscle)	BF002321
AP-hu03	м	17	M17	empty	0	#N/A
AP-hu03	M	18	M18	3318799	MARCKS-like protein	BF591250
AP-hu03	м	19	M19	empty	0	#N/A
AP-hu03	м	20	M20	3320058	cytochrome P450, family 3, subfamily A, polypeptide 5	BF591425
AP-hu03	м	21	M21	empty	0	#N/A
AP-hu03	м	22	M22	3393054	retinoblastoma binding protein 5	BF055070
AP-hu03	М	23	M23	empty	0	#N/A
AP-hu03	м	24	M24	3405658	transforming growth factor, beta receptor I (activin A receptor type II-like kinase,	BF439832
	N	4	A14	^	53kDa)	
AP-0003	N N	1	Dia 19	0		#N/A #N/A
	N	2	NZ N2	0	0	#N/A #N/A
AP-11003	IN .	3	145	v	v	#N/A

AP-hu03	N	4	N4	0	0	#N/A
AP-hu03	Ν	5	N5	0	0	#N/A
AP-hu03	Ν	6	N6	0	0	#N/A
AP-hu03	Ν	7	N7	0	0	#N/A
AP-hu03	Ν	8	N8	0	0	#N/A
AP-hu03	Ν	9	N9	0	0	#N/A
AP-hu03	N	10	N10	0	0	#N/A
AP-hu03	N	11	N11	0	0	#N/A
AP-hu03	N	12	N12	0	0	#IN/A #NI/A
AP-hu03	N	13	N13	0	0	#NVA #NI/A
AP-NU03	N	14	N14	0	0	#N/A
AP-NUU3	N	10	N15 N16	0	0	#N/A
AP-1003	N N	17	N17	0	0	#N/A
AP-hu03	N	18	N18	õ	0	#N/A
AP-hu03	N	19	N19	õ	0	#N/A
AP-hu03	N	20	N20	0	0	#N/A
AP-hu03	N	21	N21	ō	0	#N/A
AP-hu03	N	22	N22	0	0	#N/A
AP-hu03	Ν	23	N23	0	0	#N/A
AP-hu03	Ν	24	N24	0	0	#N/A
AP-hu03	0	1	01	empty	0	#N/A
AP-hu03	0	2	02	3476767	arylsulfatase B	BF058213
AP-hu03	0	3	03	empty	0	#N/A
AP-hu03	0	4	04	3477610	Wilms tumor associated protein	BF059076
AP-hu03	0	5	05	empty		#N/A
AP-hu03	0	6	06	3523521	integrin, beta 6	BF111900
AP-hu03	0		07	empty	U aastala teasaast asatala SEOC1 alaba aubusit isafaan 1	#NVA
AP-NU03	0	8	08	3525/40		#N/A
AP-NUU3	Ň	9 10	09	900pty 3562165	U postmeiotic segregation increased 2-like 6	#INA
AP-1003	õ	11	010	emoty		#N/A
AP-hu03	õ	12	012	3562195	V YY1 associated factor 2	BE197072
AP-hu03	ŏ	13	012	empty	0	#N/A
AP-hu03	ŏ	14	014	3085383	zinc finger protein 256	BF510334
AP-hu03	ō	15	015	empty	0	#N/A
AP-hu03	ō	16	016	3085523	spinocerebellar ataxia 7 (olivopontocerebellar atrophy with retinal degeneration)	BF510366
AP-hu03	0	17	017	empty	0	#N/A
AP-hu03	0	18	O18	3565961	chemokine (C-C motif) ligand 22	BF432854
AP-hu03	0	19	O19	empty	0	#N/A
AP-hu03	0	20	O20	3570409	B cell RAG associated protein	BF115957
AP-hu03	0	21	O21	empty	0	#N/A
AP-hu03	0	22	022	empty	0	#N/A
AP-hu03	0	23	023	empty	0	#N/A
AP-hu03	0	24	024	empty	0	#N/A
AP-hu03	2	1	P1	0	0	#N/A #N/A
	Р В	2	P2 D2	0	0	#N/A #N/A
AP-1003	r D	3	F3 P4	ñ	0	#N/A
AP-hu03	P	5	P5	õ	ů	#N/A
AP-hu03	P	6	P6	õ	0	#N/A
AP-hu03	P	7	P7	ō	0	#N/A
AP-hu03	Р	8	P8	0	0	#N/A
AP-hu03	Ρ	9	P9	0	0	#N/A
AP-hu03	Р	10	P10	0	0	#N/A
AP-hu03	Р	11	P11	0	0	#N/A
AP-hu03	Ρ	12	P12	0	0	#N/A
AP-hu03	Р	13	P13	0	0	#N/A
AP-hu03	P	14	P14	0	0	#N/A
AP-hu03	2	15	P15	0	0	#N/A
AP-NUU3	2	16	P16	0	0	#NVA
AP-0003	۲ P	17	P1/	0	0	#NVA #N1/A
AP-0003	Þ	10	P10	0	0	#N/A #N/A
AP-hu03	p	20	P20	õ	0	#N/A
AP-hu03	P	21	P21	õ	ů.	#N/A
AP-hu03	P	22	P22	õ	ō	#N/A
AP-hu03	P	23	P23	ō	Ō	#N/A
AP-hu03	Р	24	P24	0	0	#N/A

Appendix C - Sequence alignments of RT-PCR amplified splice variants of HHV-7

>U17 top band

```
EM VI:AF037218 AF037218.1 Human herpesvirus 7 strain RK, complete genome.
      Length = 153.080
 Plus Strand HSPs:
Score = 650 (103.6 bits), Expect = 3.9e-21, P = 3.9e-21
 Identities = 208/280 (74%), Positives = 208/280 (74%), Strand = Plus / Plus
         2 TTGCA-CATAGGAGATAGCAACCTATGAATTGA-CAAACACTTTCTATAACATCTTGACA 59
Ouerv:
Sbjct: 27747 TTGCAACATAGGAGATAGCAACCTATGAAAACAACAACAACATATTAGAACATCATGACA 27806
        60 CGGTGGAAATGGTTTGACTGCTTTTGTA-CTT-CATACTCTCTTGCCA-TCT-TATGAAG 115
Query:
Sbjct: 27807 CATTGGAAATGTTTTGACTGCTTTTGCAACTTACATAGG-TCTTGCAGCTCGGTTTCACT 27865
Query:
        116 ATTTCACACCTGATATGTTGATCCACG-AAAGGCCTAATAAAATTTTTCTTAAT--G-GG 171

      Image: Sbjct:
      27866 AGT--AGTTCTGAAAGGCATATCC-CGCAAAGGCGAAATGATTATTTTCCTACTCTGAGG 27922

        172 TTGTCTAACAAGGATTAAATTTCCCCAGTGTATGCCTAATAACGAAGTCACTAAGAGAATC 231
Query:
232 GCGATAAAAGGCTTGTTTAACAAGGCTTAAATTTGCGAGT 271
Query:
           Sbjct: 27980 GCGATAAAAGGCTTGTTTAACAAGGCTTAAATTTGCGAGT 28019
 Score = 349 (58.4 bits), Expect = 1.5e-07, P = 1.5e-07
 Identities = 99/131 (75%), Positives = 99/131 (75%), Strand = Plus / Plus
         89 TTCATACTCTCTTGCCATCTTATGAAGATTTCACACCTGATATGTTGATCCACGAAAGGC 148
Ouerv:
            Sbjct: 27909 TTCCTACTCTGAGGCCATCTTATGAGAATTTCACACCCACTATTTTGAGTAACGAA-GTC 27967
        149 - CTAATAAAATTTTTCTTAATGGGTTGTCTAACAAGGATTAAATTTCCCCAGTGTATGCCT 207
Query:
208 AATAACGAAGT 218
Query:
            Sbjct: 28028 AA-ATCGAACT 28037
```

>U17 mid band

```
>EM_VI:AF037218 AF037218.1 Human herpesvirus 7 strain RK, complete genome.
      Length = 153,080
 Plus Strand HSPs:
Score = 1010 (157.6 bits), Expect = 2.1e-37, P = 2.1e-37
 Identities = 202/202 (100%), Positives = 202/202 (100%), Strand = Plus / Plus
Ouery:
         1 AGAACATCATGACACATTGGAAATGTTTTGACTGCTTTTGCAACTTACATAGGTCTTGCA 60
           Sbjct: 27793 AGAACATCATGACACATTGGAAATGTTTTGACTGCTTTTGCAACTTACATAGGTCTTGCA 27852
        61 GCTCGGTTTCACTAGTAGTTCTGAAAGGCATATCCCGCAAAGGCGAAATGATTATTTTCC 120
Query:
           Sbjct: 27853 GCTCGGTTTCACTAGTAGTTCTGAAAGGCATATCCCGCAAAGGCGAAATGATTATTTCC 27912
       121 TACTCTGAGGCCATCTTATGAAGATTTCACACCCACTATTTTGAGTAACGAAGTCACTAA 180
Ouery:
Sbjct: 27913 TACTCTGAGGCCATCTTATGAAGATTTCACACCCACTATTTTGAGTAACGAAGTCACTAA 27972
        181 GAGAATCGCGATAAAAGGCTTG 202
Ouery:
           Sbjct: 27973 GAGAATCGCGATAAAAGGCTTG 27994
```

>U17 bottom band

```
EM VI:AF037218 AF037218.1 Human herpesvirus 7 strain RK, complete genome.
      Length = 153,080
 Plus Strand HSPs:
Score = 870 (136.6 bits), Expect = 4.5e-31, P = 4.5e-31
Identities = 174/174 (100%), Positives = 174/174 (100%), Strand = Plus / Plus
        15 ACATAGGTCTTGCAGGTCCGGTTTCACTAGTAGTTCTGAAAGGCATATCCCGCAAAGGCGA 74
Ouerv:
          Sbjct: 27839 ACATAGGTCTTGCAGGTCCGGTTTCACTAGTAGTTCTGAAAGGCATATCCCCGCAAAGGCGA 27898
        75 AATGATTATTTTCCTACTCTGAGGCCATCTTATGAAGATTTCACACCCACTATTTTGAGT 134
Ouery:
          Sbjct: 27899 AATGATTATTTTCCTACTCTGAGGCCATCTTATGAAGATTTCACACCCACTATTTTGAGT 27958
       135 AACGAAGTCACTAAGAGAATCGCGATAAAAGGCTTGTTTAACAAGGCTTAAATT 188
Ouery:
          Sbjct: 27959 AACGAAGTCACTAAGAGAATCGCGATAAAAGGCTTGTTTAACAAGGCTTAAATT 28012
```

>U90

>U41 top band

>CL6 = U41 bottom band

```
EM_VI:AF037218 AF037218.1 Human herpesvirus 7 strain RK, complete genome.
    Length = 153,080
 Plus Strand HSPs:
Score = 885 (138.8 bits), Expect = 9.5e-32, P = 9.5e-32
Identities = 177/177 (100%), Positives = 177/177 (100%), Strand = Plus / Plus
      Query:
       Ouerv:
      61 TTTTAACGGTTGGGAAGAAGTCATTTTCAACCGTTAGATTAAGTAACAAGGGTGAGATCA 120
       Sbjct: 66427 TTTTAACGGTTGGGAAGAAGTCATTTTCAACCGTTAGATTAAGTAACAAGGGTGAGATCA 66486
Query:
     121 CTACTGATTTTCTTTTTTCCATTAACGATAATGTAGTCAGAATTTCTATTAATTTTT 177
```

REFERENCES

Ablashi, D. V., Handy, M., Bernbaum, J., Chatlynne, L. G., Lapps, W., Kramarsky, B., Berneman, Z. N., Komaroff, A. L., and Whitman, J. E. (1998). Propagation and characterization of human herpesvirus-7 (HHV-7) isolates in a continuous T-lymphoblastoid cell line (SupT1). J Virol Methods 73, 123-140.

Agaisse, H., Burrack, L. S., Philips, J. A., Rubin, E. J., Perrimon, N., and Higgins, D. E. (2005). Genome-wide RNAi screen for host factors required for intracellular bacterial infection. Science 309, 1248-1251.

Ahn, J. H., Jang, W. J., and Hayward, G. S. (1999). The human cytomegalovirus IE2 and UL112-113 proteins accumulate in viral DNA replication compartments that initiate from the periphery of promyelocytic leukemia protein-associated nuclear bodies (PODs or ND10). J Virol 73, 10458-10471.

Ahn, J. W., Powell, K. L., Kellam, P., and Alber, D. G. (2002). Gammaherpesvirus lytic gene expression as characterized by DNA array. J Virol 76, 6244-6256.

Alba, M. M., Lee, D., Pearl, F. M., Shepherd, A. J., Martin, N., Orengo, C. A., and Kellam, P. (2001). VIDA: a virus database system for the organization of animal virus genome open reading frames. Nucleic Acids Res 29, 133-136.

Alsaker, K. V., and Papoutsakis, E. T. (2005). Transcriptional program of early sporulation and stationary-phase events in Clostridium acetobutylicum. J Bacteriol 187, 7103-7118.

Altschul, S. F., Bundschuh, R., Olsen, R., and Hwa, T. (2001). The estimation of statistical parameters for local alignment score distributions. Nucleic Acids Res 29, 351-361.

Bahnson, A. B., Dunigan, J. T., Baysal, B. E., Mohney, T., Atchison, R. W., Nimgaonkar, M. T., Ball, E. D., and Barranger, J. A. (1995). Centrifugal enhancement of retroviral mediated gene transfer. J Virol Methods 54, 131-143.

Baldick, C. J., Jr., Marchini, A., Patterson, C. E., and Shenk, T. (1997). Human cytomegalovirus tegument protein pp71 (ppUL82) enhances the infectivity of viral DNA and accelerates the infectious cycle. J Virol 71, 4400-4408.

Bassett, D. E., Jr., Eisen, M. B., and Boguski, M. S. (1999). Gene expression informatics--it's all in your mine. Nat Genet 21, 51-55.

Behr, M. A., Wilson, M. A., Gill, W. P., Salamon, H., Schoolnik, G. K., Rane, S., and Small, P. M. (1999). Comparative genomics of BCG vaccines by whole-genome DNA microarray. Science 284, 1520-1523.

Ben Mamoun, C., Gluzman, I. Y., Hott, C., MacMillan, S. K., Amarakone, A. S., Anderson, D. L., Carlton, J. M., Dame, J. B., Chakrabarti, D., Martin, R. K., *et al.* (2001). Co-ordinated programme of gene expression during asexual intraerythrocytic development of the human malaria parasite Plasmodium falciparum revealed by microarray analysis. Mol Microbiol *39*, 26-36.

Berneman, Z. N., Ablashi, D. V., Li, G., Eger-Fletcher, M., Reitz, M. S., Jr., Hung, C. L., Brus, I., Komaroff, A. L., and Gallo, R. C. (1992a). Human herpesvirus 7 is a T-lymphotropic virus and is

related to, but significantly different from, human herpesvirus 6 and human cytomegalovirus. Proc Natl Acad Sci U S A 89, 10552-10556.

Berneman, Z. N., Gallo, R. C., Ablashi, D. V., Frenkel, N., Katsafanas, G., Kramarsky, B., and Brus, I. (1992b). Human herpesvirus 7 (HHV-7) strain JI: independent confirmation of HHV-7. J Infect Dis 166, 690-691.

Bigger, C. B., Brasky, K. M., and Lanford, R. E. (2001). DNA microarray analysis of chimpanzee liver during acute resolving hepatitis C virus infection. J Virol 75, 7059-7066.

Birnboim, H. C., and Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. Nucleic Acids Res 7, 1513-1523.

Black, J. B., Burns, D. A., Goldsmith, C. S., Feorino, P. M., Kite-Powell, K., Schinazi, R. F., Krug, P. W., and Pellett, P. E. (1997). Biologic properties of human herpesvirus 7 strain SB. Virus Res 52, 25-41.

Black, J. B., and Pellett, P. E. (1999). Human herpesvirus 7. Rev Med Virol 9, 245-262.

Black, J. B., Sanderlin, K. C., Goldsmith, C. S., Gary, H. E., Lopez, C., and Pellett, P. E. (1989). Growth properties of human herpesvirus-6 strain Z29. J Virol Methods 26, 133-145.

Bolt, G., Berg, K., and Blixenkrone-Moller, M. (2002). Measles virus-induced modulation of host-cell gene expression. J Gen Virol 83, 1157-1165.

Brazma, A., Hingamp, P., Quackenbush, J., Sherlock, G., Spellman, P., Stoeckert, C., Aach, J., Ansorge, W., Ball, C. A., Causton, H. C., *et al.* (2001). Minimum information about a microarray experiment (MIAME)-toward standards for microarray data. Nat Genet 29, 365-371.

Brennan, D. C., Storch, G. A., Singer, G. G., Lee, L., Rueda, J., and Schnitzler, M. A. (2000). The prevalence of human herpesvirus-7 in renal transplant recipients is unaffected by oral or intravenous ganciclovir. J Infect Dis 181, 1557-1561.

Browne, E. P., Wing, B., Coleman, D., and Shenk, T. (2001). Altered cellular mRNA levels in human cytomegalovirus-infected fibroblasts: viral block to the accumulation of antiviral mRNAs. J Virol 75, 12319-12330.

Broyles, S. S. (2003). Vaccinia virus transcription. J Gen Virol 84, 2293-2303.

Cai, H. Y., Lu, L., Muckle, C. A., Prescott, J. F., and Chen, S. (2005). Development of a novel protein microarray method for serotyping Salmonella enterica strains. J Clin Microbiol 43, 3427-3430.

Campadelli-Fiume, G., Cocchi, F., Menotti, L., and Lopez, M. (2000). The novel receptors that mediate the entry of herpes simplex viruses and animal alphaherpesviruses into cells. Rev Med Virol 10, 305-319.

Carter, K. L., Cahir-McFarland, E., and Kieff, E. (2002). Epstein-barr virus-induced changes in B-lymphocyte gene expression. J Virol 76, 10427-10436.

Chambers, J., Angulo, A., Amaratunga, D., Guo, H., Jiang, Y., Wan, J. S., Bittner, A., Frueh, K., Jackson, M. R., Peterson, P. A., *et al.* (1999). DNA microarrays of the complex human cytomegalovirus genome: profiling kinetic class with drug sensitivity of viral gene expression. J Virol 73, 5757-5766.

Chan, P. K., Peiris, J. S., Yuen, K. Y., Liang, R. H., Lau, Y. L., Chen, F. E., Lo, S. K., Cheung, C. Y., Chan, T. K., and Ng, M. H. (1997). Human herpesvirus-6 and human herpesvirus-7 infections in bone marrow transplant recipients. J Med Virol 53, 295-305.

Chang, Y. E., and Laimins, L. A. (2000). Microarray analysis identifies interferon-inducible genes and Stat-1 as major transcriptional targets of human papillomavirus type 31. J Virol 74, 4174-4182.

Che, S., and Ginsberg, S. D. (2004). Amplification of RNA transcripts using terminal continuation. Lab Invest 84, 131-137.

Chee, M. S., Bankier, A. T., Beck, S., Bohni, R., Brown, C. M., Cerny, R., Horsnell, T., Hutchison, C. A., 3rd, Kouzarides, T., Martignetti, J. A., and et al. (1990). Analysis of the protein-coding content of the sequence of human cytomegalovirus strain AD169. Curr Top Microbiol Immunol 154, 125-169.

Chen, Y., Kamat, V., Dougherty, E. R., Bittner, M. L., Meltzer, P. S., and Trent, J. M. (2002). Ratio statistics of gene expression levels and applications to microarray data analysis. Bioinformatics 18, 1207-1215.

Chicurel, M. (2002). Bioinformatics: bringing it all together. Nature 419, 751, 753, 755 passim.

Chuck, A. S., Clarke, M. F., and Palsson, B. O. (1996). Retroviral infection is limited by Brownian motion. Hum Gene Ther 7, 1527-1534.

Churchill, G. A. (2004). Using ANOVA to analyze microarray data. Biotechniques 37, 173-175, 177.

Clark, D. A., Freeland, M. L., Mackie, L. K., Jarrett, R. F., and Onions, D. E. (1993). Prevalence of antibody to human herpesvirus 7 by age. J Infect Dis 168, 251-252.

Compton, T., Nowlin, D. M., and Cooper, N. R. (1993). Initiation of human cytomegalovirus infection requires initial interaction with cell surface heparan sulfate. Virology 193, 834-841.

Corbeil, J., Sheeter, D., Genini, D., Rought, S., Leoni, L., Du, P., Ferguson, M., Masys, D. R., Welsh, J. B., Fink, J. L., *et al.* (2001). Temporal gene regulation during HIV-1 infection of human CD4+ T cells. Genome Res 11, 1198-1204.

Cossman, J. (2001). Gene expression analysis of single neoplastic cells and the pathogenesis of Hodgkin's lymphoma. J Histochem Cytochem 49, 799-800.

Cotter, P. A., and Miller, J. F. (1998). In vivo and ex vivo regulation of bacterial virulence gene expression. Curr Opin Microbiol 1, 17-26.

Covell, D. G., Wallqvist, A., Rabow, A. A., and Thanki, N. (2003). Molecular classification of cancer: unsupervised self-organizing map analysis of gene expression microarray data. Mol Cancer Ther 2, 317-332.

Crowley, R. W., Secchiero, P., Zella, D., Cara, A., Gallo, R. C., and Lusso, P. (1996). Interference between human herpesvirus 7 and HIV-1 in mononuclear phagocytes. J Immunol 156, 2004-2008.

Cuadras, M. A., Feigelstock, D. A., An, S., and Greenberg, H. B. (2002). Gene expression pattern in Caco-2 cells following rotavirus infection. J Virol 76, 4467-4482.

Cummings, C. A., and Relman, D. A. (2000). Using DNA microarrays to study host-microbe interactions. Emerg Infect Dis 6, 513-525.

Cuomo, L., Trivedi, P., de Grazia, U., Calogero, A., D'Onofrio, M., Yang, W., Frati, L., Faggioni, A., Rymo, L., and Ragona, G. (1998). Upregulation of Epstein-Barr virus-encoded latent membrane protein by human herpesvirus 6 superinfection of EBV-carrying Burkitt lymphoma cells. J Med Virol 55, 219-226.

Dana, A. N., Hong, Y. S., Kern, M. K., Hillenmeyer, M. E., Harker, B. W., Lobo, N. F., Hogan, J. R., Romans, P., and Collins, F. H. (2005). Gene expression patterns associated with blood-feeding in the malaria mosquito Anopheles gambiae. BMC Genomics 6, 5.

Das, S., Sierra, J. C., Soman, K. V., Suarez, G., Mohammad, A. A., Dang, T. A., Luxon, B. A., and Reyes, V. E. (2005). Differential protein expression profiles of gastric epithelial cells following Helicobacter pylori infection using ProteinChips. J Proteome Res 4, 920-930.

de Brevern, A. G., Hazout, S., and Malpertuy, A. (2004). Influence of microarrays experiments missing values on the stability of gene groups by hierarchical clustering. BMC Bioinformatics 5, 114.

de La Fuente, C., Deng, L., Santiago, F., Arce, L., Wang, L., and Kashanchi, F. (2000). Gene expression array of HTLV type 1-infected T cells: Up-regulation of transcription factors and cell cycle genes. AIDS Res Hum Retroviruses 16, 1695-1700.

de la Fuente, C., Santiago, F., Deng, L., Eadie, C., Zilberman, I., Kehn, K., Maddukuri, A., Baylor, S., Wu, K., Lee, C. G., *et al.* (2002). Gene expression profile of HIV-1 Tat expressing cells: a close interplay between proliferative and differentiation signals. BMC Biochem 3, 14.

de Saizieu, A., Gardes, C., Flint, N., Wagner, C., Kamber, M., Mitchell, T. J., Keck, W., Amrein, K. E., and Lange, R. (2000). Microarray-based identification of a novel Streptococcus pneumoniae regulon controlled by an autoinduced peptide. J Bacteriol *182*, 4696-4703.

DeRisi, J., Penland, L., Brown, P. O., Bittner, M. L., Meltzer, P. S., Ray, M., Chen, Y., Su, Y. A., and Trent, J. M. (1996). Use of a cDNA microarray to analyse gene expression patterns in human cancer. Nat Genet 14, 457-460.

DeRisi, J. L., Iyer, V. R., and Brown, P. O. (1997). Exploring the metabolic and genetic control of gene expression on a genomic scale. Science 278, 680-686.

Diehn, M., and Relman, D. A. (2001). Comparing functional genomic datasets: lessons from DNA microarray analyses of host-pathogen interactions. Curr Opin Microbiol 4, 95-101.

Domachowske, J. B., Bonville, C. A., Easton, A. J., and Rosenberg, H. F. (2002). Differential expression of proinflammatory cytokine genes in vivo in response to pathogenic and nonpathogenic pneumovirus infections. J Infect Dis *186*, 8-14.

Drago, F., Ranieri, E., Malaguti, F., Losi, E., and Rebora, A. (1997). Human herpesvirus 7 in pityriasis rosea. Lancet 349, 1367-1368.

Easton, A. J., and Clements, J. B. (1980). Temporal regulation of herpes simplex virus type 2 transcription and characterization of virus immediate early mRNA's. Nucleic Acids Res 8, 2627-2645.

Ebrahimi, B., Dutia, B. M., Roberts, K. L., Garcia-Ramirez, J. J., Dickinson, P., Stewart, J. P., Ghazal, P., Roy, D. J., and Nash, A. A. (2003). Transcriptome profile of murine gammaherpesvirus-68 lytic infection. J Gen Virol 84, 99-109.

Eckmann, L., Smith, J. R., Housley, M. P., Dwinell, M. B., and Kagnoff, M. F. (2000). Analysis by high density cDNA arrays of altered gene expression in human intestinal epithelial cells in response to infection with the invasive enteric bacteria Salmonella. J Biol Chem 275, 14084-14094.

Eisen, M. B., Spellman, P. T., Brown, P. O., and Botstein, D. (1998). Cluster analysis and display of genome-wide expression patterns. Proc Natl Acad Sci U S A 95, 14863-14868.

Fakhari, F. D., and Dittmer, D. P. (2002). Charting latency transcripts in Kaposi's sarcoma-associated herpesvirus by whole-genome real-time quantitative PCR. J Virol 76, 6213-6223.

Fernandez, C., Boutolleau, D., Manichanh, C., Mangeney, N., Agut, H., and Gautheret-Dejean, A. (2002). Quantitation of HHV-7 genome by real-time polymerase chain reaction assay using MGB probe technology. J Virol Methods 106, 11-16.

Fielden, M. R., Halgren, R. G., Dere, E., and Zacharewski, T. R. (2002). GP3: GenePix postprocessing program for automated analysis of raw microarray data. Bioinformatics 18, 771-773.

Florens, L., Washburn, M. P., Raine, J. D., Anthony, R. M., Grainger, M., Haynes, J. D., Moch, J. K., Muster, N., Sacci, J. B., Tabb, D. L., *et al.* (2002). A proteomic view of the Plasmodium falciparum life cycle. Nature 419, 520-526.

Frenkel, N., Schirmer, E. C., Wyatt, L. S., Katsafanas, G., Roffman, E., Danovich, R. M., and June, C. H. (1990). Isolation of a new herpesvirus from human CD4+ T cells. Proc Natl Acad Sci U S A 87, 748-752.

Frohman, M. A., Dush, M. K., and Martin, G. R. (1988). Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer. Proc Natl Acad Sci U S A 85, 8998-9002.

Fujisaki, H., Tanaka-Taya, K., Tanabe, H., Hara, T., Miyoshi, H., Okada, S., and Yamanishi, K. (1998). Detection of human herpesvirus 7 (HHV-7) DNA in breast milk by polymerase chain reaction and prevalence of HHV-7 antibody in breast-fed and bottle-fed children. J Med Virol 56, 275-279.

Furukawa, M., Yasukawa, M., Yakushijin, Y., and Fujita, S. (1994). Distinct effects of human herpesvirus 6 and human herpesvirus 7 on surface molecule expression and function of CD4+ T cells. J Immunol 152, 5768-5775.

Geballe, A. P., Leach, F. S., and Mocarski, E. S. (1986). Regulation of cytomegalovirus late gene expression: gamma genes are controlled by posttranscriptional events. J Virol 57, 864-874.

Geiss, G. K., An, M. C., Bumgarner, R. E., Hammersmark, E., Cunningham, D., and Katze, M. G. (2001). Global impact of influenza virus on cellular pathways is mediated by both replication-dependent and -independent events. J Virol 75, 4321-4331.

Geiss, G. K., Bumgarner, R. E., An, M. C., Agy, M. B., van 't Wout, A. B., Hammersmark, E., Carter, V. S., Upchurch, D., Mullins, J. I., and Katze, M. G. (2000). Large-scale monitoring of host cell gene expression during HIV-1 infection using cDNA microarrays. Virology 266, 8-16.

Geiss, G. K., Carter, V. S., He, Y., Kwieciszewski, B. K., Holzman, T., Korth, M. J., Lazaro, C. A., Fausto, N., Bumgarner, R. E., and Katze, M. G. (2003). Gene expression profiling of the cellular transcriptional network regulated by alpha/beta interferon and its partial attenuation by the hepatitis C virus nonstructural 5A protein. J Virol 77, 6367-6375.

Geng, Y. Q., Chandran, B., Josephs, S. F., and Wood, C. (1992). Identification and characterization of a human herpesvirus 6 gene segment that trans activates the human immunodeficiency virus type 1 promoter. J Virol 66, 1564-1570.

Gerhold, D., Lu, M., Xu, J., Austin, C., Caskey, C. T., and Rushmore, T. (2001). Monitoring expression of genes involved in drug metabolism and toxicology using DNA microarrays. Physiol Genomics 5, 161-170.

Gibson, W. (1983). Protein counterparts of human and simian cytomegaloviruses. Virology 128, 391-406.

Goebel, S. J., Johnson, G. P., Perkus, M. E., Davis, S. W., Winslow, J. P., and Paoletti, E. (1990). The complete DNA sequence of vaccinia virus. Virology 179, 247-266, 517-263.

Goff, L. A., Bowers, J., Schwalm, J., Howerton, K., Getts, R. C., and Hart, R. P. (2004). Evaluation of sense-strand mRNA amplification by comparative quantitative PCR. BMC Genomics 5, 76.

Gompels, U. A., Nicholas, J., Lawrence, G., Jones, M., Thomson, B. J., Martin, M. E., Efstathiou, S., Craxton, M., and Macaulay, H. A. (1995). The DNA sequence of human herpesvirus-6: structure, coding content, and genome evolution. Virology 209, 29-51.

Graveel, C. R., Jatkoe, T., Madore, S. J., Holt, A. L., and Farnham, P. J. (2001). Expression profiling and identification of novel genes in hepatocellular carcinomas. Oncogene 20, 2704-2712.

Gray, N. S., Wodicka, L., Thunnissen, A. M., Norman, T. C., Kwon, S., Espinoza, F. H., Morgan, D. O., Barnes, G., LeClerc, S., Meijer, L., *et al.* (1998). Exploiting chemical libraries, structure, and genomics in the search for kinase inhibitors. Science 281, 533-538.

Guerra, S., Lopez-Fernandez, L. A., Conde, R., Pascual-Montano, A., Harshman, K., and Esteban, M. (2004). Microarray analysis reveals characteristic changes of host cell gene expression in response to attenuated modified vaccinia virus Ankara infection of human HeLa cells. J Virol 78, 5820-5834.

Guerra, S., Lopez-Fernandez, L. A., Pascual-Montano, A., Munoz, M., Harshman, K., and Esteban, M. (2003). Cellular gene expression survey of vaccinia virus infection of human HeLa cells. J Virol 77, 6493-6506.

Hacia, J. G. (1999). Resequencing and mutational analysis using oligonucleotide microarrays. Nat Genet 21, 42-47.

Hakenbeck, R., Balmelle, N., Weber, B., Gardes, C., Keck, W., and de Saizieu, A. (2001). Mosaic genes and mosaic chromosomes: intra- and interspecies genomic variation of Streptococcus pneumoniae. Infect Immun 69, 2477-2486.

Hannon, G. J. (2002). RNA interference. Nature 418, 244-251.

Harrington, C. A., Rosenow, C., and Retief, J. (2000). Monitoring gene expression using DNA microarrays. Curr Opin Microbiol 3, 285-291.

Harris, T. M., and Childs, G. (2002). Global gene expression patterns during differentiation of F9 embryonal carcinoma cells into parietal endoderm. Funct Integr Genomics 2, 105-119.

Hauser, M. A., Li, Y. J., Xu, H., Noureddine, M. A., Shao, Y. S., Gullans, S. R., Scherzer, C. R., Jensen, R. V., McLaurin, A. C., Gibson, J. R., et al. (2005). Expression profiling of substantia nigra in

Parkinson disease, progressive supranuclear palsy, and frontotemporal dementia with parkinsonism. Arch Neurol 62, 917-921.

Hayashi, M. L., Blankenship, C., and Shenk, T. (2000). Human cytomegalovirus UL69 protein is required for efficient accumulation of infected cells in the G1 phase of the cell cycle. Proc Natl Acad Sci U S A 97, 2692-2696.

Hayward, R. E. (2000). Plasmodium falciparum phosphoenolpyruvate carboxykinase is developmentally regulated in gametocytes. Mol Biochem Parasitol 107, 227-240.

Hedenfalk, I., Duggan, D., Chen, Y., Radmacher, M., Bittner, M., Simon, R., Meltzer, P., Gusterson, B., Esteller, M., Kallioniemi, O. P., *et al.* (2001). Gene-expression profiles in hereditary breast cancer. N Engl J Med 344, 539-548.

Hegde, P., Qi, R., Abernathy, K., Gay, C., Dharap, S., Gaspard, R., Hughes, J. E., Snesrud, E., Lee, N., and Quackenbush, J. (2000). A concise guide to cDNA microarray analysis. Biotechniques 29, 548-550, 552-544, 556 passim.

Heller, M. J. (2002). DNA microarray technology: devices, systems, and applications. Annu Rev Biomed Eng 4, 129-153.

Helmberg, A. (2001). DNA-microarrays: novel techniques to study aging and guide gerontologic medicine. Exp Gerontol 36, 1189-1198.

Hertel, L., and Mocarski, E. S. (2004). Global Analysis of Host Cell Gene Expression Late during Cytomegalovirus Infection Reveals Extensive Dysregulation of Cell Cycle Gene Expression and Induction of Pseudomitosis Independent of US28 Function. J Virol 78, 11988-12011.

Hinchliffe, S. J., Isherwood, K. E., Stabler, R. A., Prentice, M. B., Rakin, A., Nichols, R. A., Oyston, P. C., Hinds, J., Titball, R. W., and Wren, B. W. (2003). Application of DNA microarrays to study the evolutionary genomics of Yersinia pestis and Yersinia pseudotuberculosis. Genome Res 13, 2018-2029.

Ho, W. Z., Cherukuri, R., Ge, S. D., Cutilli, J. R., Song, L., Whitko, S., and Douglas, S. D. (1993). Centrifugal enhancement of human immunodeficiency virus type 1 infection and human cytomegalovirus gene expression in human primary monocyte/macrophages in vitro. J Leukoc Biol 53, 208-212.

Hobbs, W. E., 2nd, and DeLuca, N. A. (1999). Perturbation of cell cycle progression and cellular gene expression as a function of herpes simplex virus ICP0. J Virol 73, 8245-8255.

Hodgkin, P. D., Scalzo, A. A., Swaminathan, N., Price, P., and Shellam, G. R. (1988). Murine cytomegalovirus binds reversibly to mouse embryo fibroblasts: implications for quantitation and explanation of centrifugal enhancement. J Virol Methods 22, 215-230.

Honda, M., Kaneko, S., Kawai, H., Shirota, Y., and Kobayashi, K. (2001). Differential gene expression between chronic hepatitis B and C hepatic lesion. Gastroenterology 120, 955-966.

Honda, M., Kawai, H., Shirota, Y., Yamashita, T., and Kaneko, S. (2005). Differential gene expression profiles in stage I primary biliary cirrhosis. Am J Gastroenterol 100, 2019-2030.

Honess, R. W., and Roizman, B. (1974). Regulation of herpesvirus macromolecular synthesis. I. Cascade regulation of the synthesis of three groups of viral proteins. J Virol 14, 8-19.

Honess, R. W., and Watson, D. H. (1977). Herpes simplex virus resistance and sensitivity to phosphonoacetic acid. J Virol 21, 584-600.

Huang, S., and Hanson, L. A. (1998). Temporal gene regulation of the channel catfish virus (Ictalurid herpesvirus 1). J Virol 72, 1910-1917.

Hudson, J. B. (1988). Further studies on the mechanism of centrifugal enhancement of cytomegalovirus infectivity. J Virol Methods 19, 97-108.

Hudson, J. B., Misra, V., and Mosmann, T. R. (1976). Cytomegalovirus infectivity: analysis of the phenomenon of centrifugal enhancement of infectivity. Virology 72, 235-243.

Hummerich, L., Muller, R., Hess, J., Kokocinski, F., Hahn, M., Furstenberger, G., Mauch, C., Lichter, P., and Angel, P. (2005). Identification of novel tumour-associated genes differentially expressed in the process of squamous cell cancer development. Oncogene.

Iizuka, N., Oka, M., Yamada-Okabe, H., Mori, N., Tamesa, T., Okada, T., Takemoto, N., Sakamoto, K., Hamada, K., Ishitsuka, H., *et al.* (2005). Self-organizing-map-based molecular signature representing the development of hepatocellular carcinoma. FEBS Lett 579, 1089-1100.

Imanishi, T., Itoh, T., Suzuki, Y., O'Donovan, C., Fukuchi, S., Koyanagi, K. O., Barrero, R. A., Tamura, T., Yamaguchi-Kabata, Y., Tanino, M., *et al.* (2004). Integrative annotation of 21,037 human genes validated by full-length cDNA clones. PLoS Biol 2, e162.

Inoue, N., Winter, J., Lal, R. B., Offermann, M. K., and Koyano, S. (2003). Characterization of entry mechanisms of human herpesvirus 8 by using an Rta-dependent reporter cell line. J Virol 77, 8147-8152.

Irizarry, R. A., Warren, D., Spencer, F., Kim, I. F., Biswal, S., Frank, B. C., Gabrielson, E., Garcia, J. G., Geoghegan, J., Germino, G., *et al.* (2005). Multiple-laboratory comparison of microarray platforms. Nat Methods 2, 345-350.

Iskenderian, A. C., Huang, L., Reilly, A., Stenberg, R. M., and Anders, D. G. (1996). Four of eleven loci required for transient complementation of human cytomegalovirus DNA replication cooperate to activate expression of replication genes. J Virol 70, 383-392.

Itoh, K., Kawasaki, S., Kawamoto, S., Seishima, M., Chiba, H., Michibata, H., Wakimoto, K., Imai, Y., Minesaki, Y., Otsuji, M., and Okubo, K. (2005). Identification of differentially expressed genes in psoriasis using expression profiling approaches. Exp Dermatol 14, 667-674.

Jenner, R. G., Alba, M. M., Boshoff, C., and Kellam, P. (2001). Kaposi's sarcoma-associated herpesvirus latent and lytic gene expression as revealed by DNA arrays. J Virol 75, 891-902.

Jenner, R. G., and Young, R. A. (2005). Insights into host responses against pathogens from transcriptional profiling. Nat Rev Microbiol 3, 281-294.

Johannes, G., Carter, M. S., Eisen, M. B., Brown, P. O., and Sarnow, P. (1999). Identification of eukaryotic mRNAs that are translated at reduced cap binding complex eIF4F concentrations using a cDNA microarray. Proc Natl Acad Sci U S A 96, 13118-13123.

Jones, J. O., and Arvin, A. M. (2003). Microarray analysis of host cell gene transcription in response to varicella-zoster virus infection of human T cells and fibroblasts in vitro and SCIDhu skin xenografts in vivo. J Virol 77, 1268-1280.

Kane, M. D., Jatkoe, T. A., Stumpf, C. R., Lu, J., Thomas, J. D., and Madore, S. J. (2000). Assessment of the sensitivity and specificity of oligonucleotide (50mer) microarrays. Nucleic Acids Res 28, 4552-4557.

Kato-Maeda, M., Rhee, J. T., Gingeras, T. R., Salamon, H., Drenkow, J., Smittipat, N., and Small, P. M. (2001). Comparing genomes within the species Mycobacterium tuberculosis. Genome Res 11, 547-554.

Keil, G. M., Ebeling-Keil, A., and Koszinowski, U. H. (1984). Temporal regulation of murine cytomegalovirus transcription and mapping of viral RNA synthesized at immediate early times after infection. J Virol 50, 784-795.

Kellam, P. (2001). Post-genomic virology: the impact of bioinformatics, microarrays and proteomics on investigating host and pathogen interactions. Rev Med Virol 11, 313-329.

Kempf, W., Adams, V., Kleinhans, M., Burg, G., Panizzon, R. G., Campadelli-Fiume, G., and Nestle, F. O. (1999). Pityriasis rosea is not associated with human herpesvirus 7. Arch Dermatol 135, 1070-1072.

Kempf, W., Adams, V., Mirandola, P., Menotti, L., Di Luca, D., Wey, N., Muller, B., and Campadelli-Fiume, G. (1998). Persistence of human herpesvirus 7 in normal tissues detected by expression of a structural antigen. J Infect Dis 178, 841-845.

Kennedy, P. G., Grinfeld, E., Craigon, M., Vierlinger, K., Roy, D., Forster, T., and Ghazal, P. (2005). Transcriptomal analysis of varicella-zoster virus infection using long oligonucleotide-based microarrays. J Gen Virol *86*, 2673-2684.

Kerr, M. K., Martin, M., and Churchill, G. A. (2000). Analysis of variance for gene expression microarray data. J Comput Biol 7, 819-837.

Khodarev, N. N., Advani, S. J., Gupta, N., Roizman, B., and Weichselbaum, R. R. (1999). Accumulation of specific RNAs encoding transcriptional factors and stress response proteins against a background of severe depletion of cellular RNAs in cells infected with herpes simplex virus 1. Proc Natl Acad Sci U S A 96, 12062-12067.

Kitahara, O., Furukawa, Y., Tanaka, T., Kihara, C., Ono, K., Yanagawa, R., Nita, M. E., Takagi, T., Nakamura, Y., and Tsunoda, T. (2001). Alterations of gene expression during colorectal carcinogenesis revealed by cDNA microarrays after laser-capture microdissection of tumor tissues and normal epithelia. Cancer Res *61*, 3544-3549.

Kotani, H., Newton, P. B., 3rd, Zhang, S., Chiang, Y. L., Otto, E., Weaver, L., Blaese, R. M., Anderson, W. F., and McGarrity, G. J. (1994). Improved methods of retroviral vector transduction and production for gene therapy. Hum Gene Ther 5, 19-28.

Kothapalli, R., Yoder, S. J., Mane, S., and Loughran, T. P., Jr. (2002). Microarray results: how accurate are they? BMC Bioinformatics 3, 22.

Kuwabara, P. E., and Coulson, A. (2000). RNAi--prospects for a general technique for determining gene function. Parasitol Today 16, 347-349.

LaBaer, J., and Ramachandran, N. (2005). Protein microarrays as tools for functional proteomics. Curr Opin Chem Biol 9, 14-19. Landazuri, N., and Le Doux, J. M. (2004). Complexation of retroviruses with charged polymers enhances gene transfer by increasing the rate that viruses are delivered to cells. J Gene Med 6, 1304-1319.

Larkin, J. E., Frank, B. C., Gavras, H., Sultana, R., and Quackenbush, J. (2005). Independence and reproducibility across microarray platforms. Nat Methods 2, 337-344.

Larsson, O., Wahlestedt, C., and Timmons, J. A. (2005). Considerations when using the significance analysis of microarrays (SAM) algorithm. BMC Bioinformatics δ , 129.

Leong, W. F., Tan, H. C., Ooi, E. E., Koh, D. R., and Chow, V. T. (2005). Microarray and real-time RT-PCR analyses of differential human gene expression patterns induced by severe acute respiratory syndrome (SARS) coronavirus infection of Vero cells. Microbes Infect 7, 248-259.

Levesque, K., Finzi, A., Binette, J., and Cohen, E. A. (2004). Role of CD4 receptor down-regulation during HIV-1 infection. Curr HIV Res 2, 51-59.

Lieberman, J., Song, E., Lee, S. K., and Shankar, P. (2003). Interfering with disease: opportunities and roadblocks to harnessing RNA interference. Trends Mol Med 9, 397-403.

Little, S. J., Riggs, N. L., Chowers, M. Y., Fitch, N. J., Richman, D. D., Spina, C. A., and Guatelli, J. C. (1994). Cell surface CD4 downregulation and resistance to superinfection induced by a defective provirus of HIV-1. Virology 205, 578-582.

Liu, B., and Stinski, M. F. (1992). Human cytomegalovirus contains a tegument protein that enhances transcription from promoters with upstream ATF and AP-1 cis-acting elements. J Virol 66, 4434-4444.

Lockhart, D. J., Dong, H., Byrne, M. C., Follettie, M. T., Gallo, M. V., Chee, M. S., Mittmann, M., Wang, C., Kobayashi, M., Horton, H., and Brown, E. L. (1996). Expression monitoring by hybridization to high-density oligonucleotide arrays. Nat Biotechnol 14, 1675-1680.

Lockhart, D. J., and Winzeler, E. A. (2000). Genomics, gene expression and DNA arrays. Nature 405, 827-836.

Lopetegui, P., Campo-Vera, H., and Yamanishi, K. (1985). Varicella-zoster virus (VZV)-specific polypeptides detected in cells treated with metabolic inhibitors. Microbiol Immunol 29, 569-575.

Lundquist, C. A., Tobiume, M., Zhou, J., Unutmaz, D., and Aiken, C. (2002). Nef-mediated downregulation of CD4 enhances human immunodeficiency virus type 1 replication in primary T lymphocytes. J Virol 76, 4625-4633.

Luo, L., Salunga, R. C., Guo, H., Bittner, A., Joy, K. C., Galindo, J. E., Xiao, H., Rogers, K. E., Wan, J. S., Jackson, M. R., and Erlander, M. G. (1999). Gene expression profiles of laser-captured adjacent neuronal subtypes. Nat Med 5, 117-122.

Lusso, P., Secchiero, P., Crowley, R. W., Garzino-Demo, A., Berneman, Z. N., and Gallo, R. C. (1994). CD4 is a critical component of the receptor for human herpesvirus 7: interference with human immunodeficiency virus. Proc Natl Acad Sci U S A 91, 3872-3876.

MacFarlane, R., Bhattacharya, D., and Singh, U. (2005a). Genomic DNA microarrays for Entamoeba histolytica: applications for use in expression profiling and strain genotyping. Exp Parasitol 110, 196-202.

MacFarlane, R. C., Shah, P. H., and Singh, U. (2005b). Transcriptional profiling of Entamoeba histolytica trophozoites. Int J Parasitol 35, 533-542.

Marko, N. F., Frank, B., Quackenbush, J., and Lee, N. H. (2005). A robust method for the amplification of RNA in the sense orientation. BMC Genomics 6, 27.

Mayne, M., Cheadle, C., Soldan, S. S., Cermelli, C., Yamano, Y., Akhyani, N., Nagel, J. E., Taub, D. D., Becker, K. G., and Jacobson, S. (2001). Gene expression profile of herpesvirus-infected T cells obtained using immunomicroarrays: induction of proinflammatory mechanisms. J Virol 75, 11641-11650.

McClintick, J. N., Jerome, R. E., Nicholson, C. R., Crabb, D. W., and Edenberg, H. J. (2003). Reproducibility of oligonucleotide arrays using small samples. BMC Genomics 4, 4.

McPherson, J. D., Marra, M., Hillier, L., Waterston, R. H., Chinwalla, A., Wallis, J., Sekhon, M., Wylie, K., Mardis, E. R., Wilson, R. K., *et al.* (2001). A physical map of the human genome. Nature 409, 934-941.

Megaw, A. G., Rapaport, D., Avidor, B., Frenkel, N., and Davison, A. J. (1998). The DNA sequence of the RK strain of human herpesvirus 7. Virology 244, 119-132.

Mendez, J. C., Dockrell, D. H., Espy, M. J., Smith, T. F., Wilson, J. A., Harmsen, W. S., Ilstrup, D., and Paya, C. V. (2001). Human beta-herpesvirus interactions in solid organ transplant recipients. J Infect Dis 183, 179-184.

Menegazzi, P., Galvan, M., Rotola, A., Ravaioli, T., Gonelli, A., Cassai, E., and Di Luca, D. (1999). Temporal mapping of transcripts in human herpesvirus-7. J Gen Virol 80 (Pt 10), 2705-2712.

Merrell, D. S., Butler, S. M., Qadri, F., Dolganov, N. A., Alam, A., Cohen, M. B., Calderwood, S. B., Schoolnik, G. K., and Camilli, A. (2002). Host-induced epidemic spread of the cholera bacterium. Nature 417, 642-645.

Miller, J. C., Zhou, H., Kwekel, J., Cavallo, R., Burke, J., Butler, E. B., Teh, B. S., and Haab, B. B. (2003). Antibody microarray profiling of human prostate cancer sera: antibody screening and identification of potential biomarkers. Proteomics 3, 56-63.

Mirandola, P., Menegazzi, P., Merighi, S., Ravaioli, T., Cassai, E., and Di Luca, D. (1998). Temporal mapping of transcripts in herpesvirus 6 variants. J Virol 72, 3837-3844.

Mitchell, R., Chiang, C. Y., Berry, C., and Bushman, F. (2003). Global analysis of cellular transcription following infection with an HIV-based vector. Mol Ther 8, 674-687.

Mocarski, E. S., Courcelle, C.T (2001). Cytomegaloviruses and their replication. In Fields Virology, D. M. Knipe, Howley, P. M., ed. (Philadelphia, Lippincott Williams & Wilkins), pp. 2629-2674.

Morgan, F. W., Stewart, J. A., Smith, A. N., and Tarnuzzer, R. W. (2005). Differential expression of stress response genes in the H-Tx rat model of congenital hydrocephalus. Brain Res Mol Brain Res 138, 273-290.

Morgan, R. W., Sofer, L., Anderson, A. S., Bernberg, E. L., Cui, J., and Burnside, J. (2001). Induction of host gene expression following infection of chicken embryo fibroblasts with oncogenic Marek's disease virus. J Virol 75, 533-539.

Moses, A. V., Jarvis, M. A., Raggo, C., Bell, Y. C., Ruhl, R., Luukkonen, B. G., Griffith, D. J., Wait, C. L., Druker, B. J., Heinrich, M. C., et al. (2002). A functional genomics approach to Kaposi's sarcoma. Ann N Y Acad Sci 975, 180-191.

Mossman, K. L., Macgregor, P. F., Rozmus, J. J., Goryachev, A. B., Edwards, A. M., and Smiley, J. R. (2001). Herpes simplex virus triggers and then disarms a host antiviral response. J Virol 75, 750-758.

Murphy, N., Millar, E., and Lee, C. S. (2005). Gene expression profiling in breast cancer: towards individualising patient management. Pathology 37, 271-277.

Nam, D. K., Lee, S., Zhou, G., Cao, X., Wang, C., Clark, T., Chen, J., Rowley, J. D., and Wang, S. M. (2002). Oligo(dT) primer generates a high frequency of truncated cDNAs through internal poly(A) priming during reverse transcription. Proc Natl Acad Sci U S A 99, 6152-6156.

Nicholas, J. (1996). Determination and analysis of the complete nucleotide sequence of human herpesvirus 7. J Virol 70, 5975-5989.

Nyberg-Hoffman, C., Shabram, P., Li, W., Giroux, D., and Aguilar-Cordova, E. (1997). Sensitivity and reproducibility in adenoviral infectious titer determination. Nat Med 3, 808-811.

O'Doherty, U., Swiggard, W. J., and Malim, M. H. (2000). Human immunodeficiency virus type 1 spinoculation enhances infection through virus binding. J Virol 74, 10074-10080.

Ohyashiki, J. H., Takaku, T., Ojima, T., Abe, K., Yamamoto, K., Zhang, Y., and Ohyashiki, K. (2005). Transcriptional profiling of human herpesvirus type B (HHV-6B) in an adult T cell leukemia cell line as in vitro model for persistent infection. Biochem Biophys Res Commun 329, 11-17.

Okabe, H., Satoh, S., Kato, T., Kitahara, O., Yanagawa, R., Yamaoka, Y., Tsunoda, T., Furukawa, Y., and Nakamura, Y. (2001). Genome-wide analysis of gene expression in human hepatocellular carcinomas using cDNA microarray: identification of genes involved in viral carcinogenesis and tumor progression. Cancer Res *61*, 2129-2137.

Okada, Y., Sahara, T., Mitsubayashi, H., Ohgiya, S., and Nagashima, T. (2005). Knowledge-assisted recognition of cluster boundaries in gene expression data. Artif Intell Med 35, 171-183.

Okamoto, T., Suzuki, T., and Yamamoto, N. (2000). Microarray fabrication with covalent attachment of DNA using bubble jet technology. Nat Biotechnol 18, 438-441.

Ordway, J. M., Fenster, S. D., Ruan, H., and Curran, T. (2005). A transcriptome map of cellular transformation by the fos oncogene. Mol Cancer 4, 19.

Osborn, J. E., and Walker, D. L. (1968). Enhancement of infectivity of murine cytomegalovirus in vitro by centrifugal inoculation. J Virol 2, 853-858.

Osman, H. K., Peiris, J. S., Taylor, C. E., Warwicker, P., Jarrett, R. F., and Madeley, C. R. (1996). "Cytomegalovirus disease" in renal allograft recipients: is human herpesvirus 7 a co-factor for disease progression? J Med Virol 48, 295-301.

Pan, L. Z., Werner, A., and Levy, J. A. (1993). Detection of plasma viremia in human immunodeficiency virus-infected individuals at all clinical stages. J Clin Microbiol 31, 283-288.

Pandey, A., and Mann, M. (2000). Proteomics to study genes and genomes. Nature 405, 837-846.

Pandya, U., Allen, C. A., Watson, D. A., and Niesel, D. W. (2005). Global profiling of Streptococcus pneumoniae gene expression at different growth temperatures. Gene 360, 45-54.

Pang, A. L., Johnson, W., Ravindranath, N., Dym, M., Rennert, O. M., and Chan, W. Y. (2005). Expression Profiling of Purified Male Germ Cells: Stage Specific Expression Patterns Related to Meiosis and Post-Meiotic Development. Physiol Genomics.

Pari, G. S., Kacica, M. A., and Anders, D. G. (1993). Open reading frames UL44, IRS1/TRS1, and UL36-38 are required for transient complementation of human cytomegalovirus oriLyt-dependent DNA synthesis. J Virol 67, 2575-2582.

Paulose-Murphy, M., Ha, N. K., Xiang, C., Chen, Y., Gillim, L., Yarchoan, R., Meltzer, P., Bittner, M., Trent, J., and Zeichner, S. (2001). Transcription program of human herpesvirus 8 (kaposi's sarcoma-associated herpesvirus). J Virol 75, 4843-4853.

Pease, A. C., Solas, D., Sullivan, E. J., Cronin, M. T., Holmes, C. P., and Fodor, S. P. (1994). Lightgenerated oligonucleotide arrays for rapid DNA sequence analysis. Proc Natl Acad Sci U S A 91, 5022-5026.

Peden, K. W. C., Martin, M. A. (1995). Virological and molecular genetic techniques for studies of established HIV isolates. In HIV: A Practical Approach - Virology & Immunology, J. Karn, ed. (New York, Oxford University Press), pp. 21-46.

Pellett, P. E., Dominguez, G. (2001). Human herpesviruses 6A, 6A, and 7 and their replication. In Fields Virology, D. M. Knipe, Howley, P. M., ed. (Philadelphia, Lippincott Williams & Wilkins), pp. 2769-2784.

Penfold, M. E., and Mocarski, E. S. (1997). Formation of cytomegalovirus DNA replication compartments defined by localization of viral proteins and DNA synthesis. Virology 239, 46-61.

Petalidis, L., Bhattacharyya, S., Morris, G. A., Collins, V. P., Freeman, T. C., and Lyons, P. A. (2003). Global amplification of mRNA by template-switching PCR: linearity and application to microarray analysis. Nucleic Acids Res 31, e142.

Petrik, J. (2001). Microarray technology: the future of blood testing? Vox Sang 80, 1-11.

Pietiainen, V., Huttunen, P., and Hyypia, T. (2000). Effects of echovirus 1 infection on cellular gene expression. Virology 276, 243-250.

Pietroboni, G. R., Harnett, G. B., and Bucens, M. R. (1989). Centrifugal enhancement of human immunodeficiency virus (HIV) and human herpesvirus type 6 (HHV-6) infection in vitro. J Virol Methods 24, 85-90.

Piguet, V., Schwartz, O., Le Gall, S., and Trono, D. (1999). The downregulation of CD4 and MHC-I by primate lentiviruses: a paradigm for the modulation of cell surface receptors. Immunol Rev 168, 51-63.

Poggioli, G. J., DeBiasi, R. L., Bickel, R., Jotte, R., Spalding, A., Johnson, G. L., and Tyler, K. L. (2002). Reovirus-induced alterations in gene expression related to cell cycle regulation. J Virol 76, 2585-2594.

Polson, A. G., Wang, D., DeRisi, J., and Ganem, D. (2002). Modulation of host gene expression by the constitutively active G protein-coupled receptor of Kaposi's sarcoma-associated herpesvirus. Cancer Res 62, 4525-4530.
Poole, L. J., Yu, Y., Kim, P. S., Zheng, Q. Z., Pevsner, J., and Hayward, G. S. (2002). Altered patterns of cellular gene expression in dermal microvascular endothelial cells infected with Kaposi's sarcoma-associated herpesvirus. J Virol 76, 3395-3420.

Puskas, L. G., Zvara, A., Hackler, L., Jr., and Van Hummelen, P. (2002). RNA amplification results in reproducible microarray data with slight ratio bias. Biotechniques 32, 1330-1334, 1336, 1338, 1340.

Qian, G., Wood, C., and Chandran, B. (1993). Identification and characterization of glycoprotein gH of human herpesvirus-6. Virology 194, 380-386.

Quackenbush, J. (2001). Computational analysis of microarray data. Nat Rev Genet 2, 418-427.

Quackenbush, J. (2002). Microarray data normalization and transformation. Nat Genet 32 Suppl, 496-501.

Rajeevan, M. S., Dimulescu, I. M., Vernon, S. D., Verma, M., and Unger, E. R. (2003). Global amplification of sense RNA: a novel method to replicate and archive mRNA for gene expression analysis. Genomics 82, 491-497.

Rapp, J. C., Krug, L. T., Inoue, N., Dambaugh, T. R., and Pellett, P. E. (2000). U94, the human herpesvirus 6 homolog of the parvovirus nonstructural gene, is highly conserved among isolates and is expressed at low mRNA levels as a spliced transcript. Virology 268, 504-516.

Rathod, P. K., Ganesan, K., Hayward, R. E., Bozdech, Z., and DeRisi, J. L. (2002). DNA microarrays for malaria. Trends Parasitol 18, 39-45.

Reading, P. C., Moore, J. B., and Smith, G. L. (2003). Steroid hormone synthesis by vaccinia virus suppresses the inflammatory response to infection. J Exp Med 197, 1269-1278.

Reading, S. A., Edwards, M. J., and Dimmock, N. J. (2001). Increasing the efficiency of virus infectivity assays: small inoculum volumes are as effective as centrifugal enhancement. J Virol Methods 98, 167-169.

Renne, R., Barry, C., Dittmer, D., Compitello, N., Brown, P. O., and Ganem, D. (2001). Modulation of cellular and viral gene expression by the latency-associated nuclear antigen of Kaposi's sarcoma-associated herpesvirus. J Virol 75, 458-468.

Richmond, C. S., Glasner, J. D., Mau, R., Jin, H., and Blattner, F. R. (1999). Genome-wide expression profiling in Escherichia coli K-12. Nucleic Acids Res 27, 3821-3835.

Roizman, B., Knipe, D. M. (2001a). Herpes simplex viruses and their replication. In Fields Virology, D. M. Knipe, Howley, P. M., ed. (Philadelphia, Lippincott Williams & Wilkins), pp. 2399-2460.

Roizman, B., Pellet, P.E. (2001b). The family herpesviridae: A brief introduction. In Fields Virology, D. M. Knipe, Howley, P. M., ed. (Philadelphia, Lippincott Willams & Wilkins), pp. 2381-2398.

Romanowski, M. J., and Shenk, T. (1997). Characterization of the human cytomegalovirus irs1 and trs1 genes: a second immediate-early transcription unit within irs1 whose product antagonizes transcriptional activation. J Virol 71, 1485-1496.

Rosenberger, C. M., Scott, M. G., Gold, M. R., Hancock, R. E., and Finlay, B. B. (2000). Salmonella typhimurium infection and lipopolysaccharide stimulation induce similar changes in macrophage gene expression. J Immunol *164*, 5894-5904.

Ross, D. T., Scherf, U., Eisen, M. B., Perou, C. M., Rees, C., Spellman, P., Iyer, V., Jeffrey, S. S., Van de Rijn, M., Waltham, M., et al. (2000). Systematic variation in gene expression patterns in human cancer cell lines. Nat Genet 24, 227-235.

Rozen, S., Skaletsky, H. J. (2000). Primer3 on the WWW for general users and for biologist programmers. In Bioinformatics Methods and Protocols: Methods in Molecular Biology, S. Krawetz, Misener, S., ed. (Totowa, NJ, Humana Press), pp. 365-386.

Ryo, A., Suzuki, Y., Arai, M., Kondoh, N., Wakatsuki, T., Hada, A., Shuda, M., Tanaka, K., Sato, C., Yamamoto, M., and Yamamoto, N. (2000). Identification and characterization of differentially expressed mRNAs in HIV type 1-infected human T cells. AIDS Res Hum Retroviruses 16, 995-1005.

Sada, E., Yasukawa, M., Ito, C., Takeda, A., Shiosaka, T., Tanioka, H., and Fujita, S. (1996). Detection of human herpesvirus 6 and human herpesvirus 7 in the submandibular gland, parotid gland, and lip salivary gland by PCR. J Clin Microbiol 34, 2320-2321.

Saeed, A. I., Sharov, V., White, J., Li, J., Liang, W., Bhagabati, N., Braisted, J., Klapa, M., Currier, T., Thiagarajan, M., *et al.* (2003). TM4: a free, open-source system for microarray data management and analysis. Biotechniques 34, 374-378.

Salahuddin, S. Z., Ablashi, D. V., Markham, P. D., Josephs, S. F., Sturzenegger, S., Kaplan, M., Halligan, G., Biberfeld, P., Wong-Staal, F., Kramarsky, B., and et al. (1986). Isolation of a new virus, HBLV, in patients with lymphoproliferative disorders. Science 234, 596-601.

Salama, N., Guillemin, K., McDaniel, T. K., Sherlock, G., Tompkins, L., and Falkow, S. (2000). A whole-genome microarray reveals genetic diversity among Helicobacter pylori strains. Proc Natl Acad Sci U S A 97, 14668-14673.

Sanger, F., Nicklen, S., and Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. Proc Natl Acad Sci U S A 74, 5463-5467.

Sarisky, R. T., and Hayward, G. S. (1996). Evidence that the UL84 gene product of human cytomegalovirus is essential for promoting oriLyt-dependent DNA replication and formation of replication compartments in cotransfection assays. J Virol 70, 7398-7413.

Sasada, A., Takaori-Kondo, A., Shirakawa, K., Kobayashi, M., Abudu, A., Hishizawa, M., Imada, K., Tanaka, Y., and Uchiyama, T. (2005). APOBEC3G targets human T-cell leukemia virus type 1. Retrovirology 2, 32.

Satterthwaite, G., Francis, S. E., Suvarna, K., Blakemore, S., Ward, C., Wallace, D., Braddock, M., and Crossman, D. (2005). Differential gene expression in coronary arteries from patients presenting with ischemic heart disease: further evidence for the inflammatory basis of atherosclerosis. Am Heart J 150, 488-499.

Schena, M., Shalon, D., Davis, R. W., and Brown, P. O. (1995). Quantitative monitoring of gene expression patterns with a complementary DNA microarray. Science 270, 467-470.

Scherf, U., Ross, D. T., Waltham, M., Smith, L. H., Lee, J. K., Tanabe, L., Kohn, K. W., Reinhold, W. C., Myers, T. G., Andrews, D. T., *et al.* (2000). A gene expression database for the molecular pharmacology of cancer. Nat Genet 24, 236-244.

Schindler, H., Wiese, A., Auer, J., and Burtscher, H. (2005). cRNA target preparation for microarrays: comparison of gene expression profiles generated with different amplification procedures. Anal Biochem 344, 92-101.

Seal, B. S., Irving, J. M., and Whetstone, C. A. (1991). Transcriptional analysis of the bovine herpesvirus 1 Cooper isolate. Temporal analysis and characterization of immediate-early, early, and late RNA. Arch Virol 121, 55-73.

Secchiero, P., Berneman, Z. N., Gallo, R. C., and Lusso, P. (1994). Biological and molecular characteristics of human herpesvirus 7: in vitro growth optimization and development of a syncytia inhibition test. Virology 202, 506-512.

Secchiero, P., Bertolaso, L., Casareto, L., Gibellini, D., Vitale, M., Bemis, K., Aleotti, A., Capitani, S., Franchini, G., Gallo, R. C., and Zauli, G. (1998a). Human herpesvirus 7 infection induces profound cell cycle perturbations coupled to disregulation of cdc2 and cyclin B and polyploidization of CD4(+) T cells. Blood 92, 1685-1696.

Secchiero, P., Bertolaso, L., Gibellini, D., Ricci, D., Bemis, K., Capitani, S., Gallo, R. C., and Zauli, G. (1998b). Enforced expression of human bcl-2 in CD4+ T cells enhances human herpesvirus 7 replication and induction of cytopathic effects. Eur J Immunol 28, 1587-1596.

Secchiero, P., Flamand, L., Gibellini, D., Falcieri, E., Robuffo, I., Capitani, S., Gallo, R. C., and Zauli, G. (1997a). Human Herpesvirus 7 induces CD4(+) T-cell death by two distinct mechanisms: necrotic lysis in productively infected cells and apoptosis in uninfected or nonproductively infected cells. Blood 90, 4502-4512.

Secchiero, P., Gibellini, D., Flamand, L., Robuffo, I., Marchisio, M., Capitani, S., Gallo, R. C., and Zauli, G. (1997b). Human herpesvirus 7 induces the down-regulation of CD4 antigen in lymphoid T cells without affecting p56lck levels. J Immunol 159, 3412-3423.

Secchiero, P., Mirandola, P., Zella, D., Celeghini, C., Gonelli, A., Vitale, M., Capitani, S., and Zauli, G. (2001). Human herpesvirus 7 induces the functional up-regulation of tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) coupled to TRAIL-R1 down-modulation in CD4(+) T cells. Blood 98, 2474-2481.

Secchiero, P., Sun, D., De Vico, A. L., Crowley, R. W., Reitz, M. S., Jr., Zauli, G., Lusso, P., and Gallo, R. C. (1997c). Role of the extracellular domain of human herpesvirus 7 glycoprotein B in virus binding to cell surface heparan sulfate proteoglycans. J Virol 71, 4571-4580.

Secchiero, P., Zella, D., Barabitskaja, O., Reitz, M. S., Capitani, S., Gallo, R. C., and Zauli, G. (1998c). Progressive and persistent downregulation of surface CXCR4 in CD4(+) T cells infected with human herpesvirus 7. Blood 92, 4521-4528.

Seth, D., Gorrell, M. D., McGuinness, P. H., Leo, M. A., Lieber, C. S., McCaughan, G. W., and Haber, P. S. (2003). SMART amplification maintains representation of relative gene expression: quantitative validation by real time PCR and application to studies of alcoholic liver disease in primates. J Biochem Biophys Methods 55, 53-66.

Shafti-Keramat, S., Handisurya, A., Kriehuber, E., Meneguzzi, G., Slupetzky, K., and Kirnbauer, R. (2003). Different heparan sulfate proteoglycans serve as cellular receptors for human papillomaviruses. J Virol 77, 13125-13135.

Shaheduzzaman, S., Krishnan, V., Petrovic, A., Bittner, M., Meltzer, P., Trent, J., Venkatesan, S., and Zeichner, S. (2002). Effects of HIV-1 Nef on cellular gene expression profiles. J Biomed Sci 9, 82-96.

Shen, R., Fan, J. B., Campbell, D., Chang, W., Chen, J., Doucet, D., Yeakley, J., Bibikova, M., Wickham Garcia, E., McBride, C., *et al.* (2005). High-throughput SNP genotyping on universal bead arrays. Mutat Res 573, 70-82.

Shiraki, K., Mukai, T., Okuno, T., Yamanishi, K., and Takahashi, M. (1991). Physicochemical characterization of human herpesvirus 6 infectivity. J Gen Virol 72 (Pt 1), 169-172.

Shyamsundar, R., Kim, Y. H., Higgins, J. P., Montgomery, K., Jorden, M., Sethuraman, A., van de Rijn, M., Botstein, D., Brown, P. O., and Pollack, J. R. (2005). A DNA microarray survey of gene expression in normal human tissues. Genome Biol 6, R22.

Simmen, K. A., Singh, J., Luukkonen, B. G., Lopper, M., Bittner, A., Miller, N. E., Jackson, M. R., Compton, T., and Fruh, K. (2001). Global modulation of cellular transcription by human cytomegalovirus is initiated by viral glycoprotein B. Proc Natl Acad Sci U S A 98, 7140-7145.

Simmons, A., Aluvihare, V., and McMichael, A. (2001). Nef triggers a transcriptional program in T cells imitating single-signal T cell activation and inducing HIV virulence mediators. Immunity 14, 763-777.

Skrincosky, D., Hocknell, P., Whetter, L., Secchiero, P., Chandran, B., and Dewhurst, S. (2000). Identification and analysis of a novel heparin-binding glycoprotein encoded by human herpesvirus 7. J Virol 74, 4530-4540.

Song, J. H., Kim, J. M., Kim, S. H., Kim, H. J., Lee, J. J., Sung, M. H., Hwang, S. Y., and Kim, T. S. (2003). Comparison of the gene expression profiles of monocytic versus granulocytic lineages of HL-60 leukemia cell differentiation by DNA microarray analysis. Life Sci 73, 1705-1719.

Song, Y. J., and Stinski, M. F. (2002). Effect of the human cytomegalovirus IE86 protein on expression of E2F-responsive genes: a DNA microarray analysis. Proc Natl Acad Sci U S A 99, 2836-2841.

Soukas, A., Cohen, P., Socci, N. D., and Friedman, J. M. (2000). Leptin-specific patterns of gene expression in white adipose tissue. Genes Dev 14, 963-980.

Spear, P. G., and Longnecker, R. (2003). Herpesvirus entry: an update. J Virol 77, 10179-10185.

Spector, D. H. (1996). Activation and regulation of human cytomegalovirus early genes. Intervirology 39, 361-377.

Stamminger, T., Puchtler, E., and Fleckenstein, B. (1991). Discordant expression of the immediateearly 1 and 2 gene regions of human cytomegalovirus at early times after infection involves posttranscriptional processing events. J Virol 65, 2273-2282.

Stasiak, P. C., and Mocarski, E. S. (1992). Transactivation of the cytomegalovirus ICP36 gene promoter requires the alpha gene product TRS1 in addition to IE1 and IE2. J Virol 66, 1050-1058.

Steenman, M., Lamirault, G., Le Meur, N., Le Cunff, M., Escande, D., and Leger, J. J. (2005). Distinct molecular portraits of human failing hearts identified by dedicated cDNA microarrays. Eur J Heart Fail 7, 157-165.

Steller, S., Angenendt, P., Cahill, D. J., Heuberger, S., Lehrach, H., and Kreutzberger, J. (2005). Bacterial protein microarrays for identification of new potential diagnostic markers for Neisseria meningitidis infections. Proteomics 5, 2048-2055.

Stenberg, R. M. (1996). The human cytomegalovirus major immediate-early gene. Intervirology 39, 343-349.

Stenberg, R. M., Depto, A. S., Fortney, J., and Nelson, J. A. (1989). Regulated expression of early and late RNAs and proteins from the human cytomegalovirus immediate-early gene region. J Virol 63, 2699-2708.

Stingley, S. W., Ramirez, J. J., Aguilar, S. A., Simmen, K., Sandri-Goldin, R. M., Ghazal, P., and Wagner, E. K. (2000). Global analysis of herpes simplex virus type 1 transcription using an oligonucleotide-based DNA microarray. J Virol 74, 9916-9927.

Stinski, M. F. (1978). Sequence of protein synthesis in cells infected by human cytomegalovirus: early and late virus-induced polypeptides. J Virol 26, 686-701.

Stinski, M. F., Mocarski, E. S., and Thomsen, D. R. (1979). DNA of human cytomegalovirus: size heterogeneity and defectiveness resulting from serial undiluted passage. J Virol 31, 231-239.

Stoughton, R. B. (2005). Applications of DNA microarrays in biology. Annu Rev Biochem 74, 53-82.

Sullivan, D. C., Gray, W. L., Caughman, G. B., Robertson, A. T., and O'Callaghan, D. J. (1990). Temporal regulation of equine herpesvirus type 3 transcription. Virus Res 15, 135-148.

Takeda, K., Okuno, T., Isegawa, Y., and Yamanishi, K. (1996). Identification of a variant A-specific neutralizing epitope on glycoprotein B (gB) of human herpesvirus-6 (HHV-6). Virology 222, 176-183.

Talbot, S. G., Estilo, C., Maghami, E., Sarkaria, I. S., Pham, D. K., P, O. c., Socci, N. D., Ngai, I., Carlson, D., Ghossein, R., *et al.* (2005). Gene expression profiling allows distinction between primary and metastatic squamous cell carcinomas in the lung. Cancer Res *65*, 3063-3071.

Tamayo, P., Slonim, D., Mesirov, J., Zhu, Q., Kitareewan, S., Dmitrovsky, E., Lander, E. S., and Golub, T. R. (1999). Interpreting patterns of gene expression with self-organizing maps: methods and application to hematopoietic differentiation. Proc Natl Acad Sci U S A 96, 2907-2912.

Tan, F. L., Moravec, C. S., Li, J., Apperson-Hansen, C., McCarthy, P. M., Young, J. B., and Bond, M. (2002). The gene expression fingerprint of human heart failure. Proc Natl Acad Sci U S A 99, 11387-11392.

Tanaka-Taya, K., Kondo, T., Nakagawa, N., Inagi, R., Miyoshi, H., Sunagawa, T., Okada, S., and Yamanishi, K. (2000). Reactivation of human herpesvirus 6 by infection of human herpesvirus 7. J Med Virol 60, 284-289.

Tang, B. S., Chan, K. H., Cheng, V. C., Woo, P. C., Lau, S. K., Lam, C. C., Chan, T. L., Wu, A. K., Hung, I. F., Leung, S. Y., and Yuen, K. Y. (2005). Comparative host gene transcription by microarray analysis early after infection of the Huh7 cell line by severe acute respiratory syndrome coronavirus and human coronavirus 229E. J Virol 79, 6180-6193.

Tavazoie, S., Hughes, J. D., Campbell, M. J., Cho, R. J., and Church, G. M. (1999). Systematic determination of genetic network architecture. Nat Genet 22, 281-285.

Taylor, L. A., Carthy, C. M., Yang, D., Saad, K., Wong, D., Schreiner, G., Stanton, L. W., and McManus, B. M. (2000). Host gene regulation during coxsackievirus B3 infection in mice: assessment by microarrays. Circ Res 87, 328-334.

Tenser, R. B., and Dunstan, M. E. (1980). Mechanisms of herpes simplex virus infectivity enhanced by ultracentrifugal inoculation. Infect Immun 30, 193-197.

Tian, B., Zhang, Y., Luxon, B. A., Garofalo, R. P., Casola, A., Sinha, M., and Brasier, A. R. (2002). Identification of NF-kappaB-dependent gene networks in respiratory syncytial virus-infected cells. J Virol 76, 6800-6814.

Torigoe, S., Kumamoto, T., Koide, W., Taya, K., and Yamanishi, K. (1995). Clinical manifestations associated with human herpesvirus 7 infection. Arch Dis Child 72, 518-519.

Troesch, A., Nguyen, H., Miyada, C. G., Desvarenne, S., Gingeras, T. R., Kaplan, P. M., Cros, P., and Mabilat, C. (1999). Mycobacterium species identification and rifampin resistance testing with high-density DNA probe arrays. J Clin Microbiol 37, 49-55.

Tsai, C. A., Lee, T. C., Ho, I. C., Yang, U. C., Chen, C. H., and Chen, J. J. (2005). Multi-class clustering and prediction in the analysis of microarray data. Math Biosci 193, 79-100.

Tsavachidou, D., Podrzucki, W., Seykora, J., and Berger, S. L. (2001). Gene array analysis reveals changes in peripheral nervous system gene expression following stimuli that result in reactivation of latent herpes simplex virus type 1: induction of transcription factor Bcl-3. J Virol 75, 9909-9917.

Tsunedomi, R., Iizuka, N., Hamamoto, Y., Uchimura, S., Miyamoto, T., Tamesa, T., Okada, T., Takemoto, N., Takashima, M., Sakamoto, K., *et al.* (2005). Patterns of expression of cytochrome P450 genes in progression of hepatitis C virus-associated hepatocellular carcinoma. Int J Oncol 27, 661-667.

Tusher, V. G., Tibshirani, R., and Chu, G. (2001). Significance analysis of microarrays applied to the ionizing radiation response. Proc Natl Acad Sci U S A 98, 5116-5121.

Ueda, K., Kusuhara, K., Okada, K., Miyazaki, C., Hidaka, Y., Tokugawa, K., and Yamanishi, K. (1994). Primary human herpesvirus 7 infection and exanthema subitum. Pediatr Infect Dis J 13, 167-168.

Vahey, M. T., Nau, M. E., Jagodzinski, L. L., Yalley-Ogunro, J., Taubman, M., Michael, N. L., and Lewis, M. G. (2002). Impact of viral infection on the gene expression profiles of proliferating normal human peripheral blood mononuclear cells infected with HIV type 1 RF. AIDS Res Hum Retroviruses 18, 179-192.

Van Gelder, R. N., von Zastrow, M. E., Yool, A., Dement, W. C., Barchas, J. D., and Eberwine, J. H. (1990). Amplified RNA synthesized from limited quantities of heterogeneous cDNA. Proc Natl Acad Sci U S A 87, 1663-1667.

van 't Wout, A. B., Lehrman, G. K., Mikheeva, S. A., O'Keeffe, G. C., Katze, M. G., Bumgarner, R. E., Geiss, G. K., and Mullins, J. I. (2003). Cellular gene expression upon human immunodeficiency virus type 1 infection of CD4(+)-T-cell lines. J Virol 77, 1392-1402.

Vazquez-Ortiz, G., Pina-Sanchez, P., Hidalgo, A., Lazos, M., Moreno, J., Alvarado, I., Cruz, F., Hernandez, D. M., Perez-Plascencia, C., and Salcedo, M. (2005). [Global expression analysis in uterine cervical cancer: metabolic pathways and altered genes]. Rev Invest Clin 57, 434-441.

Vertegaal, A. C., Kuiperij, H. B., van Laar, T., Scharnhorst, V., van der Eb, A. J., and Zantema, A. (2000). cDNA micro array identification of a gene differentially expressed in adenovirus type 5- versus type 12-transformed cells. FEBS Lett 487, 151-155.

Wagner, A., Doerks, A., Aboud, M., Alonso, A., Tokino, T., Flugel, R. M., and Lochelt, M. (2000). Induction of cellular genes is mediated by the Bell transactivator in foamy virus-infected human cells. J Virol 74, 4441-4447. Wagner, E. K., Swanstrom, R. I., and Stafford, M. G. (1972). Transcription of the herpes simplex virus genome in human cells. J Virol 10, 675-682.

Walduck, A., Rudel, T., and Meyer, T. F. (2004). Proteomic and gene profiling approaches to study host responses to bacterial infection. Curr Opin Microbiol 7, 33-38.

Wang, D., Coscoy, L., Zylberberg, M., Avila, P. C., Boushey, H. A., Ganem, D., and DeRisi, J. L. (2002a). Microarray-based detection and genotyping of viral pathogens. Proc Natl Acad Sci U S A 99, 15687-15692.

Wang, H., Hao, L., Shung, C. Y., Sunter, G., and Bisaro, D. M. (2003). Adenosine kinase is inactivated by geminivirus AL2 and L2 proteins. Plant Cell 15, 3020-3032.

Wang, J., Delabie, J., Aasheim, H., Smeland, E., and Myklebost, O. (2002b). Clustering of the SOM easily reveals distinct gene expression patterns: results of a reanalysis of lymphoma study. BMC Bioinformatics 3, 36.

Wang, Z. W., Sarmento, L., Wang, Y., Li, X. Q., Dhingra, V., Tseggai, T., Jiang, B., and Fu, Z. F. (2005). Attenuated rabies virus activates, while pathogenic rabies virus evades, the host innate immune responses in the central nervous system. J Virol 79, 12554-12565.

Wathen, M. W., and Stinski, M. F. (1982). Temporal patterns of human cytomegalovirus transcription: mapping the viral RNAs synthesized at immediate early, early, and late times after infection. J Virol 41, 462-477.

Wenli, M., Yan, W., Hongmin, W., and Wenling, Z. (2004). An oligonucleotide microarray for the detection of vaccinia virus. Br J Biomed Sci 61, 142-145.

Wilson, C. L., Pepper, S. D., Hey, Y., and Miller, C. J. (2004). Amplification protocols introduce systematic but reproducible errors into gene expression studies. Biotechniques 36, 498-506.

Wilson, M., DeRisi, J., Kristensen, H. H., Imboden, P., Rane, S., Brown, P. O., and Schoolnik, G. K. (1999). Exploring drug-induced alterations in gene expression in Mycobacterium tuberculosis by microarray hybridization. Proc Natl Acad Sci U S A 96, 12833-12838.

Wyatt, L. S., and Frenkel, N. (1992). Human herpesvirus 7 is a constitutive inhabitant of adult human saliva. J Virol 66, 3206-3209.

Wyatt, L. S., Rodriguez, W. J., Balachandran, N., and Frenkel, N. (1991). Human herpesvirus 7: antigenic properties and prevalence in children and adults. J Virol 65, 6260-6265.

Yadav, M., Nambiar, S., Khoo, S. P., and Yaacob, H. B. (1997). Detection of human herpesvirus 7 in salivary glands. Arch Oral Biol 42, 559-567.

Yamanishi, K., Okuno, T., Shiraki, K., Takahashi, M., Kondo, T., Asano, Y., and Kurata, T. (1988). Identification of human herpesvirus-6 as a causal agent for exanthem subitum. Lancet 1, 1065-1067.

Yang, I. V., Chen, E., Hasseman, J. P., Liang, W., Frank, B. C., Wang, S., Sharov, V., Saeed, A. I., White, J., Li, J., *et al.* (2002a). Within the fold: assessing differential expression measures and reproducibility in microarray assays. Genome Biol *3*, research0062.

Yang, Y. H., Dudoit, S., Luu, P., Lin, D. M., Peng, V., Ngai, J., and Speed, T. P. (2002b). Normalization for cDNA microarray data: a robust composite method addressing single and multiple slide systematic variation. Nucleic Acids Res 30, e15.

Yang, Y. H., and Speed, T. (2002). Design issues for cDNA microarray experiments. Nat Rev Genet 3, 579-588.

Yasukawa, M., Hasegawa, A., Sakai, I., Ohminami, H., Arai, J., Kaneko, S., Yakushijin, Y., Maeyama, K., Nakashima, H., Arakaki, R., and Fujita, S. (1999). Down-regulation of CXCR4 by human herpesvirus 6 (HHV-6) and HHV-7. J Immunol 162, 5417-5422.

Yasukawa, M., Inoue, Y., Ohminami, H., Sada, E., Miyake, K., Tohyama, T., Shimada, T., and Fujita, S. (1997). Human herpesvirus 7 infection of lymphoid and myeloid cell lines transduced with an adenovirus vector containing the CD4 gene. J Virol 71, 1708-1712.

Yasukawa, M., Yakushijin, Y., Furukawa, M., and Fujita, S. (1993). Specificity analysis of human CD4+ T-cell clones directed against human herpesvirus 6 (HHV-6), HHV-7, and human cytomegalovirus. J Virol 67, 6259-6264.

Yoshikawa, T., Asano, Y., Kobayashi, I., Nakashima, T., Yazaki, T., Suga, S., Ozaki, T., Wyatt, L. S., and Frenkel, N. (1993). Seroepidemiology of human herpesvirus 7 in healthy children and adults in Japan. J Med Virol 41, 319-323.

Yuen, T., Wurmbach, E., Pfeffer, R. L., Ebersole, B. J., and Sealfon, S. C. (2002). Accuracy and calibration of commercial oligonucleotide and custom cDNA microarrays. Nucleic Acids Res 30, e48.

Zammatteo, N., Hamels, S., De Longueville, F., Alexandre, I., Gala, J. L., Brasseur, F., and Remacle, J. (2002). New chips for molecular biology and diagnostics. Biotechnol Annu Rev 8, 85-101.

Zhang, Y., de Bolle, L., Aquaro, S., van Lommel, A., De Clercq, E., and Schols, D. (2001a). Productive infection of primary macrophages with human herpesvirus 7. J Virol 75, 10511-10514.

Zhang, Y., Hatse, S., De Clercq, E., and Schols, D. (2000). CXC-chemokine receptor 4 is not a coreceptor for human herpesvirus 7 entry into CD4(+) T cells. J Virol 74, 2011-2016.

Zhang, Y., Luxon, B. A., Casola, A., Garofalo, R. P., Jamaluddin, M., and Brasier, A. R. (2001b). Expression of respiratory syncytial virus-induced chemokine gene networks in lower airway epithelial cells revealed by cDNA microarrays. J Virol 75, 9044-9058.

Zhao, H., Hastie, T., Whitfield, M. L., Borresen-Dale, A. L., and Jeffrey, S. S. (2002). Optimization and evaluation of T7 based RNA linear amplification protocols for cDNA microarray analysis. BMC Genomics 3, 31.

Zhu, H., Bilgin, M., Bangham, R., Hall, D., Casamayor, A., Bertone, P., Lan, N., Jansen, R., Bidlingmaier, S., Houfek, T., et al. (2001). Global analysis of protein activities using proteome chips. Science 293, 2101-2105.

Zhu, H., Cong, J. P., Mamtora, G., Gingeras, T., and Shenk, T. (1998). Cellular gene expression altered by human cytomegalovirus: global monitoring with oligonucleotide arrays. Proc Natl Acad Sci U S A 95, 14470-14475.

Zhu, Z., Gershon, M. D., Ambron, R., Gabel, C., and Gershon, A. A. (1995). Infection of cells by varicella zoster virus: inhibition of viral entry by mannose 6-phosphate and heparin. Proc Natl Acad Sci U S A 92, 3546-3550.