

The association of maternal prenatal psychosocial stress with vascular function in the child at age 10–11 years: findings from the Avon longitudinal study of parents and children

Aimée E van Dijk^{1,2}, Karen Dawe³, John Deanfield⁴,
Karien Stronks⁵, Reinoud JBJ Gemke⁴, Tanja GM Vrijkotte¹ and
Debbie A Lawlor³

Abstract

Objective: To investigate whether (1) maternal psychosocial stress (depression/anxiety) during pregnancy is associated with offspring vascular function and (2) whether any association differs depending on the gestational timing of exposure to stress. We also investigated whether any association is likely to be due to intrauterine mechanisms by (3) comparing with the association of paternal stress with offspring vascular function and (4) examining whether any prenatal association is explained by maternal postnatal stress.

Methods and results: Associations were examined in a UK birth cohort, with offspring outcomes (systolic and diastolic blood pressure, SBP and DBP, endothelial function assessed by brachial artery flow-mediated dilatation (FMD); arterial stiffness assessed by carotid to radial pulse wave velocity (PWV), brachial artery distensibility (DC), and brachial artery diameter (BD) assessed at age 10–11 years ($n = 4318$). Maternal depressive symptoms and anxiety were assessed at 18 and 32 weeks gestation and 8 months postnatally. Paternal symptoms were assessed at week 19. With the exception of DBP and BD, there were no associations of maternal depressive symptoms with any of the vascular outcomes. Maternal depressive and anxiety symptoms were associated with lower offspring DBP and wider BD, though the latter attenuated to the null with adjustment for confounding factors. Paternal symptoms were not associated with offspring outcomes. Maternal postnatal depressive symptoms were associated with lower offspring SBP.

Conclusions: We found no evidence to support the hypothesis that maternal stress during pregnancy adversely affects offspring vascular function at age 10–12 years via intrauterine mechanisms.

Keywords

Arterial stiffness, blood pressure, child, developmental origins of health and disease, fetal development, pregnancy, psychological, stress

Received 6 July 2012; accepted 21 March 2013

Introduction

Maternal stress or the administration of glucocorticoids ('stress hormones') have been linked to altered fetal growth and might have permanent effects on the tissue structure and function of the offspring.¹ For example, the hypothalamic–pituitary–adrenal axis, which is a key part of the neuroendocrine stress response, and the renin–angiotensin system may be affected by intrauterine exposure to maternal stress,

¹University of Amsterdam, Amsterdam, The Netherlands

²Public Health Service of Amsterdam (GGD), Amsterdam, The Netherlands

³University of Bristol, Bristol, UK

⁴University College London, London, UK

⁵VU University Medical Center, Amsterdam, The Netherlands

Corresponding author:

Aimée E van Dijk, Academic Medical Center, Dept. of Public Health, PO Box 22660, 1100 DD Amsterdam, The Netherlands.
Email: a.e.vandijk@amc.uva.nl

and this exposure may lead to adult risk of cardiovascular disease.^{2–17}

Convincing evidence for the effects of fetal programming on cardiovascular risk factors comes from animal studies, in which prenatal exposure to excess glucocorticoids results in persistent elevation of arterial blood pressure (BP) in adulthood, with potential sex differences (for review, see Drake et al.¹⁸). There is also some evidence from human studies. For example, antenatal glucocorticoid administration is linked with high BP in the fetus¹⁹ and in adolescence.²⁰ Those studies have, however, been complicated by high doses of glucocorticoids, which are not comparable to levels normally occurring during pregnancy. In addition, glucocorticoid treatment usually takes place when pre-term birth is expected, adding bias because of the complications of premature birth. Few pregnant women will be treated with glucocorticoids, but the results of these studies have been used to suggest that common mental complaints (depression and anxiety) associated with stress, if experienced by the mother during pregnancy might increase her offspring's risk of cardiovascular disease. If this is true it would have important public health implications, but to our knowledge few studies have examined this.

In a previous study in a Dutch population-based birth cohort (ABCD study), the presence of multiple psychosocial stressors during early pregnancy was associated with higher BP in the offspring at age five.²¹ However, we are unaware of any other human studies of this association and this needs further replication and additional exploration to see if any association is likely to reflect a causal intrauterine mechanism.

The aim of this study was to examine whether (1) maternal psychosocial stress (depression/anxiety) during pregnancy is associated with offspring vascular function (assessed by systolic blood pressure (SBP), diastolic blood pressure (DBP), endothelial function assessed by brachial artery flow-mediated dilatation (FMD), arterial stiffness assessed by carotid to radial pulse wave velocity (PWV), brachial distensibility coefficient (DC), and brachial artery diameter) and (2) whether any association differs depending on the gestational timing of exposure to stress. We further examined whether there is evidence that any association of maternal stress with offspring vascular function is likely to be due to intrauterine mechanisms by comparing the associations of maternal stress with offspring outcomes to those of paternal antenatal stress with the same outcomes and by examining whether any maternal prenatal association was explained by the persistence of symptoms into the postnatal period and exposure postnatally (rather than intrauterine driving associations). The paternal comparison is an important form of negative control for exploring a causal association in

observational epidemiology.^{22,23} The rationale is that if maternal stress is truly causing offspring outcomes via intrauterine mechanisms (as opposed to family based confounding) then one would expect a stronger maternal association, compared to paternal association. Examining whether any maternal pregnancy stress association with offspring outcomes is largely mediated by postnatal stress is also important as if the effect is largely due to postnatal impact then any intervention to prevent adverse offspring outcomes would want to focus on minimizing postnatal stress.

Methods

Study population

Data from the Avon Longitudinal Study of Parents and Children (ALSPAC), a prospective population-based birth cohort that recruited 14,541 pregnant women resident in Avon, UK with expected dates of delivery 1 April 1991 to 31 December 1992 were used. Additional details of the study can be found on the study website (www.alspac.bris.ac.uk) and in previous publications.^{24–26} Figure 1 shows the flow of participants through the study. In this study, we included mother–offspring pairs where there was a singleton pregnancy resulting in a live birth with the child surviving to at least 1 year of age. The eligible cohort for the current analyses was the 4318 mother–offspring pairs who had complete data on maternal depressive symptoms and anxiety at both gestational time points, offspring blood pressure (obtained at the 10–11-year follow-up clinic), and complete data on all covariates.

Parental assessment of psychosocial stress

Data on depressive symptoms and anxiety were collected through postal questionnaires, which the mothers in the current sample filled out at a median gestational age of 18 weeks (mean 19; interquartile range 18–20) and median 32 weeks (mean 32; interquartile range 32–33) antenatally and median 8 months postnatally (mean 8.6; interquartile range 8.1–9.1). Fathers completed one questionnaire at a median 19 weeks of their partners' pregnancy (mean 20; interquartile range 19–21). Maternal and paternal depressive symptoms at all time points were assessed using the Edinburgh Postnatal Depression Scale (EPDS), a widely used 10-item self-report questionnaire that has been shown to be valid in and outside the postnatal period.²⁷ A pilot study of a random sample of 54 pregnant women attending a routine check-up indicated substantial overlap between the EPDS and the Center for Epidemiological Studies Depression Scale ($r^2=0.87$, $p<0.001$). Internal consistency (Cronbach's




Figure 1. Participant flow.

Data in the yellow boxes are for mother(father)–child pairs with complete data.

alpha) of the EPDS exceeded 0.80 at each of the assessments.²⁸

Maternal and paternal anxiety at all time points was measured using the anxiety items from the Crown–Crisp experiential index (CCEI), a validated self-rating inventory (for example, see Crisp et al.²⁹); example items include ‘worry a lot’ and ‘feeling strung up inside’. In a pilot study of a random sample of 54 pregnant women attending a routine check-up, Crown–Crisp index correlated 0.70 and 0.76 with the State and Trait (respectively) subscales of the Spielberger State-trait Anxiety Inventory. Internal consistency (Cronbach’s alpha) exceeded 0.80 at each of the assessments.²⁸

To compare prenatal and postnatal maternal stress we categorized all EPDS and CCEI scores into ‘high

levels of anxiety/depressive symptoms’ (yes versus no) by dichotomizing at the 80th percentile (those classified as having ‘high levels’ included those with values at or higher than the 80th percentile; those classified as ‘normal’ included those with values lower than the 80th percentile). We then further categorized participants into those with normal levels at all time points; prenatal high levels only; postnatal high levels only, or pre- and postnatal high levels.

Vascular function in the child

Vascular phenotypes were assessed at the 10–11-year follow-up assessment and detailed descriptions of the measurements have been previously reported.^{30,31} Blood pressure was measured with the automatic

oscillometric method. For PWV, pressure pulse waveforms were recorded transcutaneously using a high-fidelity micromanometer (SPC-301; Millar Instruments, Houston, TX, USA) from the radial and carotid pulse, synchronous with the ECG signal. Integral software processed the data to calculate the mean time difference between R-wave and pressure wave on a beat-to-beat basis over 10 s, and the PWV was then calculated using the mean time difference and arterial path length between the two recording points (SphygmoConversion 7.1; Scanned, UK). Ultrasound images of the right brachial artery were used to measure FMD and DC. Images were recorded onto SVHS video using an ALOKA 5500 high-resolution ultrasound system with a 5–10 MHz linear array probe (Keymed, UK) and the measurements were undertaken later from the videos at the Vascular Physiology Unit, Institute of Child Health, London. The right brachial artery was imaged 5–10 cm above the antecubital fossa. FMD was induced by a 5-min inflation of a pneumatic cuff, placed around the forearm immediately below the medial epicondyle, to 200 mmHg followed by rapid deflation using an automatic air regulator. The diameter of the brachial artery was measured using edge detection software (Brachial Tools, MIA, IA, USA) from ECG-triggered ultrasound images captured at 3 s intervals throughout the 11-min recording protocol. FMD was expressed as an absolute value in our analyses.³¹ The distension of the artery was determined by measuring the luminal diameter excursion from diastole to systole. Distensibility coefficient was calculated from the distension and the pulse pressure and was expressed as mean per cent change in cross-sectional area per unit change in blood pressure.

Covariates

Maternal age, gestational age, birthweight, and offspring sex were obtained from obstetric records. Self-reported weight and height at the time of recruitment were used to calculate maternal pre-pregnancy body mass index (BMI; Pearson's correlation with measured weight at first antenatal clinic $r^2=0.95$, $p<0.0001$). Parity was obtained from a maternal completed questionnaire at 18 weeks gestation and for these analyses we dichotomized this (any previous pregnancies yes/no). Based on questionnaire responses, the highest parental occupation was used to allocate the children to head of household occupational social class groups (classes I (professional/managerial) to V (unskilled manual workers), using the 1991 British Office of Population and Census Statistics classification). Ethnicity was based on the mothers' self-report (White; Black Caribbean; Black African; Other Black; Indian; Pakistani; Bangladeshi; Chinese; Other).

Mothers were repeatedly asked about their smoking throughout pregnancy and these data were used to generate a categorical variable (never smoked; smoked before pregnancy or in the first trimester and then stopped; smoked throughout pregnancy). A categorical variable on alcohol consumption was defined based on a question in the questionnaire at gestational week 18 (no drinking; less than 1 glass per week; more than 1 glass per week; more than 1 glass per day).

Age of the child at the time of vascular function assessment was recorded in months as they arrived at the assessment clinics. Weight was measured to the nearest 0.1 kg using Tanita scales. Height was measured to the nearest 0.1 cm using a Harpenden stadiometer.

Statistics

Associations between the stress measures (at all time points) and offspring vascular outcome measures were explored using multivariable linear regression with the statistical package SPSS version 18.0 (SPSS, Chicago, USA). In model 1 we adjusted for sex, height, and age of the child at the time of vascular assessment. In model 2 we additionally adjusted for maternal confounders (age, ethnicity, pre-pregnancy BMI, nulliparity, social class, smoking, and alcohol consumption). In model 3 we adjusted for potential mediation of any association by gestational age at birth and birthweight, and in model 4 for potential mediation by BMI of the child at the time of vascular assessment. We explored whether there were any differences in associations by sex of the child by testing for an interaction between sex and the exposure (depressive symptoms or anxiety) in model 2, using the Wald test. The equality of betas of a psychosocial measure at two gestational time points, or between mothers and fathers, was tested using the Wald-statistic. Associations are also checked for linearity using restricted cubic splines within ordinary least squares linear regression analysis (OLS) in R (R Foundation for Statistical Computing).

Results

Table 1 shows the characteristics of the mothers and children included in this study. Table 2 shows the correlations between the maternal depressive symptoms and anxiety scores at all three time points and the fathers' equivalent measurements. There were strong correlations between depressive symptoms and anxiety at each time point in the mothers, and also in the fathers. Over time, in the mothers, these measurements correlated strongly with each other. There were weak to modest correlations between maternal depressive symptoms and anxiety and paternal equivalent outcomes.

Table 1. Participant characteristics

	Study population (<i>n</i> = 43 18)
Maternal	
Age at delivery (years)	29.3 ± 4.4 (26.3–32.3)
White ethnicity (yes)	98.7
Pre-pregnancy body mass index (kg/m ²)	22.8 ± 3.6 (21.0–24.7)
Obesity (body mass index ≥30 kg/m ²) (yes)	4.5
Social class	
I (professional/managerial)	7.1
II	35.9
III (non-manual)	41.7
III (manual)	6.3
IV	7.6
V (unskilled manual workers)	1.4
Primiparous (yes)	49.4
Smoking	
Non-smoking	77.3
Stopped before pregnancy or in first trimester	11.1
Smoked during pregnancy	11.6
Alcohol	
None	43.7
<1 glass/week	41.0
>1 glass/week	13.7
>1 glass/day	1.5
Pregnancy-related hypertension (yes)	15.5
Maternal stress	
Anxiety at gestational week 18 as a continuous score	5 ± 3 (3–7)
Anxiety at gestational week 32 as a continuous score	5 ± 3 (2–7)
Depressive symptoms at gestational week 18 as a continuous score	6 ± 5 (3–9)
Depressive symptoms at gestational week 32 as a continuous score	6 ± 5 (3–10)
Pre- and postnatal anxiety categories (<i>n</i> = 3375) ^a	
Low anxiety at all time points	59.9
Prenatal anxiety only	19.8
Postnatal anxiety only	5.8
High anxiety at all time points	14.5
Pre- and postnatal depressive symptom categories (<i>n</i> = 3421) ^a	
Low depressive symptoms at all time points	61.6
Prenatal depressive symptoms only	15.0
Postnatal depressive symptoms only	8.7
High depressive symptoms at all time points	14.6
Child at birth	
Born through caesarean section (yes)	9.6
Sex (boys)	49.4
Gestational age (weeks)	39.6 ± 1.6 (38.6–40.6)
Premature (<37 weeks) (yes)	3.7
Birthweight (g)	3456 ± 494 (1366–5546)

(continued)

Table 1. Continued

	Study population (n = 4318)
Child at age 10 assessment	
Age (years)	10.7 ± 0.2 (10.5–10.8)
Height (cm)	144 ± 7 (140–149)
Body mass index (kg/m ²)	18.1 ± 3.0 (16.3–20.0)
Systolic blood pressure (mmHg)	104 ± 9 (98–109)
Diastolic blood pressure (mmHg)	60 ± 8 (54–65)
Flow-mediated dilatation (mm) (n = 3830)	0.21 ± 0.08 (0.15–0.27)
Pulse wave velocity (m/s) (n = 4286)	7.55 ± 1.22 (6.75–8.35)
Distensibility coefficient (% per mmHg) (n = 3943)	12.5 ± 6.1 (9.0–16.0)
Brachial artery diameter (mm) (n = 3830)	2.67 ± 0.30 (2.47–2.86)

Values are mean ± SD (IQR) or %; ^aLow anxiety/depressive symptoms both pre- and postnatally: all scores for the respective exposure (anxiety or depressive symptoms) lower than the 80th percentile. Prenatal anxiety/depressive symptoms: one or both of the prenatal scores at or higher than the 80th percentile, but the postnatal score lower than the 80th percentile. Postnatal anxiety/depressive symptoms: both prenatal scores lower than the 80th percentile and the postnatal score at or higher than the 80th percentile. High anxiety/depressive symptoms: all scores at or higher than the 80th percentile. 80th percentile scores: Anxiety: 7 at gestational week 18; 7 at gestational week 32; 6 at baby age 8 months. Depressive symptoms: 10; 11; 8 respectively.

Table 2. Correlation coefficients between maternal depressive symptoms and anxiety at gestational weeks 18 and 32, and at 8 months postnatally, and paternal depressive symptoms and anxiety at partner's gestational week 19

	18 weeks		32 weeks		Postnatal		Paternal	
	EPDS	CCEI	EPDS	CCEI	EPDS	CCEI	EPDS	CCEI
EPDS depressive symptoms at week 18	1							
	n = 4318							
CCEI anxiety at week 18	0.74	1						
	n = 4318	n = 4318						
EPDS depressive symptoms at week 32	0.64	0.57	1					
	n = 4318	n = 4318	n = 4318					
CCEI anxiety at week 32	0.60	0.67	0.76	1				
	n = 4318	n = 4318	n = 4318	n = 4318				
EPDS depressive symptoms 8 months postnatal	0.50	0.46	0.54	0.50	1			
	n = 4173	n = 4173	n = 4173	n = 4173	n = 4173			
CCEI anxiety 8 months postnatal	0.47	0.53	0.49	0.56	0.76	1		
	n = 4181	n = 4181	n = 4181	n = 4181	n = 4142	n = 4181		
Paternal EPDS depressive symptoms	0.24	0.19	0.20	0.18	0.17	0.16	1	
	n = 3589	n = 3589	n = 3589	n = 3589	n = 3475	n = 3484	n = 3589	
Paternal CCEI anxiety	0.15	0.14	0.13	0.11	0.12	0.12	0.68	1
	n = 3576	n = 3576	n = 3576	n = 3576	n = 3463	n = 3473	n = 3545	n = 3576

All *p*-values are <0.001; CCEI, Crown–Crisp experiential index; EPDS, Edinburgh Postnatal Depression Scale.

There was no strong statistical evidence of interaction by sex and therefore all multivariable association results are presented with both females and males combined. Furthermore, there was no evidence for non-linear associations: we therefore present results examining linear associations.

At both gestational ages, there was no strong evidence that maternal depressive symptoms or anxiety were associated with adverse cardiovascular function in any of the multivariable models (Table 3). Maternal depressive symptoms in pregnancy at both time points were associated with larger brachial artery

Table 3. Multivariable associations of maternal depressive symptoms and anxiety at two gestational time points with offspring vascular phenotypes at mean age 10.7 years (n = 4318)

	Systolic blood pressure		Diastolic blood pressure		Flow-mediated dilatation		Pulse wave velocity		Distensibility coefficient		Brachial artery diameter							
	95% CI		95% CI		95% CI		95% CI		95% CI		95% CI							
	β	Upper	β	Upper	β	Upper	β	Upper	β	Upper	β	Upper						
Depressive symptoms																		
Gestational week 18																		
Model 1	-0.038	-0.096	0.021	-0.074*	-0.126	-0.022	0.000	-0.001	0.001	-0.004	-0.012	0.004	-0.011	-0.053	0.031	0.003*	0.001	0.005
Model 2	-0.051	-0.111	0.008	-0.083*	-0.136	-0.03	0.000	-0.001	0.001	-0.005	-0.013	0.004	-0.014	-0.057	0.029	0.002*	0.000	0.004
Model 3	-0.051	-0.110	0.009	-0.082*	-0.135	-0.029	0.000	-0.001	0.001	-0.005	-0.013	0.004	-0.014	-0.057	0.029	0.002*	0.000	0.004
Model 4	-0.053	-0.112	0.006	-0.084*	-0.136	-0.031	0.000	-0.001	0.000	-0.004	-0.012	0.004	-0.014	-0.058	0.029	0.002*	0.000	0.003
Gestational week 32																		
Model 1	-0.039	-0.095	0.016	-0.022	-0.072	0.027	0.000	0.000	0.001	-0.002	-0.010	0.006	0.012	-0.028	0.051	0.003*	0.001	0.004
Model 2	-0.052	-0.108	0.004	-0.029	-0.08	0.021	0.000	0.000	0.001	-0.003	-0.010	0.005	0.009	-0.032	0.050	0.002*	0.000	0.004
Model 3	-0.052	-0.108	0.004	-0.028	-0.078	0.022	0.000	0.000	0.001	-0.002	-0.010	0.005	0.009	-0.032	0.049	0.002	0.000	0.004
Model 4	-0.058	-0.113	-0.002	-0.032	-0.082	0.018	0.000	0.000	0.001	-0.001	-0.009	0.006	0.009	-0.032	0.049	0.001	-0.001	0.003
Wald-statistic (model 1)	p = 0.89		p = 0.02		p = 0.29		p = 0.67		p = 0.20		p = 0.94							
Anxiety																		
Gestational week 18																		
Model 1	-0.029	-0.108	0.049	-0.034	-0.104	0.036	0.000	-0.001	0.000	-0.002	-0.013	0.009	0.014	-0.043	0.070	0.004*	0.001	0.006
Model 2	-0.046	-0.126	0.034	-0.043	-0.114	0.028	-0.001	-0.001	0.000	-0.003	-0.014	0.008	0.013	-0.045	0.070	0.003*	0.000	0.006
Model 3	-0.044	-0.123	0.036	-0.041	-0.112	0.030	-0.001	-0.001	0.000	-0.003	-0.014	0.008	0.013	-0.045	0.070	0.003*	0.000	0.005
Model 4	-0.048	-0.127	0.031	-0.044	-0.115	0.027	-0.001	-0.001	0.000	-0.003	-0.013	0.008	0.012	-0.045	0.070	0.002	0.000	0.005
Gestational week 32																		
Model 1	-0.018	-0.095	0.059	-0.040	-0.109	0.028	0.000	0.000	0.001	0.003	-0.008	0.013	0.024	-0.032	0.080	0.003*	0.000	0.005
Model 2	-0.034	-0.112	0.044	-0.048	-0.118	0.022	0.000	0.000	0.001	0.002	-0.009	0.013	0.022	-0.035	0.078	0.002	0.000	0.005
Model 3	-0.030	-0.108	0.048	-0.045	-0.115	0.024	0.000	-0.001	0.001	0.002	-0.009	0.013	0.022	-0.035	0.078	0.002	0.000	0.005
Model 4	-0.038	-0.116	0.039	-0.051	-0.120	0.019	0.000	-0.001	0.001	0.003	-0.007	0.014	0.021	-0.035	0.078	0.001	-0.001	0.004
Wald-statistic (model 1)	p = 0.74		p = 0.81		p = 0.01		p = 0.30		p = 0.65		p = 0.53							

Model 1: Adjusted for offspring sex, height, and age at time of outcome measurement; Model 2: Maternal confounder-adjusted model: as model 1 plus adjustment for maternal age, ethnicity, pre-pregnancy BMI, nulliparity, social class, smoking, alcohol consumption; Model 3: Birth outcome-adjusted model: as model 2 plus adjustment for gestational age and birthweight; Model 4: Offspring adiposity-adjusted model: as model 3 plus adjustment for BMI child; *p < 0.05; The Wald-statistic tests the equality of the effects of depressive symptoms between week 18 and 32 in model 1.

diameter in the confounder-adjusted model (model 2), but with some evidence that these associations might be mediated by birthweight/gestational age and offspring adiposity at the time of assessment of vascular function (models 3 and 4). Depressive symptoms at 18 weeks gestation were also associated with lower diastolic blood pressure. Maternal anxiety at 18 weeks gestation was also associated with brachial artery diameter (model 2), again with some evidence of mediation by birthweight/gestational age and offspring adiposity.

Table 4 compares the multivariable associations of maternal depressive symptoms with offspring vascular outcomes to the equivalent associations of paternal symptoms. Paternal depressive symptoms or anxiety were not associated with any of the vascular outcomes in any models. For almost all associations, there was statistical evidence that the paternal–offspring associations were consistent with the maternal–offspring associations. There was some statistical evidence that the inverse association of maternal stress with offspring DBP was different to the paternal null association (Wald-statistic $p = 0.04$).

Table 5 shows the multivariable associations of postnatal maternal depressive symptoms and anxiety with offspring vascular function. Consistent with the findings of a lack of associations at each time point with offspring outcomes and of the strong correlations between symptoms at each time point, maternal postnatal symptoms were not strongly associated with overall adverse offspring vascular outcomes. There was some evidence that offspring of mothers with postnatal depressive symptoms or anxiety had a lower SBP and DBP. These associations were not explained by gestational duration, birthweight, or offspring age at vascular assessment.

Discussion

We have found no convincing evidence that prenatal psychosocial stress, in the form of maternal depressive symptoms or anxiety, causes adverse levels of offspring blood pressure, endothelial function, arterial stiffness, brachial artery distensibility, or brachial diameter assessed at age 10–11 years via intrauterine mechanisms. Indeed for most outcomes, there was no association. Furthermore, the point estimates for the maternal–offspring association were very similar to those for the paternal–offspring associations, providing further evidence against the hypothesis that exposure to maternal stress *in utero* causes adverse developmental effects on vascular structure/function. Maternal depressive symptoms and anxiety in the postnatal period were associated with lower blood pressure in the offspring (i.e. in the opposite direction to what might be expected), though the magnitudes of association were

small. As postnatal stress was not the prime focus of this paper, this could be a chance finding and requires further replication and detailed assessment to ascertain whether this is robust.

Our main findings seem to be in line with the only previous study of a somewhat similar association in a general population cohort that we are aware of, in which we did not find an association between prenatal maternal depressive symptoms and anxiety with offspring blood pressure in Dutch children at age 5–7.²¹ However, that previous study did report an association between the accumulation of multiple prenatal maternal stressors and higher offspring systolic and diastolic blood pressure. In the current study, we do not have a measure of the accumulation of multiple psychosocial stressors available. It is possible that the accumulation of different stressors can exert a programming effect on the offspring, which is not identified in a study such as this that assesses just two sources of stress (depression/anxiety) that are highly related to each other. Maternal cortisol early in pregnancy appears to be associated with both increased vascular resistance and lower arterial elasticity in children aged 5–7.^{32,33} Possibly, the stressors in the current study are not causing sufficient rises in maternal cortisol concentrations to exert effects on offspring arterial parameters. Of note the evidence for a direct association between psychosocial stress and cortisol, and thus the role of cortisol in fetal programming via psychosocial stress, has been a main point of discussion in a recent review by Glover et al.⁶ Nonetheless our results are important as they strongly suggest that levels of maternal stress in pregnancy that are experienced by a relatively large proportion of pregnant women are unlikely to adversely affect their offspring's vascular health.

It is possible that there are specific sensitive periods in which maternal stress has an effect on future cardiovascular function.³⁴ However, the clear lack of associations at either 18 weeks or 32 weeks and the correlation between symptoms across these two time points and with the postnatal measurement (suggesting women would be ranked similarly throughout pregnancy) argues against this as an explanation for our null results.

The associations of maternal symptoms with wider brachial diameter and lower diastolic blood pressure were in the opposite direction to what we hypothesized and require further replication before one could consider these are not due to chance.

The large sample size is a strength of the current study, but as in most cohort studies, selective loss to follow up was present. Those who have remained in the cohort tend to be of higher socioeconomic position^{24,25} and we have shown that those who were included in the analyses here had lower parental depressive and anxiety

Table 4. Comparison of multivariable associations of maternal and paternal depressive symptoms ($n = 3589$) and anxiety ($n = 3576$) at partner's gestational week 18 with offspring vascular phenotypes at mean age 10.7 years

	Systolic blood pressure		Diastolic blood pressure		Flow-mediated dilatation		Pulse wave velocity		Distensibility coefficient		Brachial artery diameter	
	β	95% CI	β	95% CI	β	95% CI	β	95% CI	β	95% CI	β	95% CI
Paternal depressive symptoms												
Model 1	-0.04	-0.12 0.04	-0.02	-0.08 0.05	0.000	-0.001 0.000	0.00	-0.01 0.01	0.01	-0.05 0.06	0.002	-0.001 0.004
Model 2	-0.04	-0.12 0.04	-0.02	-0.09 0.05	0.000	-0.001 0.000	-0.01	-0.02 0.01	0.01	-0.05 0.06	0.002	-0.001 0.004
Model 3	-0.04	-0.12 0.04	-0.02	-0.09 0.05	0.000	-0.001 0.000	-0.01	-0.02 0.01	0.01	-0.05 0.06	0.002	-0.001 0.004
Model 4	-0.04	-0.12 0.03	-0.02	-0.09 0.05	0.000	-0.001 0.000	0.00	-0.01 0.01	0.01	-0.05 0.06	0.001	-0.001 0.004
Maternal depressive symptoms												
Model 1	-0.05	-0.11 0.02	-0.10*	-0.16 -0.04	0.000	-0.001 0.001	-0.01	-0.02 0.00	-0.01	-0.06 0.03	0.003*	0.001 0.005
Wald-statistic	$p = 0.94$		$p = 0.04$		$p = 0.41$		$p = 0.70$		$p = 0.56$		$p = 0.23$	
Paternal anxiety												
Model 1	-0.04	-0.15 0.07	-0.04	-0.13 0.06	0.000	-0.001 0.001	-0.01	-0.02 0.01	0.01	-0.07 0.08	0.002	-0.002 0.005
Model 2	-0.02	-0.12 0.09	-0.03	-0.12 0.07	0.000	-0.001 0.001	-0.01	-0.02 0.01	0.00	-0.08 0.08	0.002	-0.001 0.006
Model 3	-0.01	-0.12 0.10	-0.03	-0.12 0.07	0.000	-0.001 0.001	-0.01	-0.02 0.01	0.00	-0.08 0.08	0.002	-0.001 0.006
Model 4	-0.02	-0.13 0.09	-0.03	-0.13 0.07	0.000	0.000 0.000	-0.01	-0.02 0.01	0.00	-0.08 0.08	0.002	-0.002 0.005
Maternal anxiety												
Model 1	-0.03	-0.12 0.06	-0.06	-0.13 0.02	0.000	-0.001 0.000	0.00	-0.02 0.01	0.02	-0.05 0.08	0.004*	0.001 0.007
Wald-statistic	$p = 0.80$		$p = 0.98$		$p = 0.68$		$p = 0.88$		$p = 0.77$		$p = 0.18$	

Model 1: Adjusted for offspring sex, height, and age at time of outcome measurement; Model 2: Maternal confounder-adjusted model: as model 1 plus adjustment for maternal age, ethnicity, pre-pregnancy BMI, nulliparity, social class, smoking, alcohol consumption; Model 3: Birth outcome-adjusted model: as model 2 plus adjustment for gestational age and birthweight; Model 4: Offspring adiposity-adjusted model: as model 3 plus adjustment for BMI child; * $p < 0.05$; The Wald-statistic tests the equality of the effects of depressive symptoms between mother and father in model 1.

Table 5. Multivariable associations of maternal postnatal depressive symptoms ($n = 3421$) and anxiety ($n = 3375$) with offspring vascular phenotypes at mean age 10.7 years

	Systolic blood pressure		Diastolic blood pressure		Flow-mediated dilatation		Pulse wave velocity		Distensibility coefficient		Brachial artery diameter						
	95% CI		95% CI		95% CI		95% CI		95% CI		95% CI						
	β	Upper	β	Upper	β	Upper	β	Upper	β	Upper	β	Upper					
Depressive symptoms																	
Model 1	-0.072*	-0.134	-0.011	-0.080*	-0.135	-0.025	0.000	0.001	-0.002	-0.010	0.007	-0.006	-0.051	0.038	0.002	0.000	0.004
Model 2	-0.070*	-0.132	-0.008	-0.080*	-0.135	-0.025	0.000	0.001	-0.003	-0.011	0.006	-0.010	-0.055	0.035	0.001	-0.001	0.003
Model 3	-0.071*	-0.133	-0.009	-0.080*	-0.135	-0.025	0.000	0.001	-0.003	-0.011	0.006	-0.010	-0.055	0.034	0.001	-0.001	0.003
Model 4	-0.080*	-0.141	-0.018	-0.086*	-0.141	-0.031	0.000	0.001	-0.001	-0.010	0.007	-0.011	-0.056	0.034	0.001	-0.001	0.002
Anxiety																	
Model 1	-0.096*	-0.181	-0.011	-0.114*	-0.191	-0.038	0.001	-0.001	-0.004	-0.016	0.007	-0.001	-0.063	0.060	0.002	-0.001	0.005
Model 2	-0.084	-0.170	0.002	-0.109*	-0.186	-0.032	0.000	-0.001	-0.006	-0.018	0.006	-0.007	-0.070	0.055	0.002	-0.001	0.005
Model 3	-0.086	-0.171	0.000	-0.109*	-0.186	-0.033	0.000	-0.001	-0.006	-0.018	0.006	-0.007	-0.070	0.055	0.002	-0.001	0.005
Model 4	-0.094*	-0.179	-0.009	-0.115	-0.191	-0.038	0.000	-0.001	-0.005	-0.016	0.007	-0.008	-0.070	0.055	0.001	-0.002	0.004

Model 1: Adjusted for offspring sex, height, and age at time of outcome measurement; Model 2: Maternal confounder-adjusted model: as model 1 plus adjustment for maternal age, ethnicity, pre-pregnancy BMI, nulliparity, social class, smoking, alcohol consumption; Model 3: Birth outcome-adjusted model: as model 2 plus adjustment for gestational age and birthweight; Model 4: Offspring adiposity-adjusted model: as model 3 plus adjustment for BMI child; * $p < 0.05$.

scores than those who were excluded because of missing outcome or covariable data (see supplementary table, available online). If the association of parental symptoms with offspring outcomes was strong and inverse in those lost to follow up, then we may have biased null findings due to selection bias, but in separate studies exploring selection bias due to loss to follow up/missing data any bias tends to be small^{35,36} and we doubt this would be sufficient to produce important associations here.

Whilst the parental symptoms that we have assessed here use validated tools, they are not the same as clinical diagnoses and we cannot make any conclusions about the effects of more severe clinical depression or anxiety in terms of its association with offspring vascular function.

In conclusion, our findings suggest that maternal prenatal psychosocial stress does not cause adverse offspring vascular function in 10–11 year old children via intrauterine mechanisms. These results do not exclude the possibility that severe prenatal clinical depression or anxiety cause adverse offspring vascular function via intrauterine mechanisms, but to date we are not aware of any study that has explored that possibility in humans.

Acknowledgements

We are grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses.

Funding

The Wellcome Trust, UK Medical Research Council and the University of Bristol provide core-funding support for ALSPAC; Vascular measures were funded by a British Heart Foundation programme grant (RG/20011002). AEVD was supported by the Dutch Heart Foundation (DHF-2007B103). DAL works in a centre that receives core funds from the University of Bristol and the UK Medical Research Council (RD1634). These parties played no part in the collection, analysis, interpretation or reporting of the data, nor did they play a part in the decision to submit this manuscript for publication.

Conflict of interest

The authors declare that there is no conflict of interest.

References

1. Roberts D and Dalziel S. Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. *Cochrane Database Syst Rev* 2006; (3): CD004454.

2. Berry LM, Polk DH, Ikegami M, et al. Preterm newborn lamb renal and cardiovascular responses after fetal or maternal antenatal betamethasone. *Am J Physiol* 1997; 272(6 Pt 2): R1972–R1979.
3. Docherty CC, Kalmár-Nagy J, Engelen M, et al. Effect of in vivo fetal infusion of dexamethasone at 0.75 GA on fetal ovine resistance artery responses to ET-1. *Am J Physiol Regul Integr Comp Physiol* 2001; 281: R261–R268.
4. Dodic M, Abouantoun T, O'Connor A, et al. Programming effects of short prenatal exposure to dexamethasone in sheep. *Hypertension* 2002; 40: 729–734.
5. Entringer S, Kumsta R, Hellhammer DH, et al. Prenatal exposure to maternal psychosocial stress and HPA axis regulation in young adults. *Horm Behav* 2009; 55: 292–298.
6. Glover V, O'Connor TG and O'Donnell K. Prenatal stress and the programming of the HPA axis. *Neurosci Biobehav Rev* 2010; 35: 17–22.
7. Glynn LM, Wadhwa PD, Dunkel-Schetter C, et al. When stress happens matters: effects of earthquake timing on stress responsivity in pregnancy. *Am J Obstet Gynecol* 2001; 184: 637–642.
8. Levitt NS, Lindsay RS, Holmes MC, et al. Dexamethasone in the last week of pregnancy attenuates hippocampal glucocorticoid receptor gene expression and elevates blood pressure in the adult offspring in the rat. *Neuroendocrinology* 1996; 64: 412–418.
9. McCormick CM, Smythe JW, Sharma S, et al. Sex-specific effects of prenatal stress on hypothalamic-pituitary-adrenal responses to stress and brain glucocorticoid receptor density in adult rats. *Brain Res Dev Brain Res* 1995; 84: 55–61.
10. Molnar J, Nijland MJ, Howe DC, et al. Evidence for microvascular dysfunction after prenatal dexamethasone at 0.7, 0.75, and 0.8 gestation in sheep. *Am J Physiol Regul Integr Comp Physiol* 2002; 283: R561–R567.
11. Molnar J, Howe DC, Nijland MJ, et al. Prenatal dexamethasone leads to both endothelial dysfunction and vasodilatory compensation in sheep. *J Physiol* 2003; 547(Pt 1): 61–66.
12. Moritz KM, Johnson K, Douglas-Denton R, et al. Maternal glucocorticoid treatment programs alterations in the renin-angiotensin system of the ovine fetal kidney. *Endocrinology* 2002; 143: 4455–4463.
13. Moss TJ, Sloboda DM, Gurrin LC, et al. Programming effects in sheep of prenatal growth restriction and glucocorticoid exposure. *Am J Physiol Regul Integr Comp Physiol* 2001; 281: R960–R970.
14. Noorlander CW, De Graan PN, Middeldorp J, et al. Ontogeny of hippocampal corticosteroid receptors: effects of antenatal glucocorticoids in human and mouse. *J Comp Neurol* 2006; 499: 924–932.
15. O'Donnell K, O'Connor TG and Glover V. Prenatal stress and neurodevelopment of the child: focus on the HPA axis and role of the placenta. *Dev Neurosci* 2009; 31: 285–292.
16. Ortiz LA, Quan A, Zarzar F, et al. Prenatal dexamethasone programs hypertension and renal injury in the rat. *Hypertension* 2003; 41: 328–334.
17. Ortiz LA, Quan A, Weinberg A, et al. Effect of prenatal dexamethasone on rat renal development. *Kidney Int* 2001; 59: 1663–1669.
18. Drake AJ, Tang JI and Nyirenda MJ. Mechanisms underlying the role of glucocorticoids in the early life programming of adult disease. *Clin Sci (Lond)* 2007; 113: 219–232.
19. Kari MA, Hallman M, Eronen M, et al. Prenatal dexamethasone treatment in conjunction with rescue therapy of human surfactant: a randomized placebo-controlled multicenter study. *Pediatrics* 1994; 93: 730–736.
20. Doyle LW, Ford GW, Davis NM, et al. Antenatal corticosteroid therapy and blood pressure at 14 years of age in preterm children. *Clin Sci (Lond)* 2000; 98: 137–142.
21. van Dijk AE, van Eijdsden M, Stronks K, et al. The association between prenatal psychosocial stress and blood pressure in the child at age 5–7 years. *PLoS ONE* 2012; 7: e43548.
22. Lipsitch M, Tchetgen TE and Cohen T. Negative controls: a tool for detecting confounding and bias in observational studies. *Epidemiology* 2010; 21: 383–388.
23. Davey Smith G. Negative control exposures in epidemiologic studies. *Epidemiology* 2012; 23: 350–351.
24. Boyd A, Golding J, Macleod J, et al. Cohort Profile: The 'Children of the 90s' – the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol* 2013; 42: 111–127.
25. Fraser A, Macdonald-Wallis C, Tilling K, et al. Cohort Profile: The Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol* 2013; 42: 97–110.
26. Williams DM, Fraser A, Sayers A, et al. Associations of childhood 25-hydroxyvitamin D2 and D3 and cardiovascular risk factors in adolescence: prospective findings from the Avon Longitudinal Study of Parents and Children. *Eur J Prev Cardiol* 2012 (Epub ahead of print).
27. Cox JL, Holden JM and Sagovsky R. Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatry* 1987; 150: 782–786.
28. Heron J, O'Connor TG, Evans J, et al. The course of anxiety and depression through pregnancy and the postpartum in a community sample. *J Affect Disord* 2004; 80: 65–73.
29. Crisp AH, Jones MG and Slater P. The Middlesex Hospital Questionnaire: a validity study. *Br J Med Psychol* 1978; 51: 269–280.
30. Donald AE, Charakida M, Falaschetti E, et al. Determinants of vascular phenotype in a large childhood population: the Avon Longitudinal Study of Parents and Children (ALSPAC). *Eur Heart J* 2010; 31: 1502–1510.
31. Lawlor DA, Macdonald-Wallis C, Fraser A, et al. Cardiovascular biomarkers and vascular function during childhood in the offspring of mothers with hypertensive disorders of pregnancy: findings from the Avon Longitudinal Study of Parents and Children. *Eur Heart J* 2012; 33: 335–345.
32. Rondo PH, Lemos JO, Pereira JA, et al. The relationship between cortisol concentrations in pregnancy and

- systemic vascular resistance in childhood. *Early Hum Dev* 2010; 86: 127–131.
33. Rondo PHC, Pereira JA, Lemos JO, et al. The impact of maternal cortisol concentrations on child arterial elasticity. *J Dev Origin Health Dis* 2011; 2: 56–62.
 34. Wadhwa PD and Federenko IS. Prenatal stress influences human fetal development and birth outcomes: implications for developmental origins of health and disease. In: Hodgson DM and Coe CL (eds) *Perinatal programming. Early life determinants of adult health and disease*. London and New York: Taylor & Francis, 2006, pp.29–46.
 35. Wolke D, Waylen A, Samara M, et al. Selective drop-out in longitudinal studies and non-biased prediction of behaviour disorders. *Br J Psychiatry* 2009; 195: 249–256.
 36. Howe LD, Tilling K, Galobardes B, et al. Loss to follow-up in cohort studies: bias in estimates of socioeconomic inequalities. *Epidemiology* 2013; 24: 1–9.