



**Looking but not seeing: recent perspectives on Posterior Cortical Atrophy (PCA)**

Journal:	<i>Current Directions in Psychological Science</i>
Manuscript ID	CDPS-15-0226.R2
Manuscript Type:	Invited Manuscript
Date Submitted by the Author:	n/a
Complete List of Authors:	Crutch, Sebastian; University College London, Dementia Research Centre, UCL Institute of Neurology Yong, Keir; University College London, Dementia Research Centre, UCL Institute of Neurology Shakespeare, Timothy; University College London, Dementia Research Centre, UCL Institute of Neurology
Keywords:	Posterior cortical atrophy, Alzheimer's disease, vision, oculomotor, crowding, dyslexia
Abstract:	<p>Posterior cortical atrophy (PCA) is the canonical 'visual dementia', with affected individuals experiencing a progressive disintegration of their visual world owing to dysfunction and atrophy at the back of the brain. The syndrome, which also affects literacy, numeracy and gesture, is typically caused by Alzheimer's disease (AD), but this atypical form of AD is distinguished from more common amnesic presentations by virtue of relatively preserved episodic memory and insight. Although problems with object and space perception are the most widely reported and investigated symptoms, these higher order perceptual difficulties are often underpinned by an array of changes in more basic visual and oculomotor processes. Here we review recent studies providing insights into these more elementary aspects of vision in PCA, including fixation stability, saccade generation, point localization, excessive crowding, and factors affecting the effective field of vision. It is argued that a more detailed appreciation of these fundamental changes in the early visual system will not only improve the characterization and understanding of this rare clinico-radiological syndrome, but will also guide the design of visual aids and strategies aimed at maintaining everyday abilities in individuals with PCA.</p>

SCHOLARONE™  
Manuscripts

1  
2  
3 Current Directions in Psychological Science  
4  
5  
6

7 **Looking but not seeing: recent perspectives on Posterior Cortical Atrophy**  
8 **(PCA)**  
9

10  
11  
12  
13  
14 Sebastian J. Crutch, Keir X. X. Yong & Timothy J. Shakespeare  
15  
16

17  
18  
19 Dementia Research Centre, Department of Neurodegeneration, Institute of  
20  
21 Neurology, University College London, UK  
22  
23

24  
25  
26 Address for correspondence:  
27  
28

29  
30 Sebastian Crutch  
31

32 Dementia Research Centre  
33

34 Box 16  
35

36 National Hospital for Neurology and Neurosurgery  
37

38 Queen Square  
39

40 London  
41

42 WC1N 3BG  
43

44 UK  
45  
46

47  
48  
49  
50 Tel: +44 (0)20 3448 3113  
51

52 Fax: +44 (0)20 7676 2066  
53  
54

55  
56  
57 Email: [s.crutch@ucl.ac.uk](mailto:s.crutch@ucl.ac.uk)  
58  
59  
60

**Abstract**

Posterior cortical atrophy (PCA) is the canonical 'visual dementia', with affected individuals experiencing a progressive disintegration of their visual world owing to dysfunction and atrophy at the back of the brain. The syndrome, which also affects literacy, numeracy and gesture, is typically caused by Alzheimer's disease (AD), but this atypical form of AD is distinguished from more common amnesic presentations by virtue of relatively preserved episodic memory and insight.

Although problems with object and space perception are the most widely reported and investigated symptoms, these higher order perceptual difficulties are often underpinned by an array of changes in more basic visual and oculomotor processes. Here we review recent studies providing insights into these more elementary aspects of vision in PCA, including fixation stability, saccade generation, point localization, excessive crowding, and factors affecting the effective field of vision. It is argued that a more detailed appreciation of these fundamental changes in the early visual system will not only improve the characterization and understanding of this rare clinico-radiological syndrome, but will also guide the design of visual aids and strategies aimed at maintaining everyday abilities in individuals with PCA.

**Keywords:** Posterior cortical atrophy; Alzheimer's disease; vision; oculomotor; crowding; dyslexia

1  
2  
3 Posterior cortical atrophy (PCA) is a clinico-radiological syndrome characterized  
4  
5 by insidious decline primarily in visuo-perceptual and visuo-spatial processing  
6  
7 and dysfunction or atrophy of the posterior cortices (Benson et al., 1988; Mendez  
8  
9 et al., 2002; Tang-Wai et al., 2004; see Crutch et al., 2012 for a review). Common  
10  
11 presenting symptoms include difficulties with driving, reading, typing and other  
12  
13 tasks which involve interpreting, locating, and/or reaching for objects under  
14  
15 visual guidance, with additional problems reading, writing, spelling, numeracy  
16  
17 and praxis. By contrast, episodic memory and insight are relatively spared. These  
18  
19 symptoms are typically associated with hypometabolism and atrophy  
20  
21 particularly in parietal, occipital and occipito-temporal cortex (see Figure 1).  
22  
23  
24

25  
26  
27  
28 The most common pathology in PCA is Alzheimer's disease (AD), with PCA  
29  
30 constituting a common form of atypical AD (McKhann et al., 2011; Dubois et al.,  
31  
32 2014). A small proportion of cases are due to Lewy body disease and  
33  
34 corticobasal degeneration (Hof et al., 1990; Renner et al., 2004). In those PCA  
35  
36 patients with Alzheimer's pathology, the distribution of pathological changes  
37  
38 differs from that in typical AD, with lower plaque and tangle counts in the medial  
39  
40 temporal regions and frontal cortices and relatively higher counts in posterior  
41  
42 cortices such as the occipital lobes (Hof et al., 1990; Tang-Wai et al., 2004).  
43  
44

45  
46 However on molecular imaging the focal posterior pattern of cortical  
47  
48 involvement is evident with tau-imaging but less clear cut with amyloid imaging  
49  
50 where results indicate a more global pattern of amyloid deposition  
51  
52 (Ossenkoppele et al., 2015; see Figure 1b). These findings suggest that tau is  
53  
54 more closely linked to symptomatology than amyloid. PCA is typically a younger-  
55  
56  
57  
58  
59  
60

onset, sporadic condition (age 50–65 years) (Mendez et al., 2002; Tang-Wai et al., 2004; McMonagle et al., 2006).




Figure 1. (A) Differences in gray matter volume between individuals with PCA and healthy controls revealed by voxel-based morphometry. Yellow overlay represents  $T$  scores in areas of statistically significant lower gray matter volume in PCA (FWE corrected at  $P < 0.05$ ). Images shown in neurological convention (right on right). Crosshairs and co-ordinates (in MNI space) indicate  $T$  score global maxima. Adapted from Ryan, Shakespeare et al. (2014). (B) Axial slices of fibrillary amyloid-beta ([ $^{11}\text{C}$ ] PIB: Pittsburgh compound B), regional tau load ([ $^{18}\text{F}$ ]AV-1451), and glucose metabolism ([ $^{18}\text{F}$ ]fluorodeoxyglucose (FDG) positron emission tomography) alongside structural magnetic resonance imaging (MRI) scan in a single patient with PCA. [ $^{11}\text{C}$ ]PIB binds throughout association neocortex whilst

1  
2  
3 *[(18) F]AV-1451 is selectively retained in posterior brain regions that were affected*  
4 *clinically and which show reduced [(18) F]FDG uptake. DVR = distribution volume*  
5 *ratio. SUVR = standardized uptake value ratio. Adapted from Ossenkoppele et al.*  
6 *(2015).*  
7  
8  
9  
10

11  
12  
13  
14 The current paper focuses upon studies directly examining aspects of basic  
15 visual and oculomotor function in PCA. Clinical studies have tended to examine  
16 high-level object and space perception problems (e.g. the perception of single or  
17 multiple whole objects, faces or words, of which patients complain directly)  
18 which rely on cognitive processes associated with parietal and occipito-temporal  
19 mechanisms downstream in the visual system. By contrast, more basic visual  
20 functions (e.g. edge detection, form and motion coherence) which may underpin  
21 many such downstream deficits and which are mediated largely by upstream  
22 occipital mechanisms, have been largely overlooked. Understanding these basic  
23 visual aspects is of interest not just to better characterize the syndrome, but also  
24 in order to develop a better understanding of the underpinnings of visual  
25 impairments in PCA and develop rehabilitative or compensatory strategies.  
26 Furthermore, a sound understanding of the basic visual and oculomotor deficits  
27 present in these patients is critical for the design and interpretation of clinical  
28 and neuroscientific studies involving individuals with PCA.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 Here we consider the insights provided by recent studies of fixation stability,  
51 saccade generation and smooth pursuit aspects of oculomotor function, and  
52 point localization, visual crowding, and factors affecting the effective field of  
53 vision. We also consider current translational research informed by these  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 insights including attempts to facilitate reading and room navigation in PCA. The  
4  
5 paper concludes with discussion of three avenues for future research illustrating  
6  
7 aspects of the syndrome that extend beyond visual deficits, namely visuo-  
8  
9 vestibular interactions, motor deficits and representation of concepts related to  
10  
11 quantity.  
12  
13



### 14 15 16 17 **Oculomotor function**

18  
19  
20 A recent systematic evaluation of oculomotor function has shown that eye  
21  
22 movement abnormalities are near-ubiquitous in PCA (Shakespeare et al., 2015a).  
23  
24 Using tests of fixation, saccade (fixation/target gap and overlap conditions) and  
25  
26 smooth pursuit eye movements, eye movement abnormalities were detected in  
27  
28 80% of PCA patients compared to 17% of typical Alzheimer's disease (tAD) and  
29  
30 5% of controls. On fixation stability tasks, PCA patients exhibit micro-saccadic  
31  
32 and large saccadic intrusions, which may account for experience of apparent  
33  
34 motion or unsteadiness amongst static objects reported by many PCA patients  
35  
36 (Crutch et al., 2011). Saccade generation was particularly affected, with PCA  
37  
38 patients making significantly shorter saccades especially for distant targets. They  
39  
40 also exhibited a significant exacerbation of the normal gap/overlap effect,  
41  
42 consistent with 'sticky fixation' (see Figure 2a). PCA patients also showed lower  
43  
44 gain in smooth pursuit than controls (see Figure 2b). These oculomotor deficits  
45  
46 may reflect weak input from degraded occipito-parietal spatial representations  
47  
48 of stimulus location into a superior collicular spatial map for eye movement  
49  
50 regulation. In other words, in PCA it may be the identification of oculomotor  
51  
52 targets rather than the generation of oculomotor movements which is  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

particularly problematic. If this hypothesis is correct, precise stimulus localization and spatial attention may play a particularly critical role in determining the adequacy of real world perception in everyday life.

For Peer Review



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5 *Figure 2. Abnormalities of fixation and pursuit in PCA (figures adapted from*  
6 *Shakespeare et al., 2015a). (A) Representative traces from the saccade task for a*  
7 *healthy control, a typical AD (tAD) patient and a PCA patient in an ‘overlap’ trial.*  
8 *The upper plot (grey line) for each participant shows gaze position in the y*  
9 *(vertical) axis, the lower plot (black line) shows gaze position in the x (horizontal)*  
10 *axis. Gridlines show displacement of 1° of visual angle. Positive values of gaze*  
11 *position indicate rightward gaze. A central fixation point was present from the*  
12 *start of the trial until timepoint B (500ms). The target appeared at 10°*  
13 *horizontally to the right of the central fixation point at timepoint A (300ms) and*  
14 *remained present until the end of the trial. The healthy control and tAD patients*  
15 *make a single saccade towards the target. The PCA patient takes a long time to*  
16 *initiate their first saccade (in the incorrect direction), followed by a number of*  
17 *small saccades to reach the target location. (B) Example traces from the pursuit*  
18 *task. The figure shows a cycle towards the middle of the trial (seconds 4–8 from a*  
19 *trial of 10 s). Positive values of gaze position indicate rightward gaze. The upper*  
20 *plot (grey line) for each participant shows gaze position in the y (vertical) axis, the*  
21 *lower plot (black line) shows gaze position in the x (horizontal) axis. Target*  
22 *position is represented by a faint blue line. Gridlines show displacement of 1° of*  
23 *visual angle.*  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

### 51 **Single point localization**

55 In an examination of basic visual functions in 21 patients (mean MMSE 22/30)  
56 with PCA, Lehmann et al. (2011) identified single point localization deficits  
57  
58  
59  
60

1  
2  
3 (evaluated by means of pointing to a single dot displayed for a limited duration)  
4  
5 in 60% of individuals. With regard to other basic visual processes, point  
6  
7 localization performance only correlated with detection of motion coherence  
8  
9 (and not with measures of basic form or colour processing). Similarly with  
10  
11 regard to higher order visual skills, point localization performance correlated  
12  
13 significantly only with space perception such as VOSP dot counting and number  
14  
15 location (and not object perception measures such as fragmented image or  
16  
17 silhouette identification), suggesting this up-stream basic deficit in calculating  
18  
19 the 3D location of single points in space may partially underpin down-stream  
20  
21 difficulties in processing more complex spatial relationships.  
22  
23  
24  
25  
26  
27

28 In a further study of visuomotor function, PCA patients demonstrated many of  
29  
30 the deficits previously reported in individuals with optic ataxia, including  
31  
32 'magnetic misreaching' (a pathological reaching bias toward the point of visual  
33  
34 fixation when grasping peripheral targets; Meek et al., 2013). However, in  
35  
36 contrast to many reported cases of optic ataxia resulting from non-degenerative  
37  
38 disease, PCA patients showed additional deficits (e.g. in grip scaling during  
39  
40 memory guided grasping, face and object identification), consistent with the  
41  
42 widespread posterior cortical tissue loss common in PCA.  
43  
44  
45  
46  
47

48 Point localization deficits should also be borne in mind when evaluating visual  
49  
50 fields using standard perimetry techniques; though detection rates may vary  
51  
52 between quadrants (reflecting variations in the extent of atrophy/dysfunction  
53  
54 across the visual cortices), high rates of inconsistency of response to visual  
55  
56 stimulation particularly in the periphery suggest the influence of spatial and  
57  
58  
59  
60

1  
2  
3 attentional factors in contrast to classical dense hemi or quadrantinopias (e.g.  
4 Pelak et al., 2011). Point localization deficits are also not restricted to the visual  
5 modality; PCA and tAD patients show impairments in stationary sound position  
6 discrimination (Golden et al., 2015).  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

### 21 **Visual crowding**

22  
23  
24  
25 Crowding is a breakdown in the ability to identify objects in clutter, and  
26 particularly impairs object perception in peripheral, amblyopic and possibly  
27 developing vision (see Pelli et al., 2004). Excessive crowding has recently been  
28 shown to particularly limit object perception in the central vision of PCA patients  
29 (Yong et al., 2014). Using centrally-presented tests of letter identification under  
30 different flanking and spacing conditions (see Figure 3a), PCA patients were less  
31 accurate and slower to identify targets in the condensed than spaced condition  
32 even when the target letters were surrounded by flankers of a different category  
33 (see Figure 3b). The magnitude of the spacing effect was significantly associated  
34 with lower grey matter volume in the right collateral sulcus (see Figure 3c), a  
35 region approximating V4 and consistent with fMRI localization data related to  
36 peripheral crowding in healthy individuals (Anderson et al., 2012). The spacing  
37 effect was also observed for same, but not reverse, polarity flankers; this reverse  
38 polarity effect is characteristic of crowding and may reflect low-level segregation  
39 of visual information via on and off pathways (Kooi et al., 1994; but see  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Chakravarthi and Cavanagh, 2007). The error responses of PCA patients were  
4  
5 closely related to averaged target and flanker stimuli (but not individual target  
6  
7 or flanker stimuli); this is more consistent with averaging (Parkes et al., 2001;  
8  
9 Pelli et al., 2004; Greenwood et al., 2009) than substitution (Wolford, 1975)  
10  
11 accounts of crowding. These data indicate that PCA patients experience excessive  
12  
13 crowding, reflecting a pre-attentive process that uses averaging to regularize the  
14  
15 pathologically noisy representation of stimulus features presented in central  
16  
17 vision. Further evidence that excessive crowding is a critical factor underpinning  
18  
19 reading problems in PCA comes from studies (both cross-sectional and  
20  
21 longitudinal) of two patients who exhibited preserved reading (despite basic  
22  
23 visual, visuo-perceptual and visuo-spatial deficits) until the point at which they  
24  
25 also developed excessive crowding (Yong et al., 2013, submitted).  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 3. *Prominent effects and neural correlates of visual crowding in PCA* (Yong et al., 2014). (A) Target/flanker arrays used in the letter, shape and number flanker tasks under different spacing conditions. (B) Accuracy and naming latency data for the PCA and tAD group for letter, shape and number flankers in both

1  
2  
3 spatial conditions (RT=reaction time, Con=condensed, Spa=spaced). Error bars  
4 show standard error. (C) Statistical parametric map (SPM) of grey matter volume  
5 associated with a measure of crowding (spacing [shapes/numbers]). The SPM is  
6 displayed on axial, coronal and sagittal sections of the custom template in MNI  
7 space. When restricting analysis to a pre-specified region of interest (outlined in  
8 blue), there was an association between a greater degree of crowding and lower  
9 grey matter volume in the collateral sulcus (FWE corrected:  $p < .05$ ; peak  $t = 6.61$ ,  
10 location:  $x = 30$   $y = -58$   $z = -8$ ): the colour-map indicates  $t$ -values for this association.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

### 25 **Reduced effective field of vision**

26  
27  
28  
29  
30 A common yet striking complaint amongst PCA patients is a difficulty in reading  
31 newspaper headlines and other large font sizes whilst continuing to be able to  
32 read the small print. This apparent reverse size effect has been reported in a  
33 number of individuals with progressive visual impairment (e.g. Coslett et al.,  
34 1995; Saffran et al., 1990; Stark et al., 1997), and increased font size has been  
35 demonstrated to reduce reading accuracy or speed in 46% of PCA patients  
36 (compared to *increasing* reading speed in 18% of tAD patients and 7% of healthy  
37 controls; Yong et al., 2014). These effects have been attributed to a reduction in  
38 the effective visual field (Crutch et al., 2011; Russell et al., 2004). The influence of  
39 stimulus size on single object perception is also reflected in multi-object and  
40 complex scene perception, with many PCA patients exhibiting difficulty  
41 perceiving the visual world as a coherent whole (Shakespeare et al., 2015b). For  
42 example, one PCA patient's description of a picture of Brighton pier and beach  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 resembled the piecing together of a puzzle: 'It looks like a park or maybe a  
4 station or a building site, it looks a bit like the thing they're trying to erect for the  
5 Olympics...or it could be the beach because down here looks a bit sandy; yes it  
6 looks a bit like Brighton or somewhere like that' (see Crutch, 2014, Figure 3).  
7  
8 This impairment in multi-object and complex scene perception (meeting some  
9 definitions of simultanagnosia) has been argued to reflect a combination of  
10 attentional and spatial working memory deficits (Pisella et al., 2013).  
11  
12 Rehabilitative or compensatory strategies are still required, but one elegant  
13 study has shown that global processing in such individuals can be enabled by  
14 psychophysical biasing of visual pathways (Thomas et al., 2012).  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

### 28 **Translation and application**

29  
30  
31  
32 One natural progression from the improving characterization of basic visual  
33 function in PCA is to couple this knowledge with a detailed cognitive analysis of  
34 everyday activities in order to develop aids and strategies that support  
35 individuals with PCA in important everyday functional abilities. One example is  
36 an aid currently in development to facilitate reading in PCA (Yong et al., 2015a).  
37 Text reading is an early casualty for most people with PCA, with patients missing  
38 words, struggling to locate the beginning of the next line, and perceiving motion  
39 amongst static words and letters (see Figure 4a and 4b). The aid aims to create  
40 the optimal conditions for text reading in PCA by applying perceptual  
41 manipulations that minimise or evade many of the oculomotor, basic visual and  
42 visuospatial deficits outlined above (fixation instability, visual disorientation,  
43 excessive crowding, spatial agnosia and attentional deficits, reduced effective  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 field of vision). Words are serially presented in a single, central location in order  
4  
5 to reduce the challenge of localising words within sentences or identifying the  
6  
7 onset of subsequent lines of text (the aid moves the words to where the eyes are,  
8  
9 rather than the eyes having to move to where the words are; see Figure 4c).  
10  
11 Marking this location with a fixation box serves as a permanent cue to location  
12  
13 and hence aid fixation stability. Following reports of PCA patients being able to  
14  
15 better localize moving than static stimuli (Crutch, 2014; Midorikawa et al., 2008),  
16  
17 the aid successively moves words to fixation as a motion cue to assist  
18  
19 disorientated readers. The presentation of individual words was also designed to  
20  
21 prevent crowding effects from adjacent words. Furthermore, any risk of the  
22  
23 horizontal and vertical line features of the fixation box themselves crowding the  
24  
25 target text was attenuated by using opposite contrast polarity (Kooi et al., 1994;  
26  
27 Chakravarthi and Cavanagh, 2007). Early data suggest considerable gains in  
28  
29 reading accuracy for PCA patients (PCA mean reading accuracy: single-word  
30  
31 reading aid = 96%; individual patient improvement range: 6%-270%) relative to  
32  
33 standard paragraph reading (PCA 57%; tAD 96%), accompanied by increases in  
34  
35 self-rated measures of reading ease, pleasantness and understandability.  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 4. (A) Heatmap of PCA accuracy data from a sample passage (mean group accuracy rates: PCA=57%, tAD=98%, controls=100%). Lighter colours indicate the location of words most commonly missed or misread by PCA patients, and indicate spatial biases towards worse reading in later paragraphs and lines and in the centre of dense, crowded passages of text. (B) Word order from a sample passage for a PCA patient (MMSE score = 22/30). Arrows outline reading order; red arrows

1  
2  
3 *indicate omission of subsequent words through reading later sections of text, and*  
4 *yellow arrows indicate reading of earlier sections of text. (C) Single- and double-*  
5 *word presentations; words appear in the fixation box (1); following participants'*  
6 *responses, successive words move rapidly (2) into the fixation box (3). (D)*  
7  
8  
9  
10  
11 *Individual PCA patients' reading accuracy (percentage correct) for baseline*  
12 *(standard paragraph) and both reading aids (single word and double word),*  
13 *ordered in terms of severity of baseline impairment.*  
14  
15  
16  
17  
18  
19  
20

21 Beyond the dramatic visual deficits seen in PCA, perceptual impairments in  
22 people with typical AD and other dementias are greatly under-recognized, and  
23 little research has examined the benefits to patient and carer quality of life and  
24 mental and physical health that may be achievable by compensating for the  
25 effects of cortical visual impairment. This is despite evidence that even simple  
26 adaptations to the environment (e.g. using red rather than white tableware to  
27 increase visual contrast between food and plate) can have a significant impact on  
28 aspects of well-being (e.g. improved eating and drinking in advanced AD; Dunne  
29 et al., 2004; see also <http://dementia.stir.ac.uk/design/virtual-environments>).  
30  
31 Several current research projects are focused upon the quantitative evaluation of  
32 optometric and cortical visual function in order to improve visual health services  
33 for people with dementia (e.g. Hancock et al., 2015) or developing aids to  
34 facilitate locomotion and object localization within the home (e.g. Yong et al.,  
35 2015b).  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Future exploration

PCA is arguably the canonical 'visual dementia'. Nonetheless there remain a number of aspects of vision and visual experience in PCA that remain poorly understood. One outstanding issue relates to the status of visual fields in PCA. Some studies have claimed visual field defects in PCA (e.g. Pelak et al., 2011; Shea et al., 2014; Wan et al., 2015), but computerized visual field perimetry (Humphrey) has revealed poor false negative reliability indices (when a patient fails to see a significantly brighter stimulus at a location than was previously seen) making it difficult to determine to what extent performance reflects attentional deficits and/or true field defects (false negative rate [N=28 evaluations]: mean [SD] 22.3% [19.4%], range 0-66%; Pelak et al., 2011). Another issue is the impact of the cortical degeneration seen in PCA upon the retina and ophthalmic tracts. Preliminary serial diffusion tensor imaging evidence has indicated reductions in the integrity of the posterior thalamic radiations in PCA (Keihaninejad et al., 2012), and investigations of the possible impact of PCA upon the retina are underway (Lengyel et al., 2015). Further work is also required to investigate whether PCA may lead to functional deafferentation of and/or disease propagation along the optic tracts and more anterior parts of the visual system.

Despite these areas of limited understanding, evidence gathered to date and described above regarding basic visual functions (i.e. crowding, point localization, saccade generation, fixation stability) give rise to several future directions for pure and applied psychological research. For example, having

1  
2  
3 established PCA as a plausible neurodegenerative model of crowding of form  
4  
5 information, it remains to be seen whether other forms of visual information  
6  
7 (e.g. orientation, colour) exhibit crowding in PCA central vision comparable to  
8  
9 that of normal peripheral vision (Wilkinson et al., 1997; van den Berg et al.,  
10  
11 2007; Greenwood et al., 2012). Clinically, such information on basic visual  
12  
13 dysfunction in PCA will also influence both the design of appropriate visual  
14  
15 assessments (see Pelli et al., 2016) and a broader, more informed approach to  
16  
17 the evaluation and management of dementia-related visual impairment (College  
18  
19 of Optometrists, 2016).  
20  
21  
22  
23

24  
25 Although deteriorating vision is the presenting complaint for the majority of  
26  
27 individuals, it is critical that PCA is not pigeon-holed as solely a visual disorder,  
28  
29 and that multisensory and non-visual symptoms and experiences are also  
30  
31 investigated and addressed. Three brief examples are illustrative. First, the  
32  
33 interactions between vision and other sensory systems have been scarcely  
34  
35 examined in PCA despite the key role of the parietal lobes in integrating,  
36  
37 ordering and spatially representing information from multiple different senses.  
38  
39 Clinical descriptions such as patients asking “Am I the right way up?” (suggesting  
40  
41 an inability to relate body position to the gravitational vertical), or reporting an  
42  
43 even more extreme complete 180° room tilt illusion (reversal of vision  
44  
45 metamorphopsia) which could not be explained by impairment of the primary  
46  
47 vestibular senses (Crutch et al., 2011), suggest understanding of visuo-vestibular  
48  
49 interactions and therapeutic approaches to balance problems in PCA as key  
50  
51 areas for future study.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Second, approximately 30% of PCA patients exhibit asymmetrical left upper limb  
4 rigidity in addition to their visual dysfunction (Ryan, Shakespeare et al., 2014).  
5  
6 These motor deficits are associated with greater asymmetry of atrophy relative  
7 to 'pure' PCA patients, particularly involving right frontoparietal and peri-  
8 rolandic cortices, putamen, and thalamus. Despite evidence that such individuals  
9 are still likely to have underlying Alzheimer's disease, such heterogeneity within  
10 PCA offers insights into factors driving phenotypic variation in AD more  
11 generally and motivates clarification of the relationship between PCA and other  
12 related syndromes such as corticobasal syndrome (CBS).  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

24  
25 Finally, individuals with PCA may shed light on processes underpinning the  
26 representation of quantity-related verbal concepts. Acalculia is a well-recognised  
27 early symptom of PCA, and patients perform poorly on tests of cognitive  
28 estimation (Bisbing et al., 2015; Lehmann et al., 2011). However PCA patients  
29 also seem to exhibit greater difficulty than typical AD patients processing the  
30 precise meaning of non-numerical measurement unit terms (e.g. kilogram,  
31 centimetre; Suarez Gonzalez et al., personal communication). These preliminary  
32 findings suggest that parietal magnitude and number representations influence  
33 or shape anterior temporal lobe-dependent semantic representations of  
34 measurement unit terms. Such influence may be analogous to the idea that  
35 semantic representations of geographical place names (e.g. America, Cornwall)  
36 are spatially coded, that is fundamentally linked to ego- and allocentric  
37 representations of the actual geographical location and other spatial concepts  
38 (e.g. 'west'; Crutch and Warrington, 2003, 2010).  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Summary

Posterior cortical atrophy (PCA) is a focal degenerative condition that dramatically compromises the ability to see what and where things are whilst leaving episodic memory and insight relatively preserved in the early stages. PCA, particularly when attributable to Alzheimer's disease, has attracted much attention from clinical scientists for the questions it raises (e.g. how one disease can affect different individuals in such different ways) and the potential insights offered into fundamental disease mechanisms. Careful observation, systematic neuropsychological investigation and the opportunity to hear the experiences of those living with and caring for someone with PCA have already shed light on a range of classic and less well-studied neuropsychological deficits, and established PCA as a viable neurodegenerative disease model of certain visual phenomena. However, translating this emerging knowledge into compensatory and management strategies and aids which can effect meaningful change in the lives of those affected by dementia-related visual impairment remains a field in its relative infancy.

## Acknowledgements

This work was undertaken at UCLH/ UCL which received a proportion of funding from the Department of Health's NIHR Biomedical Research Centres funding scheme. The Dementia Research Centre is an Alzheimer's Research UK Co-

1  
2  
3 ordinating Centre (Grant no. ARUK-SRF2013-8). This work was supported by the  
4  
5 NIHR Queen Square Dementia Biomedical Research Unit. The Dementia Research  
6  
7 Centre is supported by Alzheimer's Research UK, Brain Research Trust (Grant  
8  
9 no. EQW 121 301), and The Wolfson Foundation. This work was supported by an  
10  
11 Alzheimer's Research UK Senior Research Fellowship and ESRC/NIHR  
12  
13 (ES/K006711/1) and EPSRC (EP/M006093/1) grants to SC; The Dunhill Medical  
14  
15 Trust [grant number R337/0214] to SC and KY. TS is supported by an  
16  
17 Alzheimer's Research UK Research Fellowship.  
18  
19

### 20 21 22 23 **Recommended readings**

24  
25  
26  
27  
28 Benson, D. F., Davis, R. J., & Snyder, B. D. (1988). Posterior cortical atrophy.  
29  
30 *Archives of Neurology*, 45, 789-793.

- 31  
32 - The original case series which described and established the clinical  
33  
34 concept of PCA.  
35  
36

37  
38  
39 Crutch, S. J., Lehmann, M., Schott, J. M., Rabinovici, G. D., Rossor, M. N.,  
40  
41 Fox, N. C. (2012). Posterior cortical atrophy. *Lancet Neurology*, 11, 170-8.

- 42  
43 - A thorough review of the clinical, neuroimaging, genetic and pathological  
44  
45 literature of PCA, which also emphasizes the need for consensus criteria  
46  
47 for clinical diagnosis and research.  
48  
49

50  
51  
52 Thomas, C., Kveraga, K., Huberle, E., Karnath, H. O., & Bar, M. (2012). Enabling  
53  
54 global processing in simultanagnosia by psychophysical biasing of visual  
55  
56 pathways. *Brain*, 135, 1578-85.  
57  
58  
59  
60

- An intriguing manipulation of a well-established cognitive effect.

Suárez-González, A., Henley, S. M., Walton, J., & Crutch, S. J. (2015). Posterior cortical atrophy: an atypical variant of Alzheimer disease. *Psychiatric Clinics of North America*, 38, 211-220.

- A recent review of the neuropsychiatric and psychological features of PCA and appropriate models of patient and carer support.

## References

Anderson, E. J., Dakin, S. C., Schwarzkopf, D. S., Rees, G., & Greenwood, J. A. (2012). The neural correlates of crowding-induced changes in appearance. *Current Biology*, 22, 1199-1206.

Benson, D. F., Davis, R. J., & Snyder, B. D. (1988). Posterior cortical atrophy. *Archives of Neurology*, 45, 789-793.

Bisbing, T. A., Olm, C. A., McMillan, C. T., Rascovsky, K., Baehr, L., Ternes, K., Irwin, D. J., Clark, R., & Grossman, M. (2015). Estimating frontal and parietal involvement in cognitive estimation: a study of focal neurodegenerative diseases. *Frontiers in Human Neuroscience*, 9, 317.

Chakravarthi, R., & Cavanagh, P. (2007). Temporal properties of the polarity advantage effect in crowding. *Journal of Vision*, 7, 11.

College of Optometrists (2016). *Visual Impairment and Dementia (VIDem) Summit*. In press.

Coslett, H. B., Stark, M., Rajaram, S., & Saffran, E. M. (1995). Narrowing the spotlight: a visual attentional disorder in presumed Alzheimer's disease. *Neurocase*, 1, 305-318.

Crutch, S. J. (2014). Seeing why they cannot see: understanding the syndrome and causes of posterior cortical atrophy. *Journal of Neuropsychology*, 8, 157-70

Crutch, S. J., Lehmann, M., Gorgoraptis, N., Kaski, D., Ryan, N., Husain, M., & et-al. (2011). Abnormal visual phenomena in posterior cortical atrophy. *Neurocase*, 17, 160-177.

Crutch, S. J., Lehmann, M., Schott, J. M., Rabinovici, G. D., Rossor, M. N., Fox, N. C. (2012). Posterior cortical atrophy. *Lancet Neurology*, 11, 170-8.

1  
2  
3 Crutch, S. J., & Warrington, E. K. (2003). Spatial coding of semantic information:  
4 knowledge of country and city names depends on their geographical proximity.  
5 *Brain*, *126*, 1821–1829.  
6

7 Crutch, S. J., & Warrington, E. K. (2010). Spatially coded semantic information  
8 about geographical terms. *Neuropsychologia*, *48*, 2120–2129.  
9

10 Dubois, B., Feldman, H. H., Jacova, C., Hampel, H., Molinuevo, J. L., Blennow, K., et  
11 al. Advancing research diagnostic criteria for Alzheimer's disease: the IWG-2  
12 criteria. *Lancet Neurology*, *13*, 614–29.  
13

14 Golden, H. L., Nicholas, J. M., Yong, K. X. X., Downey, L. E., Schott, J. M., Mummery,  
15 C. J., Crutch, S. J., & Warren, J. D. (2015). Auditory spatial processing in  
16 Alzheimer's disease. *Brain*, *138*, 189-202.  
17

18 Greenwood, J. A., Bex, P. J., & Dakin, S. C. (2009). Positional averaging explains  
19 crowding with letter-like stimuli. *Proceedings of the National Academy of Sciences*  
20 *USA*, *106*, 13130–5.  
21

22 Greenwood, J. A., Bex, P. J., & Dakin, S. C. (2012). Crowding follows the binding of  
23 relative position and orientation. *Journal of Vision*, *12*(3):18, 1–20.  
24

25 Hancock, B., Shah, R. et al. (2015). A proposal for a UK dementia eye care  
26 pathway. *Optometry in Practice*, *16*, 71-76.  
27

28 Hof, P. R., Bouras, C., Constantinidis, J., & Morrison, J. H. (1990). Selective  
29 disconnection of specific visual association pathways in cases of Alzheimer's  
30 disease presenting with Balint's syndrome. *Journal of Neuropathology and*  
31 *Experimental Neurology*, *49*, 168–84.  
32

33 Keihaninejad, S., Zhang, H., Shakespeare, T., Ryan, N., Malone, I., Frost, C.,  
34 Cardoso, M., Modat, M., Lehmann, M., Mancini, L., Thornton, J., Warren, J., Cash, D.,  
35 Crutch, S., Ourselin, S., & Fox, N. (2012). Longitudinal changes in fiber tract  
36 integrity in posterior cortical atrophy: Serial diffusion tensor imaging.  
37 Alzheimer's Association International Conference.  
38

39 Kooi, F. L., Toet, A., Tripathy, S. P., & Levi, D. M. (1994). The effect of similarity  
40 and duration on spatial interaction in peripheral vision. *Spatial Vision*, *8*, 255-  
41 279.  
42

43 Lehmann, M., Barnes, J., Ridgway, G. R., Wattam-Bell, J., Warrington, E. K., Fox, N.  
44 C., & Crutch, S. J. (2011). Basic visual function and cortical thickness patterns  
45 in posterior cortical atrophy. *Cerebral Cortex*, *21*, 2122–32.  
46

47 Lengyel, I., Quinn, N. B., Csincsik, L., Ritchie, C., Hogg, R. E., Crutch, S. J., Peto, T., &  
48 Shakespeare, T. J. (2015). Retinal image analysis and Alzheimer's disease.  
49 *Investigative Ophthalmology & Visual Science*, *56*(7), 2795.  
50  
51  
52

1  
2  
3 McKhann, G. M., Knopman, D. S., Chertkow, H., Hyman, B. T., Jack, C. R., Kawas, C.  
4 H., et al. (2011). The diagnosis of dementia due to Alzheimer's disease:  
5 recommendations from the National Institute on Aging-Alzheimer's Association  
6 workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's and*  
7 *Dementia*, 7, 263e269.

8  
9  
10 McMonagle, P., Deering, F., Berliner, Y., & Kertesz, A. (2006). The cognitive profile  
11 of posterior cortical atrophy. *Neurology*, 66, 331-338.

12  
13 Meek, B. P., Shelton, P., & Marotta, J. J. (2013). Posterior cortical atrophy:  
14 visuomotor deficits in reaching and grasping. *Frontiers in Human Neuroscience*, 7,  
15 294.

16  
17  
18 Mendez, M. F., Ghajarania, M., & Perryman, K. M. (2002). Posterior cortical  
19 atrophy: clinical characteristics and differences compared to Alzheimer's  
20 disease. *Dementia and Geriatric Cognitive Disorders*, 14, 33e40.

21  
22  
23 Midorikawa, A., Nakamura, K., Nagao, T., & Kawamura, M. (2008). Residual  
24 perception of moving objects: dissociation of moving and static objects in a case  
25 of posterior cortical atrophy. *European Neurology*, 59, 152-8.

26  
27  
28 Ossenkoppele, R., Schonhaut, D. R., Baker, S. L., O'Neil, J. P., Janabi, M., Ghosh, P.  
29 M., Santos, M., Miller, Z. A., Bettcher, B. M., Gorno-Tempini, M. L., Miller, B. L.,  
30 Jagust, W. J., Rabinovici, G. D. (2015). Tau, amyloid, and hypometabolism in a  
31 patient with posterior cortical atrophy. *Annals of Neurology*, 77, 338-42.

32  
33  
34 Parkes, L., Lund, J., Angelucci, A., Solomon, J. A., & Morgan, M. (2001). Compulsory  
35 averaging of crowded orientation signals in human vision. *Nature Neuroscience*,  
36 4, 739-44.

37  
38  
39 Pelak, V. S., Smyth, S. F., Boyer, P. J., & Filley, C. M. (2011). Computerized visual  
40 field defects in posterior cortical atrophy. *Neurology*, 77, 2119-2122.

41  
42  
43 Pelli, D. G., Palomares, M., & Majaj, N. J. (2004). Crowding is unlike ordinary  
44 masking: Distinguishing feature integration from detection. *Journal of Vision*, 4,  
45 1136-1169.

46  
47  
48 Pelli, D. G., Waugh, S. J., Martelli, M., Crutch, S. J., Primativo, S., Yong, K. X., Rhodes,  
49 M., Yee, K., Wu, X., Famira, H. F., Yiltiz, H. (2016) A clinical test for visual  
50 crowding. *F1000Research*. In press.

51  
52  
53 Pisella, L., Sergio, L., Blangero, A., Torchin, H., Vighetto, A., & Rossetti, Y. (2009).  
54 Optic ataxia and the function of the dorsal stream: contributions to perception  
55 and action. *Neuropsychologia*, 47, 3033-3044.

56  
57  
58 Renner, J. A., Burns, J. M., Hou, C. E., McKeel, D. W., Storandt, M., & Morris, J. C.  
59 (2004). Progressive posterior cortical dysfunction: a clinicopathologic series.  
60 *Neurology*, 63, 1175-80.

1  
2  
3 Russell, C., Malhotra, P., & Husain, M. (2004). Attention modulates the visual field  
4 in healthy observers and parietal patients. *Neuroreport*, *15*, 2189e2193.

5  
6 Ryan, N. S., Shakespeare, T. J., Lehmann, M., Keihaninejad, S., Nicholas, J. M.,  
7 Leung, K. K., Fox, N. C., & Crutch, S. J. (2014). Motor features in posterior cortical  
8 atrophy and their imaging correlates. *Neurobiology of Aging*, *35*, 2845-57.

9  
10 Saffran, E. M., Fitzpatrick-DeSalme, E. J., & Coslett, H. B. (1990). Visual  
11 disturbances in dementia. *Modular deficits in Alzheimer-type dementia*, *44*, 297-  
12 327.

13  
14 Shakespeare, T. J., Kaski, D., Yong, K. X. X., Paterson, R. W., Slattery, C. F., Ryan, N.  
15 S., Schott, J. M., & Crutch, S. J. (2015a). Abnormalities of fixation, saccade and  
16 pursuit in posterior cortical atrophy. *Brain*. Apr 19. pii: awv103.

17  
18 Shakespeare, T. J., Pertzov, Y., Yong, K. X. X., Nicholas, J., & Crutch SJ. (2015b).  
19 Reduced modulation of scanpaths in response to task demands in posterior  
20 cortical atrophy. *Neuropsychologia*, *68*, 190-200.

21  
22 Shea, Y. F., & Chu, L. W. (2014). Visual field defect as an uncommon presentation  
23 of posterior cortical atrophy. *Geriatrics & Gerontology International*, *14*, 1003-4.

24  
25 Stark, M. E., Grafman, J., & Fertig, E. (1997). A restricted 'spotlight' of attention in  
26 visual object recognition. *Neuropsychologia*, *35*, 1233-1249.

27  
28 Tang-Wai, D. F., Graff-Radford, N. R., Boeve, B. F., Dickson, D. W., Parisi, J. E.,  
29 Crook, R., & et-al. (2004). Clinical, genetic, and neuropathologic characteristics of  
30 posterior cortical atrophy. *Neurology*, *63*, 1168-1174.

31  
32 Thomas, C., Kveraga, K., Huberle, E., Karnath, H. O., & Bar, M. (2012). Enabling  
33 global processing in simultanagnosia by psychophysical biasing of visual  
34 pathways. *Brain*, *135*, 1578-85.

35  
36 van den Berg, R., Roerdink, J. B. T. M., & Cornelissen, F. W. (2007). On the  
37 generality of crowding: Visual crowding in size, saturation, and hue compared to  
38 orientation. *Journal of Vision*, *7*(2):14, 1-11, [http://](http://www.journalofvision.org/content/7/2/14)  
39 [www.journalofvision.org/content/7/2/14](http://www.journalofvision.org/content/7/2/14), doi:10.1167/7.2.14.

40  
41 Wan, S. L., Bukowska, D. M., Ford, S., & Chen, F. K. (2015). Posterior Cortical  
42 Atrophy Presenting with Superior Arcuate Field Defect. *Case Reports in*  
43 *Ophthalmological Medicine*, 2015:796381.

44  
45 Wilkinson, F., Wilson, H. R., & Elleberg, D. (1997). Lateral interactions in  
46 peripherally viewed texture arrays. *Journal of the Optical Society of America. A,*  
47 *Optics, Image Science, and Vision*, *14*, 2057-2068.

48  
49 Worf, G. (1975). Perturbation model for letter identification. *Psychological*  
50 *Review*, *82*, 184-99.

1  
2  
3 Yong, K. X. X., Warren, J. D., Warrington, E. K., & Crutch, S. J. (2013). Intact reading  
4 in patients with profound early visual dysfunction. *Cortex*, *49*, 2294-306.

6 Yong, K. X. X., Shakespeare, T. J., Cash, D., Henley, S. M., Warren, J. D., & Crutch, S. J.  
7 (2014). (Con)text-specific effects of visual dysfunction on reading in posterior  
8 cortical atrophy. *Cortex*, *57*, 92-106.

10 Yong, K. X. X., Shakespeare, T. J., Cash, D., Henley, S. M., Nicholas, J. M., Ridgway, G.  
11 R., Golden, H. L., Warrington, E. K., Carton, A. M., Kaski, D., Schott, J. M., Warren, J.  
12 D., & Crutch, S. J. (2014). Prominent effects and neural correlates of visual  
13 crowding in a neurodegenerative disease population. *Brain*, *137*, 3284-99.

16 Yong, K. X. X., Rajdev, K., Shakespeare, T. J., Leff, A. P., & Crutch, S. J. (2015a).  
17 Facilitating text reading in posterior cortical atrophy. *Neurology*, *85*, 339-48.

20 Yong, K. X. X., Holloway, C., Carton, A., Yang, B., Suzuki, T., Serougne, R.,  
21 Boampong, D., Voyce, D., Langham, L., Frost, C., Sullivan, M. P., McCarthy, I., Tyler,  
22 N., & Crutch, S. J. (2015b). Effects of Cortical Visual Impairment on Navigational  
23 Ability in Posterior Cortical Atrophy and Typical Alzheimer's Disease.  
24 Alzheimer's Association International Conference.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60