

Adjuvant FOLFOX +/- cetuximab in full *RAS* and *BRAF* wildtype stage III colon cancer patients

J. Taieb¹, R. Balogoun², K. Le Malicot³, J. Tabernero⁴, E. Mini⁵, G. Folprecht⁶, J-L. Van Laethem⁷, J-F. Emile⁸, C. Mulot⁹, S. Fratté¹⁰, C-B. Levaché¹¹, L. Saban-Roche¹², J. Thaler¹³, L N. Petersen¹⁴, J. Bridgewater¹⁵, G. Perkins^{1,2}, C. Lepage¹⁶, E. Van Cutsem¹⁷, A. Zaanan^{1,2} and P. Laurent-Puig² for PETACC8 investigators*

1- Department of Gastroenterology and GI oncology, Sorbonne Paris Cité, Université Paris Descartes, Hôpital Européen Georges Pompidou, Paris, France

2- Department of Biology, Université Paris Descartes, Sorbonne Paris Cité, France; Assistance Publique Hôpitaux de Paris, Hôpital Européen Georges Pompidou, Paris, France; INSERM UMR-S1147,

3 - Fédération francophone de cancérologie digestive (FFCD), Dijon, France

4- Medical Oncology Department, Vall d'Hebron University Hospital and Institute of Oncology (VHIO), Barcelona, Universitat Autònoma de Barcelona, Spanish Gastrointestinal Tumours TTD Group, Spain.

5- Department of Experimental and Clinical Medicine, Section of Internal Medicine, University of Florence, Florence, Italy

6- Medical Department I, University Hospital Carl Gustav Carus, Dresden, Germany

7- Department of Gastroenterology, Hôpital Universitaire Erasme, Brussels, Belgium

8- Pathology Department, Ambroise Paré Hospital, Boulogne, France

9- Université Paris Descartes, Sorbonne Paris Cité, CRB EPIGENETEC, INSERM UMR-S1147

10- Department of Gastroenterology, Centre Hospitalier de Belfort-Montbéliard, France

11 - Department of Radiotherapy and medical oncology , Polyclinique Francheville, Périgueux, France

12 - Department of medical Oncology, Institut de Cancerologie de la Loire, Saint-Priest-En-Jarez, France

13- Department of Internal Medicine IV, Klinikum Wels-Grieskirchen, Wels, Austria;

14 - Department of Oncology, Rigshospitalet, København, Denmark;

15- UCL Cancer Institute, University College London, London, United Kingdom;

16 - Hepato-Gastroenterology Department, Dijon University Hospital and INSERM U 866, Dijon, France;

17- Digestive Oncology, University Hospitals Leuven and KU Leuven, Leuven, Belgium

*Please see the Supplementary Appendix for a list of the Investigators/Collaborators.

Word count: 2411

Correspondence:

Pr Julien Taieb, Georges Pompidou European Hospital, Department of hepatogastroenterology and GI oncology, Sorbonne Paris Cité / Université Paris Descartes, 20 rue Leblanc 75015 Paris, France Phone: 33 1 56093551; Fax: 33 1 56093441 julien.taieb@egp.aphp.fr / jtaieb75@gmail.com

Background: RAS mutations have been shown to confer resistance to anti-EGFR treatment. We analyzed the results of the PETACC8 trial (cetuximab + FOLFOX vs FOLFOX) in full RAS and BRAF wildtype (WT) patients (pts) with resected stage III colon cancer.

Methods: Exons 2, 3 and 4 of KRAS and NRAS, and BRAF exons 11 and 15, were sequenced using the Ampliseq colon-lung cancer panel version 2, in PETACC8 trial pts who consented to translational research. The impact of cetuximab on time to recurrence (TTR), disease-free survival (DFS) and overall survival (OS) was investigated in pts with tumors harboring RAS & BRAF WT and RAS mutations. The prognostic value of each individual mutation was also tested.

Results: Among the 2559 pts analyzed, 745 pts (29%) were known to have KRAS exon 2 mutations and 163 pts (6.4%) the BRAF V600E mutation. Of the remaining 1651 pts, 1054 were assessed by NGS, showing that a further 227 pts (21%) had KRAS exon 2,3,4 or NRAS exon 2,3,4 mutations, and that 46 pts (4.4%) had a newly diagnosed BRAF mutation. Cetuximab added to FOLFOX did not significantly improve TTR, DFS or OS in pts with RAS WT or RAS & BRAF WT tumors (HR 0.77 to 1.03, all $P > 0.05$). Cetuximab addition was not either significantly deleterious in RAS mutant pts or in pts with rare RAS or BRAF mutations. In the overall trial population, NRAS and KRAS codon 61 mutations were the only rare mutations with the same pejorative prognostic value as KRAS exon 2 or BRAF V600E mutations.

Conclusion

Though not significant, the clinically relevant 0.76 adjusted HR observed for DFS in favor of adding cetuximab to FOLFOX, in full RAS and BRAF WT stage III colon

cancer pts, may justify a new randomized controlled trial testing EGFR inhibitors in this setting.

Clinical trial number: This is an ancillary study of the PETACC8 trial: EUDRACT 2005-003463-23.

Keywords: stage III colon cancer; *RAS* mutations; *BRAF* mutations; cetuximab; FOLFOX; phase III; adjuvant; prognosis

Key message: In 1900 stage III colon cancer patients fully characterized by NGS for *RAS* and *BRAF*, adding cetuximab to the standard FOLFOX results in a trend to a better outcome in *RAS* & *BRAF* wildtype patients. Though not significant, the clinically relevant 0.76 adjusted/HR observed for DFS in favor of adding cetuximab may justify a new randomized trial with anti-EGFRs in this setting.

Introduction

The Pan-European Trials in Alimentary traCt Cancer 8 (PETACC-8) study tested FOLFOX4, with or without cetuximab, after curative resection of stage III colon cancer [1]. Promising phase II and III studies of cetuximab adjunction to FOLFOX4 in metastatic colorectal cancer showed impressive response and disease-control rates, suggesting possible synergy of this new combination [2, 3]. The PETACC-8 protocol was amended on 17 June 2008, restricting enrolment to patients with *KRAS* exon 2 WT tumours and increasing the sample size. The first analysis of the trial results was negative, with no improvement in disease-free survival (DFS) or overall survival (OS) when cetuximab was added to FOLFOX [1].

KRAS exon 2 mutations are predictive of resistance to anti-epidermal growth factor receptor (EGFR) therapy in patients with metastatic colorectal cancer [3-6], as are activating mutations in *KRAS* exon 3 or 4 and in *NRAS* exon 2, 3 or 4 [7, 8].

BRAF mutations are typically exclusive of *RAS* mutations, and clinical data suggest that the *BRAF* V600E mutation is predictive of poorer survival but not of anti-EGFR efficacy in patients with metastatic colorectal cancer [9, 10], however, the low prevalence of these mutations makes it difficult to evaluate their possible biomarker status.

Patient selection based on tumour mutational status might thus improve the harm-benefit profile of anti-EGFR therapy. This has been largely demonstrated in metastatic colorectal cancer [7, 8] but not yet in the adjuvant setting. We and others recently found that *BRAF* V600E and *KRAS* exon 2 mutations were prognostic in stage III colon cancer, being associated with shorter time to recurrence (TTR), OS, and survival after relapse [11-14]. However, anti-EGFR efficacy has not yet been evaluated in selected patients with *RAS* WT and *BRAF* WT resected stage III colon

cancer.

We used the Ampliseq colon-lung cancer panel version 2 to sequence exons 2, 3 and 4 of *KRAS* and *NRAS*, as well as *BRAF* exons 11 and 15, amongst those PETACC8 trial participants who consented to translational research. TTR, DFS and OS were analyzed in full *RAS* WT patients and full *RAS* and *BRAF* WT patients. The prognostic impact of individual rare *RAS* and *BRAF* mutations was also investigated.

Materials and methods

Patients

PETACC8 trial participants underwent complete resection of histologically proven stage III colon adenocarcinoma, and were then randomly assigned to receive 6 months of either FOLFOX or FOLFOX+cetuximab, with regular monitoring, as described elsewhere [1]. The trial started in December 2005. The protocol was amended in June 2008 to enroll only patients with *KRAS* exon 2 WT tumours, and the sample size was increased to maintain power of statistical analyses. The study ended on 9 November 2009. Specific written informed consent was required from each patient included in the planned translational program of the trial.

DNA Extraction and Mutation Analysis

Tumour samples were prospectively banked. Tumour DNAs were extracted from FFPE tissues containing more than 50% of tumour cells by using the QIAamp® DNA Mini Kit (Qiagen®). Molecular analysis, centralised at Georges Pompidou European Hospital, was performed retrospectively for the 2096 patients included before the trial amendment and prospectively for the other 463 patients. *KRAS* hotspot mutations

(c.34G>A/p.G12S, c.34G>C/p.G12R, c.34G>T/p.G12C, c.35G>A/p.G12D, c.35G>C/p.G12A, c.35G>T/p.G12V and c.38G>A p.G13D) and the *BRAF* V600E mutation (c.1799T>A/p.V600E) were detected by real-time PCR with TaqMan® probes (Applied Biosystems). The assays are alteration-specific and robustly detect 10% of mutated alleles for all the mutations tested.

Exons 2, 3 and 4 of *KRAS* and *NRAS*, as well as *BRAF* exons 11 and 15, were sequenced with the Ampliseq colon-lung cancer panel version 2 in the PETACC8 trial participants who consented to translational research.

Statistical Analyses

TTR, DFS and OS were analysed in patients with any *RAS* or *BRAF* mutations, *RAS* & *BRAF* WT status, and rare *RAS* mutations. The individual prognostic value of each mutation was also analysed.

TTR was defined as the time between randomization and local or metastatic recurrence or death related to disease recurrence, whichever occurred first. DFS was defined as the time between randomization and local or metastatic recurrence or diagnosis of a second colorectal cancer, or death from any cause, whichever occurred first. OS was defined as the time between randomization and death from any cause.

For baseline comparisons, categorical factors were compared with χ^2 tests and continuous factors with standard parametric or non-parametric tests, depending on their normality. Continuous variables are reported as mean (SD) and median (interquartile range, IQR) values.

TTR, DFS and OS curves were estimated using the Kaplan–Meier method. Differences between groups of patients were analysed with log-rank tests. Cox models, Kaplan-Meier curves and forest plots were used for all analyses. Factors included in multivariate analyses were the treatment group and baseline prognostic factors that were clinically relevant or significant in univariate analysis, namely tumour grade, pT stage, pN stage, venous embolism, lymphatic invasion (VELI), bowel obstruction/perforation, and tumour location.

A two-sided significance level of 5% was applied for all analyses. Results were not adjusted for multiple comparisons. All statistical analyses were done by FFCD statisticians using SAS statistical software (version 9.4). The database was locked in July 2015.

Results

Study population

Among the 2559 patients included in the PETACC8 phase III study, 741 were *KRAS* exon 2 mutated and 167 were *BRAF* V600E mutated. Of the remaining 1651 patients, 1054 gave their written consent for translational research and had sufficient tumour material for NGS analyses. NGS failed in 62 cases. The remaining 992 patients were fully analysed. A total of 1900 patients (including RAS mutated patients) met all the criteria for full molecular analysis (informed consent, sufficient material, and technical success) (Supplementary Figure 1). The patients' baseline and tumour characteristics are summarized in Table 1. The demographic and clinical characteristics of the patients included in the molecular study ($N=1900$) were not

significantly different from those of the entire randomized population ($N=2559$) (Supplementary Table 1).

RAS and BRAF mutational status

Amongst the 1900 patients included in the molecular study, 719 (38%) were double wildtype, 968 (51%) were *RAS* mutated and 213 (11%) were *BRAF* mutated (Figure 1). *KRAS*, *NRAS* and *BRAF* mutation frequencies are summarized in Figure 1.

The most frequently mutated *KRAS* exon was exon 2 (80.9%), followed by exons 4 (8.3%) and 3 (4.6%); two tumours (0.2%) were mutated on two different exons (Table 2). As expected, codon 12 was the most frequently mutated codon (75.9%), followed by codon 13 (18.4%). *NRAS* exons 2, 3 (codon 61) and 4 were mutated in respectively 30, 31 and 2 cases.

BRAF was mutated in 213 tumours, including 192 tumours (90%) harbouring the V600E mutation. The second most frequent mutation affected codon 469, in 8 cases (3.8%). The mutations were grouped for analysis into V600E and non V600E.

KRAS and *BRAF* mutations were both present in 8 tumours (4 V600E and 4 non V600E). *KRAS* and *NRAS* mutations were both present in two tumours (*KRAS* pA146T associated with *NRAS* p.G12D and with *NRAS* p.A146V in one case each). *NRAS* and *BRAF* mutations were both present in 3 tumours, all with non V600E *BRAF* mutations.

Clinical outcomes according to RAS and BRAF mutational status

As previously reported, adding cetuximab to FOLFOX did not improve TTR in the whole trial population [1] (Figure 2-A). In the *RAS* WT and *BRAF* WT population, a trend towards better outcomes was seen in the cetuximab group but the difference did not reach statistical significance for TTR (HR:0.77 (0.55-1.08); $P=0.12$) (Figure 2-B), DFS (HR:0.85 (0.63-1.14); $P=0.27$) or OS (HR:1.03 (0.70-1.50); $P=0.89$). In multivariate analyses, the results were better but still not significant: TTR (HR=0.70 (95%CI: 0.48 - 1.03); $P=0.07$), DFS (HR=0.76 (95%CI: 0.54 - 1.06); $P=0.11$), and OS (HR=0.90 (95%CI: 0.59 - 1.36); $P=0.60$).

In patients with *RAS*-mutated tumours, the addition of cetuximab to FOLFOX was associated with a trend towards poorer TTR (HR:1.14 (0.91-1.44); $P=0.25$) (Figure 2-C), DFS (HR:1.13 (0.91-1.40); $P=0.27$) and OS (HR:1.29 (0.99-1.69); $P=0.061$). These trends were less pronounced in multivariate analyses: TTR (HR=1.09 (95%CI: 0.84 - 1.41); $P=0.51$), DFS (HR=1.06 (95%CI: 0.83 - 1.35); $P=0.64$), OS (HR= 1.17 (95%CI: 0.87 - 1.57); $P=0.30$).

As DFS is the usual endpoint for adjuvant trials, DFS Kaplan Meier curves are shown in supplementary figure 2-A, 2-B and 2-C.

Rare *RAS* and *BRAF* mutations (i.e. *KRAS* exon 3,4 ; *NRAS* exon 2,3,4 and *BRAF* non V600E) tended to be associated with a deleterious effect of cetuximab, with HRs of 1.6 for TTR ($P=0.09$) and 1.61 for OS ($P=0.13$).

Prognostic value of RAS and BRAF mutations

In the overall study population, *KRAS* exon 2 and *BRAF* V600E mutations were associated with worse outcomes when compared to *RAS* and *BRAF* WT status, as

previously described [12]. This was also the case of *KRAS* and *NRAS* (exon 3) codon 61 rare mutants with respect to TTR and OS, contrary to other rare *RAS* or *BRAF* mutants (Figure 3).

The number of rare mutations was too small for meaningful multivariable analysis.

Discussion

KRAS and *NRAS* are closely related to *RAS* oncogene family members, and mutations at codon 12, 13, 61, 117 or 146 of either gene result in increased levels of guanosine triphosphate-bound *RAS* proteins [15, 16]. *KRAS* and *NRAS* mutations at these codons tend to be mutually exclusive in colorectal tumours, suggesting functional redundancy [17]. Mutations in *HRAS*, the third member of the *RAS* family, are infrequent in colorectal cancer [17, 18]. Clinical data suggest that *RAS* genes mutations are also associated with worse outcomes in the adjuvant setting [11-14].

Previous trials of anti-EGFR therapies combined with irinotecan or oxaliplatin-containing regimens showed no benefit in patients with *KRAS* exon 2 mutations [2, 6].

Randomized phase 3 trials of panitumumab, given alone [19] or in combination with FOLFOX or FOLFIRI [3, 5, 7], showed no response to this anti-EGFR therapy in patients with metastatic colorectal tumours harboring a mutation in *KRAS* or *NRAS*.

This was also the case in recent analyses of randomized trials with cetuximab [8, 20].

All these studies involved patients with metastatic colorectal cancer. By contrast, we assessed here the effect of adjuvant cetuximab plus FOLFOX in patients with fully resected primary stage III colon tumours and full *KRAS*, *NRAS* and *BRAF* characterization.

Removing patients with rare RAS and BRAF mutations, with a poor outcome, from the target efficacy population reveals a trend to a positive effect of the addition of cetuximab to standard FOLFOX in patients with RAS & BRAF wildtype tumours. Although the impact of cetuximab was not statistically significant, it might be clinically relevant. In the MOSAIC pivotal trial, adding oxaliplatin to 5-FU improved DFS, with an HR of 0.8. Here, multivariate analysis adjusted for pT, pN, histological grade, VELL and tumour location yielded an HR of 0.76 (95%CI:0.54-1.06). This suggests that a new randomized trial powered to demonstrate such a difference in wildtype colon cancer patients may be relevant, especially after a 12-year period with no advances in adjuvant treatment of stage III colon cancer. If such trial is not forthcoming, our results would have to be confirmed using at least internally (other sequencing approaches for example) and externally (on other datasets) before discussing any practice change. New markers of colon cancer sensitivity to anti-EGFRs are emerging and could in future also be assessed in samples from PETACC8 and other adjuvant trials of anti-EGFRs, such as the NCCTG N0147 study, in order to generate hypotheses for future trials [21, 22].

Although adding cetuximab to FOLFOX tended to be beneficial in terms of TTR and DFS, this was not the case for OS (HR of 0.9 in adjusted analyses). This discordance between OS and TTR/DFS suggests that survival after relapse may differ between patients who do and do not receive adjuvant cetuximab, possibly because of lower cetuximab prescription rates in the metastatic setting when patients have received adjuvant cetuximab. Further analyses of survival after recurrence, and of treatments received at recurrence, are needed to clarify this point.

A deleterious effect of cetuximab and panitumumab has been reported in some patients with *RAS*-mutated tumours treated with FOLFOX in the metastatic setting [7,

20]. This was not the case of patients with *RAS*-mutated metastatic colorectal cancer receiving irinotecan-based backbone chemotherapy [8]. In our study of stage III colon cancer, there was only a non-significant trend towards worse outcomes with cetuximab in *RAS*-mutant patients.

This trend towards a deleterious effect of cetuximab was even stronger in patients with rare *RAS* mutations, but again it did not reach statistical significance, possibly owing to the small number of patients with rare *RAS* mutations (n=185).

We and others have shown that *KRAS* exon 2 and *BRAF* V600E mutations are associated with a poor prognosis in stage III colon cancer and especially in the 90% of patients with MSS tumours [14]. However, the prognostic value of rare *KRAS*, *NRAS* and *BRAF* mutations has rarely been studied in this setting. Gavin et al reported in 2299 stage II and III colon tumours a similar frequency of *NRAS* mutations (2.9%) that were associated with a worse TTR (HR=1.53; 95%CI, 1.01-2.31; $P=0.04$), but this difference disappeared in multivariate analysis and was not significant for OS [23]. A recent retrospective study of rare *KRAS* mutations at codons 12, 13 and 61 in stage II-III colon cancer patients showed no significant impact on DFS or OS [24]. However, the impact of individual *KRAS* mutations was not studied, the sample was quite small, and the study was retrospective. Modest et al. very recently studied the prognostic impact of *RAS* mutations in metastatic patients and found that only G13D and G12C had prognostic value and not rare mutations [25]. We found no recent data on the prognostic value of rare *BRAF* mutations in the adjuvant setting. In the metastatic setting, *BRAF* non V600E-mutated tumours seem to carry a better prognosis [26]. In the present work, we found that only *KRAS* and *NRAS* codon 61 mutations had significant negative prognostic

value, while other rare *RAS* or *BRAF* non V600E mutations did not seem to affect patient outcome. However, these results need to be confirmed in larger series with full *RAS* and *BRAF* mutational analyses.

In conclusion, adding cetuximab to standard FOLFOX adjuvant therapy in stage III colon cancer results in a non significant trend towards better outcomes in *RAS* & *BRAF* WT patients. No significant detrimental effect was observed in *RAS* mutant patients. Though not significant, the clinically relevant 0.76 adjusted HR observed for DFS in favor of adding cetuximab to FOLFOX in full *RAS* and *BRAF* WT stage III colon cancer pts, may justify a new randomized controlled trial testing EGFR inhibitors in this setting.

Acknowledgements :

The study was sponsored by “Fédération Francophone de Cancérologie Digestive” (FFCD), which was responsible for the study management for design and conduct of the study; for collection, management, analysis, and interpretation of the data; for preparation, review, or approval of the manuscript; and for decision to submit the manuscript for publication. We thank all participating patients and their families, and the study groups and investigators from the participating countries (appendix); we also thank the team from the FFCD data centre. The FFCD also thanks the Ligue Nationale Contre le Cancer for its grant.

Funding:

Merck KGaA and Sanofi-Aventis supported the study, Merck provided the study cetuximab and financial support for study management, Sanofi-Aventis provided financial support for the provision of oxaliplatin to Belgian sites when necessary. Merck KGaA reviewed the manuscript for medical accuracy only before journal submission. The authors are fully responsible for the content of this manuscript, and the views and opinions described in the publication reflect solely those of the authors.

Conflict of Interest Disclosures:

Dr Taieb has participated in consulting or/and advisory boards for Merck, Sanofi, Roche Genentech, Pfizer and Amgen; Dr Zaanani for Roche, Merck Serono, Amgen, Sanofi and Lilly; Dr Tabernero from Amgen, ImClone Systems, Lilly, Millennium, Novartis, Roche/Genentech, Sanofi, Celgene, Chugai Pharma, Taiho Pharmaceutical, Boehringer Ingelheim and Merck Serono; Dr Folprecht from Roche, Merck KGaA, Lilly and Bristol; Dr Laurent-Puig from Sanofi, Merck Serono, Amgen, Roche, Genomic Health, Myriad Genetics and Pfizer.

Dr Folprecht from Merck KGaA, Lilly and Bayer; Dr Laurent-Puig from Sanofi, Merck Serono, Amgen, Roche, Genomic Health, Myriad Genetics and Pfizer.

Dr Bridgewater received honoraria from Merck.

Dr Folprecht has received research funding from Merck KGaA. Dr Van Cutsem received research funding from Amgen, Bayer, Boehringer, Celgene, Ipsen, Merck KGA, Merck, Novartis, Roche and Sanofi.

No other disclosures are reported.

References

1. Taieb J, Tabernero J, Mini E et al. Oxaliplatin, fluorouracil, and leucovorin with or without cetuximab in patients with resected stage III colon cancer (PETACC-8): an open-label, randomised phase 3 trial. *Lancet Oncol* 2014; 15: 862-873.
2. Bokemeyer C, Bondarenko I, Makhson A et al. Fluorouracil, leucovorin, and oxaliplatin with and without cetuximab in the first-line treatment of metastatic colorectal cancer. *J Clin Oncol* 2009; 27: 663-671.
3. Douillard JY, Siena S, Cassidy J et al. Randomized, phase III trial of panitumumab with infusional fluorouracil, leucovorin, and oxaliplatin (FOLFOX4) versus FOLFOX4 alone as first-line treatment in patients with previously untreated metastatic colorectal cancer: the PRIME study. *J Clin Oncol* 2010; 28: 4697-4705.
4. Amado RG, Wolf M, Peeters M et al. Wild-type KRAS is required for panitumumab efficacy in patients with metastatic colorectal cancer. *J Clin Oncol* 2008; 26: 1626-1634.
5. Peeters M, Price TJ, Cervantes A et al. Randomized phase III study of panitumumab with fluorouracil, leucovorin, and irinotecan (FOLFIRI) compared with FOLFIRI alone as second-line treatment in patients with metastatic colorectal cancer. *J Clin Oncol* 2010; 28: 4706-4713.
6. Van Cutsem E, Kohne CH, Hitre E et al. Cetuximab and chemotherapy as initial treatment for metastatic colorectal cancer. *N Engl J Med* 2009; 360: 1408-1417.
7. Douillard JY, Oliner KS, Siena S et al. Panitumumab-FOLFOX4 treatment and RAS mutations in colorectal cancer. *N Engl J Med* 2013; 369: 1023-1034.
8. Van Cutsem E, Lenz HJ, Kohne CH et al. Fluorouracil, leucovorin, and irinotecan plus cetuximab treatment and RAS mutations in colorectal cancer. *J Clin Oncol* 2015; 33: 692-700.
9. Rowland A, Dias MM, Wiese MD et al. Meta-analysis of BRAF mutation as a predictive biomarker of benefit from anti-EGFR monoclonal antibody therapy for RAS wild-type metastatic colorectal cancer. *Br J Cancer* 2015; 112: 1888-1894.
10. Pietrantonio F, Petrelli F, Coiu A et al. Predictive role of BRAF mutations in patients with advanced colorectal cancer receiving cetuximab and panitumumab: a meta-analysis. *Eur J Cancer* 2015; 51: 587-594.
11. Blons H, Emile JF, Le Malicot K et al. Prognostic value of KRAS mutations in stage III colon cancer: post hoc analysis of the PETACC8 phase III trial dataset. *Ann Oncol* 2014; 25: 2378-2385.
12. Taieb J, Zaanan A, Le Malicot K et al. Prognostic Effect of BRAF and KRAS Mutations in Patients With Stage III Colon Cancer Treated With Leucovorin, Fluorouracil, and Oxaliplatin With or Without Cetuximab: A Post Hoc Analysis of the PETACC-8 Trial. *JAMA Oncol* 2016; 1-11.

13. Sinicrope FA, Shi Q, Smyrk TC et al. Molecular markers identify subtypes of stage III colon cancer associated with patient outcomes. *Gastroenterology* 2015; 148: 88-99.
14. Taieb J, Le Malicot K, Shi Q et al. Prognostic value of BRAF and KRAS mutations in MSI and MSS subgroups of stage III colon cancer from adjuvant chemotherapy trials. *J Natl Cancer Inst* 2016; In press.
15. Janakiraman M, Vakiani E, Zeng Z et al. Genomic and biological characterization of exon 4 KRAS mutations in human cancer. *Cancer Res* 2010; 70: 5901-5911.
16. Karnoub AE, Weinberg RA. Ras oncogenes: split personalities. *Nat Rev Mol Cell Biol* 2008; 9: 517-531.
17. Fernandez-Medarde A, Santos E. Ras in cancer and developmental diseases. *Genes Cancer* 2011; 2: 344-358.
18. Forbes SA, Bindal N, Bamford S et al. COSMIC: mining complete cancer genomes in the Catalogue of Somatic Mutations in Cancer. *Nucleic Acids Res* 2011; 39: D945-950.
19. Peeters M, Oliner KS, Parker A et al. Massively parallel tumor multigene sequencing to evaluate response to panitumumab in a randomized phase III study of metastatic colorectal cancer. *Clin Cancer Res* 2013; 19: 1902-1912.
20. Bokemeyer C, Kohne CH, Ciardiello F et al. FOLFOX4 plus cetuximab treatment and RAS mutations in colorectal cancer. *Eur J Cancer* 2015; 51: 1243-1252.
21. Laurent-Puig P, Grisoni ML, Heinemann V et al. MiR 31 3p as a predictive biomarker of cetuximab efficacy effect in metastatic colorectal cancer (mCRC) patients enrolled in FIRE-3 study. *J Clin Oncol* 2016; 34: suppl; abstr 3516.
22. Jacobs B, De Roock W, Piessevaux H et al. Amphiregulin and epiregulin mRNA expression in primary tumors predicts outcome in metastatic colorectal cancer treated with cetuximab. *J Clin Oncol* 2009; 27: 5068-5074.
23. Gavin PG, Colangelo LH, Fumagalli D et al. Mutation profiling and microsatellite instability in stage II and III colon cancer: an assessment of their prognostic and oxaliplatin predictive value. *Clin Cancer Res* 2012; 18: 6531-6541.
24. Shen Y, Han X, Wang J et al. Prognostic impact of mutation profiling in patients with stage II and III colon cancer. *Sci Rep* 2016; 6: 24310.
25. Modest DP, Ricard I, Heinemann V et al. Outcome according to KRAS-, NRAS- and BRAF-mutation as well as KRAS mutation variants: pooled analysis of five randomized trials in metastatic colorectal cancer by the AIO colorectal cancer study group. *Ann Oncol* 2016; 27: 1746-1753.
26. Cremolini C, Di Bartolomeo M, Amatu A et al. BRAF codons 594 and 596 mutations identify a new molecular subtype of metastatic colorectal cancer at favorable prognosis. *Ann Oncol* 2015; 26: 2092-2097.

Figure legends

Figure 1:

Distribution of mutations.

Figure 2:

Kaplan Meier curves for time to relapse according to study treatment (A) in the *KRAS* exon 2 WT intention-to-treat population, (B) In patients with *RAS* WT and *BRAF* WT tumours, and (C) In patients with *RAS*-mutated tumours. TTR=time to recurrence. HR=hazard ratio.

Figure 3:

Prognostic impact of individual *RAS* and *BRAF* mutations on recurrence (A) and survival (B).

Supplementary Figure 1:

Flow chart of PETACC8 trial molecular study evaluating the impact of full *RAS* and *BRAF* wildtype mutations

Supplementary Figure 2:

Kaplan Meier curves for disease-free survival according to study treatment (A) in the *KRAS* exon 2 WT intention-to-treat population, (B) In patients with *RAS* WT and *BRAF* WT tumours, and (C) In patients with *RAS*-mutated tumours. DFS=Disease-Free Survival. HR=hazard ratio.

Table 1- Baseline patient and tumour characteristics in the *RAS* mutant, *BRAF* mutant and double wildtype subpopulations


		Double WT		<i>RAS</i> Mutant		<i>BRAF</i> Mutant	
		Folfox	Folfox+Cetux	Folfox	Folfox+Cetux	Folfox	Folfox+Cetux
Gender	<i>n</i>	367	352	484	484	99	114
	Male	224 (61.0%)	225 (63.9%)	255 (52.7%)	270 (55.8%)	49 (49.5%)	55 (48.2%)
	Female	143 (39.0%)	127 (36.1%)	229 (47.3%)	214 (44.2%)	50 (50.5%)	59 (51.8%)
Age	<i>n</i>	367	352	484	484	99	114
	Mean (SD)	58.83 (9.21)	58.33 (10.12)	60.03 (9.47)	59.68 (9.37)	60.85 (8.86)	59.78 (9.26)
	Median	60.00	60.00	61.00	61.00	62.00	60.00
	Q1; Q3	53.00; 66.00	52.00; 66.00	54.00; 68.00	54.00; 67.00	54.00; 68.00	53.00; 67.00
	Range	25.00; 75.00	19.00; 75.00	25.00; 75.00	23.00; 74.00	28.00; 73.00	27.00; 74.00
Age	<i>n</i>	367	352	484	484	99	114
	Age <= 70 years	336 (91.6%)	318 (90.3%)	425 (87.8%)	429 (88.6%)	89 (89.9%)	99 (86.8%)
	Age > 70 years	31 (8.4%)	34 (9.7%)	59 (12.2%)	55 (11.4%)	10 (10.1%)	15 (13.2%)
	Missing	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
WHO Performance Status	<i>n</i>	355	344	463	465	96	108
	0	293 (82.5%)	284 (82.6%)	387 (83.6%)	380 (81.7%)	73 (76.0%)	83 (76.9%)
	1	60 (16.9%)	60 (17.4%)	75 (16.2%)	83 (17.8%)	23 (24.0%)	25 (23.1%)
	2	2 (0.6%)	0 (0.0)	1 (0.2%)	2 (0.4%)	.	.
Tumour Location	<i>n</i>	366	352	479	483	99	114
	Left	276 (75.4%)	260 (73.9%)	261 (54.5%)	254 (52.6%)	31 (31.3%)	32 (28.1%)
	Right	88 (24.0%)	91 (25.9%)	211 (44.1%)	216 (44.7%)	67 (67.7%)	82 (71.9%)
	Both sides	2 (0.5%)	1 (0.3%)	7 (1.5%)	13 (2.7%)	1 (1.0%)	0 (0.0)
Tumour grade	<i>n</i>	367	352	483	484	99	114
	Missing	5 (1.4%)	3 (0.9%)	7(1.4%)	6 (1.2%)	1 (1.0%)	2 (1.8%)
	Well differentiated	79 (21.5%)	80 (22.7%)	95 (19.7%)	102 (21.1%)	14 (14.1%)	16 (14.0%)
	Moderately differentiated	227 (61.9%)	207 (58.8%)	295 (61.1%)	298 (61.6%)	50 (50.5%)	54 (47.4%)
	Poorly differentiated	54 (14.7%)	60 (17.0%)	84 (17.4%)	76 (15.7%)	34 (34.3%)	40 (35.1%)
	Undifferentiated	2 (0.5%)	2 (0.6%)	2 (0.4%)	2 (0.4%)	0 (0.0)	2 (1.8%)
pN stage	<i>n</i>	367	352	484	484	99	114
	pN1	240 (65.4%)	224 (63.6%)	304 (62.8%)	305 (63.0%)	60 (60.6%)	58 (50.9%)
	pN2	127 (34.6%)	128 (36.4%)	180 (37.2%)	179 (37.0%)	39 (39.4%)	56 (49.1%)

		Double WT		RAS Mutant		BRAF Mutant	
		Folfox	Folfox+Cetux	Folfox	Folfox+Cetux	Folfox	Folfox+Cetux
pT stage	<i>n</i>	367	352	484	484	99	114
	pT1	11 (3.0%)	10 (2.8%)	14 (2.9%)	10 (2.1%)	1 (1.0%)	2 (1.8%)
	pT2	32 (8.7%)	30 (8.5%)	29 (6.0%)	30 (6.2%)	5 (5.1%)	3 (2.6%)
	pT3	259 (70.6%)	229 (65.1%)	332 (68.6%)	346 (71.5%)	72 (72.7%)	84 (73.7%)
	pT4	65 (17.7%)	83 (23.6%)	109 (22.5%)	98 (20.2%)	21 (21.2%)	24 (21.1%)
	pTis	0 (0.0)	1 (0.9%)
Bowel obstruction and perforation	<i>n</i>	367	352	484	484	99	114
	Bowel obstruction and/or perforation	64 (17.4%)	65 (18.5%)	99 (20.5%)	97 (20.0%)	17 (17.2%)	18 (15.8%)
	No bowel obstruction and no perforation	303 (82.6%)	287 (81.5%)	385 (79.5%)	387 (80.0%)	82 (82.8%)	96 (84.2%)
VELI	<i>n</i>	367	352	484	484	99	114
	Vascular Invasion or Lymphatic infiltration	210 (57.2%)	207 (58.8%)	262 (54.1%)	248 (51.2%)	67 (67.7%)	62 (54.4%)
	No Vascular Invasion and no Lymphatic infiltration	104 (28.3%)	95 (27.0%)	144 (29.8%)	154 (31.8%)	24 (24.2%)	29 (25.4%)
MMR Status	N	340	324	406	390	89	105
	pMMR	309 (90.9%)	301 (92.9%)	377 (92.9%)	372 (95.4%)	56 (62.9%)	71 (67.6%)
	dMMR	31 (9.1%)	23 (7.1%)	29 (7.1%)	18 (4.6%)	33 (37.1%)	34 (32.4%)

Abbreviations: MMR, mismatch repair; pMMR, proficient MMR; dMMR, deficient MMR


Table 2: *RAS* and *BRAF* mutations

	<i>RAS</i> Mutant (n=968)	<i>BRAF</i> Mutant (n=213)
<i>KRAS</i> mutations		
Exon 2	783 (80.9%)	-
Codon 12	594 (75.9%)	-
Codon 13	178 (18.4%)	-
Other	11 (1.4%)	-
Exon 3	42 (4.3%)	-
Codon 59	8 (19.0%)	-
Codon 61	34 (81.0%)	-
Exon 4	80 (8.3%)	-
Codon 146	68 (85.0%)	-
Codon 117	11 (13.8%)	-
Other	1 (1.2%)	-
<i>NRAS</i> mutations		
Exon 2	30 (3.1%)	-
Codon 12	26 (86.7%)	-
Codon 13	4 (13.3%)	-
Exon 3	31 (3.2%)	-
Exon 4	2 (0.2%)	-
<i>BRAF</i> mutations	-	
V600E	-	192 (90.1%)
Other mutations	-	21 (9.9%)



Distribution of mutations.


304x171mm (96 x 96 DPI)



	0	1	2	3	4	5
Folfox	556	513	451	416	377	275
Folfox+ Cetux	557	500	440	406	360	272

Kaplan Meier curves for time to relapse according to study treatment (A) in the KRAS exon 2 WT intention-to-treat population


272x160mm (144 x 144 DPI)



Folfox	367	340	299	275	249	186
Folfox+ Cetux	352	329	297	277	244	185


Kaplan Meier curves for time to relapse according to study treatment (B) In patients with RAS WT and BRAF WT tumours,

292x162mm (144 x 144 DPI)




Kaplan Meier curves for time to relapse according to study treatment (C) In patients with RAS-mutated tumours

292x160mm (144 x 144 DPI)




Prognostic impact of individual RAS and BRAF mutations on recurrence (A)

349x209mm (96 x 96 DPI)




Prognostic impact of individual RAS and BRAF mutations on overall Survival (B)

349x209mm (96 x 96 DPI)




169x127mm (144 x 144 DPI)




	0	1	2	3	4	5
Folfox	556	512	449	413	374	273
Folfox+ Cetux	557	497	438	402	356	270

408x244mm (96 x 96 DPI)



	0	1	2	3	4	5
Folfox	367	340	299	275	248	186
Folfox+ Cetux	352	327	296	273	240	183

408x244mm (96 x 96 DPI)



	0	1	2	3	4	5
Folfox	484	427	375	344	303	219
Folfox+ Cetux	484	421	349	325	273	218

408x244mm (96 x 96 DPI)

Supplementary Table 1. Study population compared to all patients enrolled in the PETACC8 trial.

		Full RAS patients (N=1900)	PETACC8 patients (N=2559)	P-value
Gender	<i>n</i>	1900	2559	χ^2 : 0.7923
	Male	1078 (56.7%)	1462 (57.1%)	.
	Female	822 (43.3%)	1097 (42.9%)	.
Treatment Group	<i>n</i>	1900	2559	χ^2 : 0.9897
	Folfox	950 (50.0%)	1279 (50.0%)	.
	Folfox+Cetux	950 (50.0%)	1280 (50.0%)	.
Age	<i>n</i>	1900	2559	<i>W</i> : 0.4962
	Mean (SD)	59.42 (9.49)	59.19 (9.67)	.
	Median	61.00	60.00	.
	Q1; Q3	54.00; 67.00	53.00; 67.00	.
	Range	19.00; 75.00	19.00; 75.00	.
Age	<i>n</i>	1900	2559	χ^2 : 0.8422
	Age <= 70 years	1696 (89.3%)	2289 (89.4%)	.
	Age > 70 years	204 (10.7%)	270 (10.6%)	.
	Missing	0 (0.0)	0 (0.0)	.
WHO performance status	<i>n</i>	1831	2441	χ^2 : 0.8073
	0	1500 (81.9%)	2004 (82.1%)	.
	1	326 (17.8%)	431 (17.7%)	.
	2	5 (0.3%)	5 (0.2%)	.
	3	0 (0.0)	1 (0.0%)	.
Tumour Location	<i>n</i>	1893	2546	χ^2 : 0.3069
	Left	1114 (58.8%)	1552 (61.0%)	.
	Right	755 (39.9%)	968 (38.0%)	.
	Both sides	24 (1.3%)	26 (1.0%)	.

		Full RAS patients	PETACC8 patients	P-value
		(N=1900)	(N=2559)	
Tumour grade	<i>n</i>	1899	2557	χ^2 : 0.9978
	.A	1 (0.1%)	1 (0.0%)	.
	.D	3 (0.2%)	5 (0.2%)	.
	.K	20 (1.1%)	25 (1.0%)	.
	Well differentiated	386 (20.3%)	527 (20.6%)	.
	Moderately differentiated	1131 (59.6%)	1527 (59.7%)	.
	Poorly differentiated	348 (18.3%)	461 (18.0%)	.
	Undifferentiated	10 (0.5%)	11 (0.4%)	.
pN grade	<i>n</i>	1900	2559	χ^2 : 0.8501
	pN1	1191 (62.7%)	1597 (62.4%)	.
	pN2	709 (37.3%)	962 (37.6%)	.
pT grade	<i>n</i>	1900	2559	χ^2 : 0.5889
	pT1	48 (2.5%)	71 (2.8%)	.
	pT2	129 (6.8%)	194 (7.6%)	.
	pT3	1322 (69.6%)	1768 (69.1%)	.
	pT4	400 (21.1%)	522 (20.4%)	.
	pTis	1 (0.1%)	1 (0.0%)	.
	pTx	0 (0.0)	3 (0.1%)	.
Bowel obstruction and perforation	<i>n</i>	1900	2559	χ^2 : 0.7152
	Bowel obstruction and/or perforation	360 (18.9%)	496 (19.4%)	.
	No bowel obstruction and no perforation	1540 (81.1%)	2063 (80.6%)	.
VELI	<i>n</i>	1900	2559	χ^2 : 0.8758
	Vascular Invasion or Lymphatic infiltration	1056 (55.6%)	1442 (56.4%)	.
	No Vascular Invasion and no Lymphatic infiltration	550 (28.9%)	729 (28.5%)	.

Abbreviations: MMR, mismatch repair; pMMR, proficient MMR; dMMR, deficient MMR ; χ^2 : ; Chi-Square Test; W: Wilcoxon Test