

## NUMERICAL ANALYSIS OF NONLINEAR SUBDIFFUSION EQUATIONS\*

BANGTI JIN<sup>†</sup>, BUYANG LI<sup>‡</sup>, AND ZHI ZHOU<sup>‡</sup>

**Abstract.** We present a general framework for the rigorous numerical analysis of time-fractional nonlinear parabolic partial differential equations, with a fractional derivative of order  $\alpha \in (0, 1)$  in time. It relies on three technical tools: a fractional version of the discrete Grönwall type inequality, discrete maximal regularity, and regularity theory of nonlinear equations. We establish a general criterion for showing the fractional discrete Grönwall inequality and verify it for the L1 scheme and convolution quadrature generated by backward difference formulas. Further, we provide a complete solution theory, e.g., existence, uniqueness, and regularity, for a time-fractional diffusion equation with a Lipschitz nonlinear source term. Together with the known results of discrete maximal regularity, we derive pointwise  $L^2(\Omega)$  norm error estimates for semidiscrete Galerkin finite element solutions and fully discrete solutions, which are of order  $O(h^2)$  (up to a logarithmic factor) and  $O(\tau^\alpha)$ , respectively, without any extra regularity assumption on the solution or compatibility condition on the problem data. The sharpness of the convergence rates is supported by the numerical experiments.

**Key words.** nonlinear fractional diffusion equation, discrete fractional Grönwall inequality, L1 scheme, convolution quadrature, error estimate

**AMS subject classifications.** 65M15, 65M60, 65M12, 45K05

**DOI.** 10.1137/16M1089320

**1. Introduction.** Time-fractional parabolic partial differential equations (PDEs) have been very popular for modeling anomalously slow transport processes in the past two decades. These models are commonly referred to as fractional diffusion or subdiffusion. At a microscopic level, the underlying stochastic process is a continuous time random walk [32]. So far they have been successfully applied in a broad range of diversified research areas, e.g., thermal diffusion in fractal domains [35], flow in highly heterogeneous aquifers [6], and single-molecular protein dynamics [20], just to name a few. Hence, the rigorous numerical analysis of such problems is of great practical importance. For the linear problem, various efficient time-stepping schemes have been proposed, which include mainly two classes: L1 type schemes and convolution quadrature (CQ).

L1 type schemes approximate the fractional derivative by replacing the integrand with its piecewise polynomial interpolation [24, 26, 37, 3] and thus generalize the classical finite difference method. The piecewise linear case has a local truncation error  $O(\tau^{2-\alpha})$  for sufficiently smooth solution, where  $\tau$  denotes the time step size. See also [31, 33] for the discontinuous Galerkin method. CQ is a flexible framework introduced by Lubich [27, 28] for constructing high-order time discretization methods

---

\*Received by the editors August 12, 2016; accepted for publication (in revised form) June 23, 2017; published electronically January 5, 2018.

<http://www.siam.org/journals/sinum/56-1/M108932.html>

**Funding:** The work of the first author was partially supported by UK EPSRC grant EP/M025160/1. The work of the second author was partially supported by a grant from the Research Grants Council of the Hong Kong Special Administrative Region (project 15300817). The work of the third author was partially supported by the AFOSR MURI Center for Material Failure Prediction Through Peridynamics and ARO MURI grant W911NF-15-1-0562.

<sup>†</sup>Department of Computer Science, University College London, London, WC1E 6BT, UK (b.jin@ucl.ac.uk, bangti.jin@gmail.com).

<sup>‡</sup>Department of Applied Mathematics, Hong Kong Polytechnic University, Kowloon, Hong Kong (buyang.li@polyu.edu.hk, libuyang@gmail.com, zhizhou@polyu.edu.hk, zhizhou0125@gmail.com).

for approximating fractional derivatives. It approximates the fractional derivative in the Laplace domain and automatically inherits the stability property of general linear multistep methods. See [10, 39, 40, 16] for CQ type schemes. Optimal error estimates have been derived for both spatially semidiscrete and fully discrete schemes, including problems with nonsmooth data [10, 14, 31, 16].

However, up to now, there has been very little work on the rigorous numerical analysis of nonlinear time-fractional diffusion equations. In this paper, we present a general framework for analyzing discretization errors of nonlinear problems. The error of the numerical solution can be split into a linear part and a nonlinear part. While the linear part has been carefully studied, the analysis of the nonlinear part requires different mathematical machineries in order to derive sharp error estimates. Besides regularity estimates for the nonlinear problem, it requires discrete maximal  $\ell^p$  regularity and a fractional version of the discrete Grönwall's inequality for time-stepping schemes. The former gives a bound on the discrete fractional derivative due to the nonlinear part, whereas the latter allows combining the nonlinear part with the linear part to obtain a global error estimate.

To the best of our knowledge, a fractional version of the discrete Grönwall's inequality for time-stepping schemes is still unavailable in the literature. We shall establish such a discrete Grönwall's inequality for both the L1 scheme and CQs generated by backward difference formulas (BDFs) up to order 6 in Theorem 2.8. Further, in Theorem 2.7, we present a general criterion under which the fractional discrete Grönwall's inequality holds.

To illustrate the main idea of this framework, we consider the following nonlinear problem in a bounded convex polygonal domain  $\Omega \subset \mathbb{R}^d$ ,  $d \geq 1$ :

$$(1.1) \quad \begin{cases} \partial_t^\alpha u - \Delta u = f(u) & \text{in } \Omega \times (0, T), \\ u = 0 & \text{on } \partial\Omega \times (0, T), \\ u = u_0 & \text{in } \Omega \times \{0\}, \end{cases}$$

where  $u_0 \in H_0^1(\Omega) \cap H^2(\Omega)$  is a given function and  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a Lipschitz continuous function, i.e.,  $|f(s) - f(t)| \leq L|s - t|$  for all  $s, t \in \mathbb{R}$ , and  $\partial_t^\alpha u$  denotes the Caputo fractional derivative of order  $\alpha \in (0, 1)$  in time [19, p. 91],

$$(1.2) \quad \partial_t^\alpha u(t) := \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-s)^{-\alpha} \frac{d}{ds} u(s) ds \quad \text{with } \Gamma(z) := \int_0^\infty s^{z-1} e^{-s} ds.$$

Let  $S_h \subset H_0^1(\Omega)$  be the continuous piecewise linear finite element space subject to a quasi-uniform shape regular triangulation of  $\Omega$ , with a mesh size  $h$ , and let  $\Delta_h : S_h \rightarrow S_h$  denote the Galerkin finite element approximation of the Dirichlet Laplacian  $\Delta$ , defined by

$$(\Delta_h w_h, v_h) := -(\nabla w_h, \nabla v_h) \quad \forall w_h, v_h \in S_h.$$

Let  $0 = t_0 < t_1 < \dots < t_N = T$  be a uniform partition of the time interval  $[0, T]$  with grid points  $t_n = n\tau$  and step size  $\tau = T/N$ . Upon rewriting the Caputo derivative  $\partial_t^\alpha u$  as a Riemann–Liouville one [19, p. 91], we consider a linearized time-stepping scheme: for the given initial value  $u_h^0 = R_h u_0$  (Ritz projection of  $u_0$ ), find  $u_h^n$ ,  $n = 1, 2, \dots, N$ , such that

$$(1.3) \quad \bar{\partial}_\tau^\alpha (u_h^n - u_h^0) - \Delta_h u_h^n = P_h f(u_h^{n-1}),$$

where  $P_h$  denotes the  $L^2$  projection onto the finite element space  $S_h$ , and  $\bar{\partial}_\tau^\alpha u_h^n$  denotes either the CQ generated by the backward Euler method or the L1 scheme; see (2.8) and (2.9) below. These methods are popular for discretizing the fractional derivative in time.

After proving the fractional discrete Grönwall’s inequality in section 2 and the regularity estimate in section 3, we present an error analysis for the fully discrete scheme (1.3) in section 4. By introducing an intermediate spatially semidiscrete Galerkin problem

$$(1.4) \quad \partial_t^\alpha u_h(t) - \Delta_h u_h(t) = P_h f(u_h(t)) \quad \forall t \in (0, T],$$

we split the error into two parts,  $u(t_n) - u_h^n = (u(t_n) - u_h(t_n)) + (u_h(t_n) - u_h^n)$ , and derive the following error estimates for each component in Theorems 4.4 and 4.5:

$$\max_{0 \leq t \leq T} \|u(t) - u_h(t)\|_{L^2(\Omega)} \leq c \ell_h^2 h^2 \quad \text{and} \quad \max_{1 \leq n \leq N} \|u_h(t_n) - u_h^n\|_{L^2(\Omega)} \leq c \tau^\alpha,$$

where  $\ell_h = \log(2 + 1/h)$ . These estimates are sharp with respect to the regularity of the solution in Theorem 3.1 (up to a logarithmic factor  $\ell_h$ ) and are confirmed by the numerical experiments in section 6. Besides, we show how to simplify the analysis of nonlinear problems by applying the fractional type discrete maximal  $\ell^p$ -regularity established in [17], an extension of the discrete maximal  $\ell^p$ -regularity of standard parabolic equations [18, 21, 25], which has been applied to numerical analysis of nonlinear parabolic equations in the literature [1, 2, 22].

Last we mention the interesting works [10, 34] on integro-differential equations, where a Riemann–Liouville fractional integral operator appears in front of the Laplacian. These models are closely related to (1.1) but have different smoothing properties. Cuesta, Lubich, and Palencia [10] proposed the CQ generated by the second-order BDF for a semilinear problem and proved an  $O(\tau^2)$  error bound of the temporal error. In [34], a Crank–Nicolson type method for a semilinear problem with variable time step size was studied. In these works, a variant of the discrete Grönwall’s inequality due to Chen, Thomée, and Wahlbin [8] plays a crucial role, which differs substantially from the discrete Grönwall’s inequality we shall establish below.

Throughout this paper, the notation  $c$  denotes a generic constant, which may vary at different occurrences, but it is always independent of the mesh size  $h$  and time step size  $\tau$ .

**2. Discrete Grönwall’s inequality for time-fractional diffusion.** In this section, we establish a fractional version of Grönwall’s inequality and its discrete analogue for time-stepping schemes. These inequalities are crucial in analyzing numerical schemes for nonlinear subdiffusion equations and are of independent interest.

**2.1. Continuous Grönwall’s inequality.** We begin with the continuous Grönwall’s inequality for fractional differential equations in a general Banach space setting.

**THEOREM 2.1** (fractional Grönwall’s inequality). *Let  $X$  be any given Banach space. For  $\alpha \in (0, 1)$  and  $p \in (1/\alpha, \infty)$ , if a function  $u \in C([0, T]; X)$  satisfies  $\partial_t^\alpha u \in L^p(0, T; X)$ ,  $u(0) = 0$ , and*

$$(2.1) \quad \|\partial_t^\alpha u\|_{L^p(0,s;X)} \leq \kappa \|u\|_{L^p(0,s;X)} + \sigma \quad \forall s \in (0, T],$$

for some positive constants  $\kappa$  and  $\sigma$ , then

$$(2.2) \quad \|u\|_{C([0,T];X)} + \|\partial_t^\alpha u\|_{L^p(0,T;X)} \leq c\sigma,$$

where the constant  $c$  is independent of  $\sigma$ ,  $u$ , and  $X$  but may depend on  $\alpha$ ,  $p$ ,  $\kappa$ , and  $T$ .

*Proof.* Due to the zero initial condition  $u(0) = 0$ , the Riemann–Liouville and Caputo fractional derivatives coincide. Hence, the function  $u(t)$  can be expressed in terms of  $\partial_t^\alpha u$  (cf. [19, Lemma 2.22, p. 96] and [19, Lemma 2.5, p. 74]):  $u(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-\xi)^{\alpha-1} \partial_\xi^\alpha u(\xi) d\xi$ . Since  $p > 1/\alpha$ , Hölder’s inequality implies

$$(2.3) \quad \|u(t)\|_X \leq c \left( \int_0^t (t-\xi)^{\frac{(\alpha-1)p}{p-1}} d\xi \right)^{\frac{p-1}{p}} \|\partial_\xi^\alpha u\|_{L^p(0,t;X)} \leq c \|\partial_\xi^\alpha u\|_{L^p(0,t;X)}.$$

Upon taking the supremum with respect to  $t \in (0, s)$  for any  $s \in (0, T]$  in (2.3), we obtain

$$\begin{aligned} \|u\|_{L^\infty(0,s;X)} &\leq c \|\partial_\xi^\alpha u\|_{L^p(0,s;X)} \leq c\kappa \|u\|_{L^p(0,s;X)} + c\sigma \\ &\leq \epsilon\kappa \|u\|_{L^\infty(0,s;X)} + c_\epsilon\kappa \|u\|_{L^1(0,s;X)} + c\sigma \quad \forall s \in [0, T], \end{aligned}$$

where  $\epsilon > 0$  can be arbitrary. By choosing  $\epsilon = \frac{1}{2\kappa}$ , the  $L^\infty$ -norm on the right-hand side can be eliminated by the left-hand side, and the last inequality reduces to

$$\|u\|_{L^\infty(0,s;X)} \leq c_\kappa \|u\|_{L^1(0,s;X)} + c\sigma \quad \forall s \in [0, T].$$

That is, we have  $\|u(s)\|_X \leq c_\kappa \int_0^s \|u(\xi)\|_X d\xi + c\sigma$  for  $s \in (0, T]$ . Now the standard Grönwall’s inequality yields

$$\max_{s \in [0, T]} \|u(s)\|_X \leq e^{c_\kappa T} c\sigma.$$

Substituting it into (2.1) yields (2.2). The proof of Theorem 2.1 is complete.  $\square$

**2.2. Discrete Grönwall’s inequality.** In this part, we establish the discrete analogue of the Grönwall’s inequality in Theorem 2.1 for time-stepping schemes that approximate the fractional derivative  $\partial_t^\alpha v(t_n)$  by a discrete convolution:

$$(2.4) \quad \bar{\partial}_\tau^\alpha v^n := \frac{1}{\tau^\alpha} \sum_{j=0}^n K_{n-j} v^j, \quad n = 0, 1, 2, \dots,$$

where  $v^n$  is an approximation of  $v(t_n)$ , and  $K_j$ ,  $j = 0, 1, 2, \dots$ , are the weights independent of the time step size  $\tau$ . Throughout, we denote by  $K(\zeta)$  the generating function of the discrete fractional derivative  $\bar{\partial}_\tau^\alpha$ , defined by

$$(2.5) \quad K(\zeta) := \frac{1}{\tau^\alpha} \sum_{j=0}^{\infty} K_j \zeta^j,$$

which is an analytic function in the (open) unit disk  $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$ , continuously differentiable up to the boundary  $\partial\mathbb{D} \setminus \{\pm 1\}$ , except for the two points  $\pm 1$ . Then we have

$$(2.6) \quad K(\zeta) \sum_{n=0}^{\infty} v^n \zeta^n = \sum_{n=0}^{\infty} (\bar{\partial}_\tau^\alpha v^n) \zeta^n.$$

*Example 2.2.* The CQ generated by the  $k$ th-order BDF [27, 10] is given by (2.4), where the coefficients  $K_j$ ,  $j = 0, 1, \dots$ , are determined by the power series expansion

$$(2.7) \quad \left( \sum_{j=1}^k \frac{1}{j} (1 - \zeta)^j \right)^\alpha = \sum_{j=0}^\infty K_j \zeta^j.$$

The special case  $k = 1$ , i.e., the backward Euler CQ, is very popular and commonly known as the Grünwald–Letnikov approximation, and the coefficients  $K_j$ ,  $j = 0, 1, 2, \dots$ , are given by

$$(2.8) \quad (1 - \zeta)^\alpha = \sum_{j=0}^\infty K_j \zeta^j.$$

*Example 2.3.* The popular L1 scheme [26] is also of the form (2.4) with [17, p. 8]

$$(2.9) \quad \frac{(1 - \zeta)^2}{\zeta \Gamma(2 - \alpha)} \text{Li}_{\alpha-1}(\zeta) = \sum_{j=0}^\infty K_j \zeta^j,$$

where  $\text{Li}_p(z) = \sum_{j=1}^\infty z^j/j^p$  is the polylogarithmic function, which is well defined for  $|z| < 1$  and can be analytically continued to the split complex plane  $\mathbb{C} \setminus [1, \infty)$  [11].

Now we turn to the discrete Grönwall’s inequality. For  $1 \leq p \leq \infty$ , we denote by  $\ell^p(X)$  the space of sequences  $v^n \in X$ ,  $n = 0, 1, \dots$ , such that  $\|(v^n)_{n=0}^\infty\|_{\ell^p(X)} < \infty$ , where

$$\|(v^n)_{n=0}^\infty\|_{\ell^p(X)} := \begin{cases} \left( \sum_{n=0}^\infty \tau \|v^n\|_X^p \right)^{\frac{1}{p}} & \text{if } 1 \leq p < \infty, \\ \sup_{n \geq 0} \|v^n\|_X & \text{if } p = \infty. \end{cases}$$

For a finite sequence  $v^n \in X$ ,  $n = 0, 1, \dots, m$ , we denote  $\|(v^n)_{n=0}^m\|_{\ell^p(X)} := \|(v^n)_{n=0}^\infty\|_{\ell^p(X)}$ , by setting  $v^n = 0$  for  $n > m$ . The following theorem is a discrete analogue of Theorem 2.1 for the backward Euler CQ. It is foundational to the proof of the discrete Grönwall’s inequalities for other time-stepping schemes.

**THEOREM 2.4** (discrete fractional Grönwall’s inequality: Backward Euler). *Let  $X$  be any given Banach space, and let  $\bar{\partial}_\tau^\alpha$  denote the backward Euler CQ given by (2.4) and (2.8). If  $\alpha \in (0, 1)$  and  $p \in (1/\alpha, \infty)$ , and a sequence  $v^n \in X$ ,  $n = 0, 1, 2, \dots$ , with  $v^0 = 0$ , satisfies*

$$(2.10) \quad \|(\bar{\partial}_\tau^\alpha v^n)_{n=1}^m\|_{\ell^p(X)} \leq \kappa \|(v^n)_{n=1}^m\|_{\ell^p(X)} + \sigma \quad \forall 0 \leq m \leq N$$

for some positive constants  $\kappa$  and  $\sigma$ , then there exists a  $\tau_0 > 0$  such that for any  $\tau < \tau_0$  there holds

$$(2.11) \quad \|(v^n)_{n=1}^N\|_{\ell^\infty(X)} + \|(\bar{\partial}_\tau^\alpha v^n)_{n=1}^N\|_{\ell^p(X)} \leq c\sigma,$$

where the constants  $c$  and  $\tau_0$  are independent of  $\sigma$ ,  $\tau$ ,  $N$ ,  $X$ , and  $v^n$  but may depend on  $\alpha$ ,  $p$ ,  $\kappa$ , and  $T$ .

To prove Theorem 2.4, we need a technical lemma, which gives a discrete analogue of the Hardy type inequality (2.3).

LEMMA 2.5 (discrete Hardy type inequality). *Let  $\alpha \in (0, 1)$ , and let  $X$  be any given Banach space. If  $v^n \in X$  and  $w^n \in X$ ,  $n = 0, 1, 2, \dots$ , satisfy*

$$(2.12) \quad \left(\frac{1-\zeta}{\tau}\right)^\alpha \sum_{n=0}^{\infty} v^n \zeta^n = \sum_{n=0}^{\infty} w^n \zeta^n,$$

*in the sense that both sides are analytic in  $\mathbb{D}$ , then for  $p \in (1/\alpha, \infty)$ , there holds*

$$(2.13) \quad \|(v^n)_{n=0}^m\|_{\ell^\infty(X)} \leq c \|(w^n)_{n=0}^m\|_{\ell^p(X)}, \quad 0 \leq m \leq N,$$

*where the constant  $c$  is independent of  $\tau$ ,  $m$ ,  $N$ , and  $X$  but may depend on  $\alpha$ ,  $p$ , and  $T$ .*

*Proof.* We define  $\phi^n$ ,  $n = 0, 1, \dots$ , to be the coefficients of the power series expansion

$$(1-\zeta)^{-\alpha} = \sum_{n=0}^{\infty} \phi^n \zeta^n.$$

Then direct calculations yield  $\phi^0 = 1$  and  $\phi^n = \prod_{j=1}^n (1 + \frac{\alpha-1}{j})$  for  $n \geq 1$ . By the trivial inequality  $\ln(1+x) \leq x$  for  $x > -1$ , we have

$$\ln \phi^n = \sum_{j=1}^n \ln \left(1 + \frac{\alpha-1}{j}\right) \leq (\alpha-1) \sum_{j=1}^n j^{-1} \leq (\alpha-1) \ln(n+1).$$

That is,  $\phi^n \leq (n+1)^{\alpha-1}$  for  $n \geq 0$ . It follows from (2.12) that

$$\sum_{n=0}^{\infty} v^n \zeta^n = \left(\frac{\tau}{1-\zeta}\right)^\alpha \sum_{n=0}^{\infty} w^n \zeta^n = \tau^\alpha \left(\sum_{n=0}^{\infty} \phi^n \zeta^n\right) \left(\sum_{n=0}^{\infty} w^n \zeta^n\right).$$

With  $p' = \frac{p}{p-1}$ , the last identity yields

$$(2.14) \quad \begin{aligned} \|v^n\|_X &= \left\| \tau^\alpha \sum_{j=0}^n \phi^{n-j} w^j \right\|_X \leq \tau^\alpha \left( \sum_{j=0}^n |\phi^{n-j}|^{p'} \right)^{\frac{1}{p'}} \left( \sum_{j=0}^n \|w^j\|_X^p \right)^{\frac{1}{p}} \\ &\leq \tau^{\alpha-1/p} \left( \sum_{j=0}^n \frac{1}{(j+1)^{p'(1-\alpha)}} \right)^{\frac{1}{p'}} \|(w^j)_{j=0}^n\|_{\ell^p(X)}. \end{aligned}$$

If  $p > 1/\alpha$ , then  $0 < p'(1-\alpha) < 1$  and so

$$\sum_{j=0}^n \frac{1}{(j+1)^{p'(1-\alpha)}} \leq \int_0^{n+1} \frac{ds}{s^{p'(1-\alpha)}} = \frac{(n+1)^{1-p'(1-\alpha)}}{1-p'(1-\alpha)}.$$

Hence, (2.14) reduces to

$$\|v^n\|_X \leq \tau^{\alpha-1/p} \frac{(n+1)^{\alpha-1/p}}{(1-p'(1-\alpha))^{1/p'}} \|(w^j)_{j=0}^n\|_{\ell^p(X)} \leq \frac{(2T)^{\alpha-1/p}}{(1-p'(1-\alpha))^{1/p'}} \|(w^j)_{j=0}^n\|_{\ell^p(X)},$$

where we have used the fact  $\tau(n+1) \leq 2T$  in the last inequality. Since the last inequality holds for all  $n = 0, \dots, m$ , it follows that (2.13) holds.  $\square$

Now we are ready to prove Theorem 2.4.

*Proof of Theorem 2.4.* For the backward Euler CQ we have  $K(\zeta) = (\frac{1-\zeta}{\tau})^\alpha$ . Since,  $v^0 = 0$ ,  $\bar{\partial}_\tau^\alpha v^0 = 0$ , and the identity (2.6) can be written as  $(\frac{1-\zeta}{\tau})^\alpha \sum_{n=0}^\infty v^n \zeta^n = \sum_{n=0}^\infty (\bar{\partial}_\tau^\alpha v^n) \zeta^n$ . Then Lemma 2.5 and (2.10) imply

$$\begin{aligned} \|(v^n)_{n=0}^m\|_{\ell^\infty(X)} &\leq c\|(\bar{\partial}_\tau^\alpha v^n)_{n=0}^m\|_{\ell^p(X)} \leq c\kappa\|(v^n)_{n=0}^m\|_{\ell^p(X)} + c\sigma \\ &\leq \epsilon\kappa\|(v^n)_{n=0}^m\|_{\ell^\infty(X)} + c_\epsilon\kappa\|(v^n)_{n=0}^m\|_{\ell^1(X)} + c\sigma \quad \forall 1 \leq m \leq N. \end{aligned}$$

By choosing  $\epsilon\kappa = 1/2$  and collecting terms, and using the fact  $v^0 = 0$ , we obtain

$$\|(v^n)_{n=1}^m\|_{\ell^\infty(X)} \leq c_\kappa\|(v^n)_{n=1}^m\|_{\ell^1(X)} + c\sigma \quad \forall 1 \leq m \leq N.$$

That is,  $\|v^m\|_X \leq c_\kappa\tau \sum_{n=1}^m \|v^n\|_X + c\sigma$  for  $1 \leq m \leq N$ . Then the standard discrete Grönwall's inequality gives, for sufficiently small step size  $\tau$ ,

$$\max_{1 \leq n \leq N} \|v^n\|_X \leq e^{c_\kappa T} c\sigma.$$

Substituting this into (2.10) yields (2.11). The proof of Theorem 2.4 is complete.  $\square$

To analyze other time-stepping schemes, we shall need the following lemma of discrete Mihlin multipliers, which is a simple consequence of Blunck's multiplier theorem [7, Theorem 1.3] through the transform  $\zeta = e^{-i\theta}$ . Here, an unconditional martingale difference (UMD) space  $X$  denotes a Banach space such that the Hilbert transform  $Hf(t) := \int_{\mathbb{R}} \frac{f(s)}{t-s} ds$  is bounded on  $L^p(\mathbb{R}; X)$  for all  $1 < p < \infty$  [23]. Examples of UMD spaces include  $\mathbb{R}^d$ ,  $d \geq 1$ , and  $L^q(\Omega)$ ,  $1 < q < \infty$ , and their closed subspaces (e.g., the finite element space  $S_h$  equipped with the  $L^q(\Omega)$  norm).

LEMMA 2.6 (discrete Mihlin multipliers). *Let  $X$  be a UMD space and let  $M : \mathbb{D} \rightarrow \mathbb{C}$  be an analytic function, continuously differentiable up to  $\partial\mathbb{D} \setminus \{\pm 1\}$ , such that the set*

$$\{M(\zeta) : \zeta \in \partial\mathbb{D} \setminus \{\pm 1\}\} \cup \{(1-\zeta)(1+\zeta)M'(\zeta) : \zeta \in \partial\mathbb{D} \setminus \{\pm 1\}\}$$

*is bounded, and denote its bound by  $c_R$ . Then for any  $1 < p < \infty$  and any sequence  $(f^n)_{n=0}^\infty \in \ell^p(X)$ , the coefficients  $u_n \in X$ ,  $n = 0, 1, \dots$ , in the power series expansion*

$$M(\zeta) \sum_{n=0}^\infty f^n \zeta^n = \sum_{n=0}^\infty u^n \zeta^n \quad \forall \zeta \in \mathbb{D}$$

*satisfy*

$$\|(u^n)_{n=0}^\infty\|_{\ell^p(X)} \leq c_{p,X} c_R \|(f^n)_{n=0}^\infty\|_{\ell^p(X)},$$

*where the constant  $c_{p,X}$  is independent of the operators  $M(\zeta)$ ,  $\zeta \in \mathbb{D}$ .*

Now other time-stepping schemes can be connected to the backward Euler CQ. The next result gives a general criterion for the discrete fractional Grönwall's inequality.

THEOREM 2.7 (general criterion for discrete fractional Grönwall's inequality). *Let  $X$  be a UMD space. If the generating function  $K(\zeta) = \frac{1}{\tau^\alpha} \sum_{n=0}^\infty K_n \zeta^n$  satisfies*

(2.15)

$$|K(\zeta)| \geq \frac{1}{c} \left| \frac{1-\zeta}{\tau} \right|^\alpha \quad \text{and} \quad |(1-\zeta)(1+\zeta)K'(\zeta)| \leq c|K(\zeta)| \quad \forall \zeta \in \partial\mathbb{D} \setminus \{\pm 1\},$$

then the discrete fractional Grönwall's inequality holds: if  $\alpha \in (0, 1)$  and  $p \in (1/\alpha, \infty)$ , and a sequence  $v^n \in X$ ,  $n = 0, 1, 2, \dots$ , with  $v^0 = 0$ , satisfies

$$(2.16) \quad \|(\bar{\partial}_\tau^\alpha v^n)_{n=1}^m\|_{\ell^p(X)} \leq \kappa \|(v^n)_{n=1}^m\|_{\ell^p(X)} + \sigma \quad \forall 1 \leq m \leq N$$

for some positive constants  $\kappa$  and  $\sigma$ , then there exists a  $\tau_0 > 0$  such that for any  $\tau < \tau_0$  there holds

$$(2.17) \quad \|(v^n)_{n=1}^N\|_{\ell^\infty(X)} + \|(\bar{\partial}_\tau^\alpha v^n)_{n=1}^N\|_{\ell^p(X)} \leq c\sigma,$$

where the constants  $c$  and  $\tau_0$  are independent of  $\sigma$ ,  $\tau$ ,  $N$ , and  $v^n$  but may depend on  $\alpha$ ,  $p$ ,  $\kappa$ ,  $X$ , and  $T$ .

*Proof.* First, we note that  $\bar{\partial}_\tau^\alpha v^n = \tau^{-\alpha} \sum_{j=0}^n K_j v^{n-j}$ ,  $n = 0, 1, 2, \dots$ , are the coefficients in the power series expansion

$$(2.18) \quad K(\zeta) \sum_{n=0}^{\infty} v^n \zeta^n = \sum_{n=0}^{\infty} (\bar{\partial}_\tau^\alpha v^n) \zeta^n;$$

it follows that

$$(2.19) \quad \sum_{n=0}^{\infty} v^n \zeta^n = \left(\frac{\tau}{1-\zeta}\right)^\alpha \left[\frac{1}{K(\zeta)} \left(\frac{1-\zeta}{\tau}\right)^\alpha\right] \sum_{n=0}^{\infty} (\bar{\partial}_\tau^\alpha v^n) \zeta^n = \left(\frac{\tau}{1-\zeta}\right)^\alpha \sum_{n=0}^{\infty} F^n \zeta^n,$$

where  $F^n$ ,  $n = 0, 1, \dots$ , are the coefficients in the expansion

$$\left[\frac{1}{K(\zeta)} \left(\frac{1-\zeta}{\tau}\right)^\alpha\right] \sum_{n=0}^{\infty} (\bar{\partial}_\tau^\alpha v^n) \zeta^n = \sum_{n=0}^{\infty} F^n \zeta^n.$$

By applying Lemma 2.5 to (2.19), we obtain

$$(2.20) \quad \|(v^n)_{n=0}^m\|_{\ell^\infty(X)} \leq c \|(F^n)_{n=0}^m\|_{\ell^p(X)} \quad \forall 1 \leq m \leq N.$$

Let  $m$  be fixed and define  $\tilde{E}^n = \bar{\partial}_\tau^\alpha v^n$  if  $n \leq m$  and  $\tilde{E}^n = 0$  if  $n > m$ . Let  $\tilde{F}^n$  be the coefficients of the power series

$$(2.21) \quad \sum_{n=0}^{\infty} \tilde{F}^n \zeta^n = \left[\frac{1}{K(\zeta)} \left(\frac{1-\zeta}{\tau}\right)^\alpha\right] \sum_{n=0}^{\infty} \tilde{E}^n \zeta^n;$$

then  $\tilde{F}^n = F^n$  for  $0 \leq n \leq m$ . Now the conditions in (2.15) imply

$$\left|\frac{1}{K(\zeta)} \left(\frac{1-\zeta}{\tau}\right)^\alpha\right| \leq c \quad \text{and} \quad \left|(1-\zeta)(1+\zeta) \frac{d}{d\zeta} \left[\frac{1}{K(\zeta)} \left(\frac{1-\zeta}{\tau}\right)^\alpha\right]\right| \leq c \quad \forall \zeta \in \partial\mathbb{D} \setminus \{\pm 1\}.$$

By choosing  $M(\zeta) = \frac{1}{K(\zeta)} \left(\frac{1-\zeta}{\tau}\right)^\alpha$  and applying Lemma 2.6 to (2.21), we obtain

$$\|(\tilde{F}^n)_{n=0}^\infty\|_{\ell^p(X)} \leq c \|(\tilde{E}^n)_{n=0}^\infty\|_{\ell^p(X)},$$

which further implies

$$\|(F^n)_{n=0}^m\|_{\ell^p(X)} = \|(\tilde{F}^n)_{n=0}^m\|_{\ell^p(X)} \leq c \|(\tilde{E}^n)_{n=0}^\infty\|_{\ell^p(X)} = c \|(\bar{\partial}_\tau^\alpha v^n)_{n=0}^m\|_{\ell^p(X)},$$

where the constant  $c$  is independent of  $m$ . The last inequality and (2.20) yield

$$\|(v^n)_{n=0}^m\|_{\ell^\infty(X)} \leq c\|(\bar{\partial}_\tau^\alpha v^n)_{n=0}^m\|_{\ell^p(X)}.$$

Substituting (2.16) into the last inequality gives

$$(2.22) \quad \begin{aligned} \|(v^n)_{n=1}^m\|_{\ell^\infty(X)} &\leq c\kappa\|(v^n)_{n=1}^m\|_{\ell^p(X)} + c\sigma \\ &\leq \epsilon\kappa\|(v^n)_{n=1}^m\|_{\ell^\infty(X)} + c_\epsilon\kappa\|(v^n)_{n=1}^m\|_{\ell^1(X)} + c\sigma \quad \forall 1 \leq m \leq N, \end{aligned}$$

where  $\epsilon > 0$  is arbitrary. By choosing  $\epsilon\kappa = 1/2$ , we obtain

$$\|(v^n)_{n=1}^m\|_{\ell^\infty(X)} \leq c_\kappa\|(v^n)_{n=1}^m\|_{\ell^1(X)} + c\sigma \quad \forall 1 \leq m \leq N.$$

That is,  $\|v^m\|_X \leq c_\kappa\tau \sum_{n=1}^m \|v^n\|_X + c\sigma$  for  $1 \leq m \leq N$ . Then the standard discrete Grönwall's inequality gives, for sufficiently small step size  $\tau$ ,

$$\max_{1 \leq n \leq N} \|v^n\|_X \leq e^{c_\kappa T} c\sigma.$$

This together with (2.16) and (2.22) yields (2.17). The proof of Theorem 2.7 is complete.  $\square$

By Theorem 2.7, the discrete fractional Grönwall's inequality can be proved for the L1 scheme and general BDF CQs.

**THEOREM 2.8** (discrete Grönwall's inequality for L1 scheme and BDF CQ). *Let  $X$  be a UMD space. For both the L1 scheme and CQ generated by the  $k$ th-order BDF, with  $1 \leq k \leq 6$ , the discrete fractional Grönwall's inequality holds: if  $\alpha \in (0, 1)$  and  $p \in (1/\alpha, \infty)$ , and a sequence  $v^n \in X$ ,  $n = 0, 1, 2, \dots$ , with  $v^0 = 0$ , satisfies*

$$\|(\bar{\partial}_\tau^\alpha v^n)_{n=1}^m\|_{\ell^p(X)} \leq \kappa\|(v^n)_{n=1}^m\|_{\ell^p(X)} + \sigma \quad \forall 1 \leq m \leq N$$

for some positive constants  $\kappa$  and  $\sigma$ , then there exists a  $\tau_0 > 0$  such that for any  $\tau < \tau_0$  there holds

$$\|(v^n)_{n=1}^N\|_{\ell^\infty(X)} + \|(\bar{\partial}_\tau^\alpha v^n)_{n=1}^N\|_{\ell^p(X)} \leq c\sigma,$$

where the constants  $c$  and  $\tau_0$  are independent of  $\sigma$ ,  $\tau$ ,  $N$ , and  $v^n$  but may depend on  $\alpha$ ,  $p$ ,  $\kappa$ ,  $X$ , and  $T$ .

*Proof.* By Theorem 2.7, it suffices to show that the generating functions  $K(\zeta)$  of the L1 scheme and CQ satisfy (2.15). We discuss them separately. First, for the L1 scheme,  $K(\zeta) = \frac{1}{\Gamma(2-\alpha)\tau^\alpha} \frac{(1-\zeta)^2}{\zeta} \text{Li}_{\alpha-1}(\zeta)$  converges for  $\zeta \in \partial\mathbb{D} \setminus \{1\}$  and has the following asymptotic expansion (cf. [11, Theorem 1] or [17, equation (4.6)]):

$$\tau^\alpha K(\zeta) = (1-\zeta)^\alpha + o((1-\zeta)^\alpha) \quad \text{as } \zeta \rightarrow 1.$$

If  $\zeta \in \partial\mathbb{D} \setminus \{1\}$  is sufficiently close to 1, then

$$\tau^\alpha |K(\zeta)| \geq \frac{1}{2}|1-\zeta|^\alpha.$$

Meanwhile, we recall the following series expansion (cf. [17, equation (4.5)]):

$$\frac{\text{Li}_{\alpha-1}(e^{-i\theta})}{\Gamma(2-\alpha)} = (2\pi)^{\alpha-2} \left( \cos\left(\frac{(2-\alpha)\pi}{2}\right) (A_\theta + B_\theta) - i \sin\left(\frac{(2-\alpha)\pi}{2}\right) (A_\theta - B_\theta) \right),$$

where  $A_\theta = \sum_{k=0}^{\infty} (k + \frac{\theta}{2\pi})^{\alpha-2}$  and  $B_\theta = \sum_{k=0}^{\infty} (k + 1 - \frac{\theta}{2\pi})^{\alpha-2}$ . Thus, if  $\zeta = e^{-i\theta}$  is away from 1, then  $\theta$  is away from 0 and  $2\pi$ , and thus  $A_\theta + B_\theta \geq c$ . This shows  $|\text{Li}_{\alpha-1}(e^{-i\theta})| > c$ . Since  $|1 - \zeta|^2 \geq c|1 - \zeta|^\alpha$  when  $\zeta = e^{-i\theta}$  is away from 1, it follows that

$$\tau^\alpha |K(\zeta)| = \frac{|\text{Li}_{\alpha-1}(\zeta)|}{\Gamma(2-\alpha)} |1 - \zeta|^2 \geq c|1 - \zeta|^2 \geq c|1 - \zeta|^\alpha.$$

Overall, the first inequality of (2.15) holds for the generating function  $K(\zeta)$  of the L1 scheme. The second inequality of (2.15) has been proved in [17, Lemma 4.3]. This shows the assertion for the L1 scheme.

Next we turn to the CQ. For the CQ generated by the  $k$ th-order BDF, the generating function  $K(\zeta)$  satisfies

$$\left(\frac{\tau}{1-\zeta}\right)^\alpha K(\zeta) = \left(\sum_{j=1}^k \frac{1}{j}(1-\zeta)^{j-1}\right)^\alpha.$$

Since the function  $\sum_{j=1}^k \frac{1}{j}(1-\zeta)^{j-1}$  has no root on the unit circle  $\partial\mathbb{D}$  for  $1 \leq k \leq 6$  (see [9, Proof of Lemma 2] or [12, pp. 246–247]), it follows that

$$\left|\left(\frac{\tau}{1-\zeta}\right)^\alpha K(\zeta)\right| = \left|\sum_{j=1}^k \frac{1}{j}(1-\zeta)^{j-1}\right|^\alpha \geq c.$$

This proves the first inequality of (2.15). Note that

$$\begin{aligned} (1+\zeta)(1-\zeta)K'(\zeta) &= -(1+\zeta)(1-\zeta) \frac{\alpha}{\tau^\alpha} \left(\sum_{j=1}^k \frac{1}{j}(1-\zeta)^j\right)^{\alpha-1} \sum_{j=1}^k (1-\zeta)^{j-1} \\ &= -\frac{\alpha}{\tau^\alpha} (1+\zeta)(1-\zeta)^\alpha \left(\sum_{j=1}^k \frac{1}{j}(1-\zeta)^{j-1}\right)^{\alpha-1} \sum_{j=0}^{k-1} (1-\zeta)^j, \end{aligned}$$

and so for any  $\zeta \in \partial\mathbb{D} \setminus \{\pm 1\}$ , there holds

$$\left|\frac{(1+\zeta)(1-\zeta)K'(\zeta)}{K(\zeta)}\right| = \left|\frac{\alpha(1+\zeta)\left(\sum_{j=1}^k \frac{1}{j}(1-\zeta)^{j-1}\right)^{\alpha-1} \sum_{j=0}^{k-1} (1-\zeta)^j}{\left(\sum_{j=1}^k \frac{1}{j}(1-\zeta)^{j-1}\right)^\alpha}\right| \leq c,$$

where the last inequality holds, since the denominator  $\left(\sum_{j=1}^k \frac{1}{j}(1-\zeta)^{j-1}\right)^\alpha$  has no root on  $\partial\mathbb{D}$ . This shows the second part of (2.15), completing the proof of the theorem.  $\square$

*Remark 2.9.* In Theorems 2.7 and 2.8, if we assume

$$\|(\bar{\partial}_\tau^\alpha v^n)_{n=1}^m\|_{\ell^p(X)} \leq \kappa \|(v^n)_{n=1}^{m-1}\|_{\ell^p(X)} + \sigma \quad \forall 1 \leq m \leq N,$$

i.e., the index on the right-hand side is slightly changed, then we have

$$\|(v^n)_{n=1}^N\|_{\ell^\infty(X)} + \|(\bar{\partial}_\tau^\alpha v^n)_{n=1}^N\|_{\ell^p(X)} \leq c\sigma$$

without any restriction on the step size  $\tau$ .

**3. Regularity of the solution.** Now we discuss the existence, uniqueness, and regularity for the solutions to (1.1) and (1.4). These results are needed in the numerical analysis in section 4. The main result of this section is the following theorem.

**THEOREM 3.1.** *Let  $u_0 \in H_0^1(\Omega) \cap H^2(\Omega)$ , and let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be Lipschitz continuous. Then problem (1.1) has a unique solution  $u$  such that*

$$(3.1) \quad u \in C^\alpha([0, T]; L^2(\Omega)) \cap C([0, T]; H_0^1(\Omega) \cap H^2(\Omega)), \quad \partial_t^\alpha u \in C([0, T]; L^2(\Omega)),$$

$$(3.2) \quad \partial_t u(t) \in L^2(\Omega) \quad \text{and} \quad \|\partial_t u(t)\|_{L^2(\Omega)} \leq ct^{\alpha-1} \quad \text{for } t \in (0, T].$$

Similarly, problem (1.4) has a unique solution  $u_h$  such that

$$(3.3) \quad \|u_h\|_{C^\alpha([0, T]; L^2(\Omega))} + \|\Delta_h u_h\|_{C([0, T]; L^2(\Omega))} + \|\partial_t^\alpha u_h\|_{C([0, T]; L^2(\Omega))} \leq c,$$

$$(3.4) \quad \|\partial_t u_h(t)\|_{L^2(\Omega)} \leq ct^{\alpha-1} \quad \text{for } t \in (0, T].$$

The constant  $c$  above is independent of the mesh size  $h$  but may depend on  $T$ .

*Remark 3.2.* For smooth initial data and right-hand side, in the absence of extra compatibility conditions, the regularity results (3.1)–(3.2) and the  $h$ -independent estimates (3.3)–(3.4) are sharp with respect to the Hölder continuity in time. The regularity (3.1) was shown in [36] for linear subdiffusion equations and in [29] for a semilinear problem with Neumann boundary conditions under certain compatibility conditions. However, we are not aware of any existing results such as (3.2) and (3.3)–(3.4) for semilinear problems without compatibility conditions, which are important for the numerical analysis in section 4.

*Remark 3.3.* If  $f$  is smooth but not Lipschitz continuous, and problems (1.1) and (1.4) have unique bounded solutions, respectively, then  $f(u)$ ,  $f'(u)$ ,  $f(u_h)$ , and  $f'(u_h)$  are still bounded. In this case, the estimates (3.1)–(3.2) and (3.3)–(3.4) are still valid, which can be seen from the proof of Theorem 3.1.

We begin with some preliminary results. Let  $L_h^2(\Omega)$  be the vector space  $S_h$  equipped with the norm of  $L^2(\Omega)$  and let  $H_h^2(\Omega)$  be the vector space  $S_h$  equipped with the norm

$$\|v_h\|_{H_h^2(\Omega)} := \|v_h\|_{L^2(\Omega)} + \|\Delta_h v_h\|_{L^2(\Omega)} \quad \forall v_h \in S_h.$$

To analyze  $u(t)$  and  $u_h(t)$  in a unified way, we consider the following abstract problem:

$$(3.5) \quad \begin{cases} \partial_t^\alpha u(t) - Au(t) = Pf(u(t)) & \text{for } t \in (0, T], \\ u(0) = u_0, \end{cases}$$

where the notation  $(X, D, A, u, P, u_0)$  denotes either  $(L^2(\Omega), H_0^1(\Omega) \cap H^2(\Omega), \Delta, u, I, u_0)$  or  $(L_h^2(\Omega), H_h^2(\Omega), \Delta_h, u_h, P_h, R_h u_0)$ , with  $I$  denoting the identity operator. On a bounded convex polygonal domain  $\Omega$ , the norm of  $D$  is equivalent to the graph norm, i.e.,

$$(3.6) \quad \|v\|_D \sim \|v\|_X + \|Av\|_X \quad \forall v \in D.$$

Let  $\|\cdot\|_{X \rightarrow X}$  be the operator norm on the space  $X$ . Then the operator  $A$  satisfies the following resolvent estimate [4, Example 3.7.5 and Theorem 3.7.11]:

$$\|(z - A)^{-1}\|_{X \rightarrow X} \leq c_\phi |z|^{-1} \quad \forall z \in \Sigma_\phi, \quad \forall \phi \in (0, \pi),$$

where for  $\phi \in (0, \pi)$ ,  $\Sigma_\phi := \{z \in \mathbb{C} \setminus \{0\} : |\arg(z)| < \phi\}$ . This further implies

$$(3.7) \quad \begin{aligned} \|(z^\alpha - A)^{-1}\|_{X \rightarrow X} &\leq c_{\phi, \alpha} |z|^{-\alpha} \quad \forall z \in \Sigma_\phi \quad \forall \phi \in (0, \pi), \\ \|A(z^\alpha - A)^{-1}\|_{X \rightarrow X} &\leq c_{\phi, \alpha} \quad \forall z \in \Sigma_\phi \quad \forall \phi \in (0, \pi). \end{aligned}$$

Let  $g(t) = Pf(u(t))$ , and  $w := u - u_0$ . Then  $w$  satisfies the following equation:

$$(3.8) \quad \partial_t^\alpha w(t) - Aw(t) = Au_0 + g(t)$$

with  $w(0) = 0$ . By means of the Laplace transform, denoted by  $\widehat{\cdot}$ , we obtain

$$z^\alpha \widehat{w}(z) - A\widehat{w}(z) = z^{-1}Au_0 + \widehat{g}(z),$$

which together with (3.7) implies  $\widehat{w}(z) = (z^\alpha - A)^{-1}(z^{-1}Au_0 + \widehat{g}(z))$ . By the inverse Laplace transform and convolution rule, the solution  $w(t)$  to (3.8) is given by

$$(3.9) \quad w(t) = F(t)Au_0 + \int_0^t E(t-s)g(s)ds,$$

where the operators  $F(t) : X \rightarrow X$  and  $E(t) : X \rightarrow X$  are defined by

$$(3.10) \quad F(t) := \frac{1}{2\pi i} \int_{\Gamma_{\theta, \delta}} e^{zt} z^{-1} (z^\alpha - A)^{-1} dz \quad \text{and} \quad E(t) := \frac{1}{2\pi i} \int_{\Gamma_{\theta, \delta}} e^{zt} (z^\alpha - A)^{-1} dz,$$

respectively. Clearly, we have  $E(t) = F'(t)$ . The contour  $\Gamma_{\theta, \delta}$  is defined by

$$(3.11) \quad \Gamma_{\theta, \delta} = \{z \in \mathbb{C} : |z| = \delta, |\arg z| \leq \theta\} \cup \{z \in \mathbb{C} : z = \rho e^{\pm i\theta}, \rho \geq \delta\},$$

oriented with an increasing imaginary part, where  $\theta \in (\pi/2, \pi)$  is fixed. In view of (3.9),  $u$  is the solution of problem (3.5) if and only if it is the solution of

$$(3.12) \quad u(t) - u_0 = F(t)Au_0 + \int_0^t E(t-s)Pf(u(s))ds.$$

The next lemma summarizes the mapping properties of the operators  $F$  and  $E$ . These are partially known [36, section 2], [30]. We only sketch the proof for completeness.

LEMMA 3.4. *For the operators  $F$  and  $E$ , the following properties hold:*

- (i)  $t^{-\alpha} \|F(t)\|_{X \rightarrow X} + t^{1-\alpha} \|F'(t)\|_{X \rightarrow X} + \|AF(t)\|_{X \rightarrow X} \leq c \quad \forall t \in (0, T]$ .
- (ii)  $F(t) : X \rightarrow D$  is continuous with respect to  $t \in [0, T]$ , and  $AF(0) = 0$ .
- (iii)  $t^{1-\alpha} \|E(t)\|_{X \rightarrow X} + t^{2-\alpha} \|E'(t)\|_{X \rightarrow X} + t \|AE(t)\|_{X \rightarrow X} \leq c \quad \forall t \in (0, T]$ .
- (iv)  $E(t) : X \rightarrow D$  is continuous with respect to  $t \in (0, T]$ .

*Proof.* First, consider (ii) in the case  $X = L^2(\Omega)$ ,  $D = H_0^1(\Omega) \cap H^2(\Omega)$ , and  $A = \Delta$ . By setting  $f(u(t)) \equiv 0$  and  $A = \Delta$  in (3.12), [36, Theorem 2.1] implies that  $\Delta F(t) = F(t)\Delta : L^2(\Omega) \rightarrow L^2(\Omega)$  is continuous with respect to  $t \in [0, T]$ . Thus,  $F(t) : L^2(\Omega) \rightarrow H_0^1(\Omega) \cap H^2(\Omega)$  is continuous with respect to  $t \in [0, T]$ . Then taking  $t \rightarrow 0$  in (3.12) yields  $\Delta F(0) = 0$ . This proves (ii) in the case  $X = L^2(\Omega)$ ,  $D = H_0^1(\Omega) \cap H^2(\Omega)$ , and  $A = \Delta$ . The proof for the case  $X = L_h^2(\Omega)$ ,  $D = H_h^2(\Omega)$ , and  $A = \Delta_h$  is similar.

For any integers  $k \geq 0$  and  $m = 0, 1$ , by choosing  $\delta = t^{-1}$  in the contour  $\Gamma_{\theta, \delta}$  and using the identity  $A(z^\alpha - A)^{-1} = -I + z^\alpha(z^\alpha - A)^{-1}$ , the resolvent estimate (3.7),

and change of variables  $z = s \cos \varphi + is \sin \varphi$ , we have (with  $|dz|$  being the arc length element of  $\Gamma_{\theta,\delta}$ )

$$\begin{aligned} \left\| A^m \frac{d^k}{dt^k} F(t) \right\|_{X \rightarrow X} &= \left\| \frac{1}{2\pi i} \int_{\Gamma_{\theta,\delta}} e^{zt} z^{k-1} A^m (z^\alpha - A)^{-1} dz \right\|_{X \rightarrow X} \\ &\leq c \int_{\Gamma_{\theta,\delta}} e^{\operatorname{Re}(z)t} |z|^{k-1+(m-1)\alpha} |dz| \\ &\leq c |\cos \theta| \int_{\delta}^{\infty} e^{st \cos \theta} s^{k-1+(m-1)\alpha} ds + c \int_{-\theta}^{\theta} e^{\cos \varphi} \delta^{k+(m-1)\alpha} d\varphi \\ &\leq ct^{-(m-1)\alpha-k}. \end{aligned}$$

Since  $E(t) = F'(t)$ , the last inequality yields (i) and (iii). The continuity of  $F(t) : X \rightarrow D$  and  $E(t) : X \rightarrow D$  for  $t \in (0, T]$  follows from the equivalent norm in (3.6), showing (iv).  $\square$

Now we are ready to present the proof of Theorem 3.1.

*Proof of Theorem 3.1.* The proof is divided into four steps.

*Step 1: Existence and uniqueness.* We denote by  $C([0, T]; X)_\lambda$  the function space  $C([0, T]; X)$  equipped with the following weighted norm:

$$\|v\|_\lambda := \max_{0 \leq t \leq T} \|e^{-\lambda t} v(t)\|_X \quad \forall v \in C([0, T]; X),$$

which is equivalent to the standard norm of  $C([0, T]; X)$  for any fixed parameter  $\lambda > 0$ . Then we define a nonlinear map  $M : C([0, T]; X)_\lambda \rightarrow C([0, T]; X)_\lambda$  by

$$Mv(t) = u_0 + F(t)Au_0 + \int_0^t E(t-s)Pf(v(s))ds.$$

For any  $\lambda > 0$ ,  $u \in C([0, T]; X)$  is a solution of (3.12) if and only if  $u$  is a fixed point of the map  $M : C([0, T]; X)_\lambda \rightarrow C([0, T]; X)_\lambda$ . It remains to prove that for some  $\lambda > 0$ , the map  $M : C([0, T]; X)_\lambda \rightarrow C([0, T]; X)_\lambda$  has a unique fixed point. In fact, the definition of  $M$  and Lemma 3.4(iii) immediately yield

$$\begin{aligned} (3.13) \quad &\|e^{-\lambda t}(Mv_1(t) - Mv_2(t))\|_X \\ &= \left\| e^{-\lambda t} \int_0^t E(t-s)(Pf(v_1(s)) - Pf(v_2(s)))ds \right\|_X \\ &\leq ce^{-\lambda t} \int_0^t (t-s)^{\alpha-1} \|v_1(s) - v_2(s)\|_X ds \\ &\leq c \int_0^t (t-s)^{\alpha-1} e^{-\lambda(t-s)} \max_{s \in [0, T]} \|e^{-\lambda s}(v_1(s) - v_2(s))\|_X ds \\ &= c\lambda^{-\alpha} \left( \int_0^1 (1-\theta)^{\alpha-1} (\lambda t)^\alpha e^{-\lambda t(1-\theta)} d\theta \right) \|v_1 - v_2\|_\lambda \quad (\text{change of variable } s = t\theta) \\ &\leq c \sup_{\substack{\lambda > 0, T \geq t > 0 \\ \theta \in [0, 1]}} \left( [\lambda t(1-\theta)]^{\alpha/2} e^{-\lambda t(1-\theta)} \right) (t/\lambda)^{\alpha/2} \left( \int_0^1 (1-\theta)^{\alpha/2-1} d\theta \right) \|v_1 - v_2\|_\lambda \\ &\leq c(T/\lambda)^{\alpha/2} \|v_1 - v_2\|_\lambda \quad \forall v_1, v_2 \in C([0, T]; X)_\lambda. \end{aligned}$$

By choosing a sufficiently large  $\lambda$ , the last inequality implies

$$\|e^{-\lambda t}(Mv_1(t) - Mv_2(t))\|_X \leq \frac{1}{2}\|v_1 - v_2\|_\lambda \quad \forall v_1, v_2 \in C([0, T]; X)_\lambda.$$

Hence, the map  $M$  is contractive on the space  $C([0, T]; X)_\lambda$ . The Banach fixed point theorem implies that  $M$  has a unique fixed point, which is also the unique solution of (3.12).

*Step 2:  $C^\alpha([0, T]; X)$  regularity.* Consider the difference quotient for  $h > 0$ ,

(3.14)

$$\begin{aligned} \frac{u(t+h) - u(t)}{h^\alpha} &= \frac{F(t+h) - F(t)}{h^\alpha} Au_0 + \frac{1}{h^\alpha} \int_t^{t+h} E(s) Pf(u(t-s)) ds \\ &\quad + \int_0^t E(s) \frac{Pf(u(t+h-s)) - Pf(u(t-s))}{h^\alpha} ds =: \sum_{i=1}^3 \mathcal{I}_i(t, h). \end{aligned}$$

A simple consequence of Lemma 3.4(i) is that  $h^{-\alpha}\|F(t+h) - F(t)\|_{X \rightarrow X} \leq c$ , which implies  $\|\mathcal{I}_1(t, h)\|_X \leq c$ . By appealing to Lemma 3.4(iii), we have

$$\begin{aligned} \|\mathcal{I}_2(t, h)\|_X &= \left\| \frac{1}{h^\alpha} \int_t^{t+h} E(s) Pf(u(t-s)) ds \right\|_X \\ &\leq c \frac{1}{h^\alpha} \int_t^{t+h} s^{\alpha-1} ds = \frac{c}{\alpha} \frac{(t+h)^\alpha - t^\alpha}{h^\alpha} \leq c. \end{aligned}$$

By the Lipschitz continuity of  $f$ , we have

$$\begin{aligned} e^{-\lambda t} \|\mathcal{I}_3(t, h)\|_X &= \left\| e^{-\lambda t} \int_0^t E(t-s) \frac{Pf(u(s+h)) - Pf(u(s))}{h^\alpha} ds \right\|_X \\ &\leq c_1 \int_0^t e^{-\lambda(t-s)} (t-s)^{\alpha-1} e^{-\lambda s} \left\| \frac{u(s+h) - u(s)}{h^\alpha} \right\|_X ds. \end{aligned}$$

By substituting the estimates of  $\mathcal{I}_i(t, h)$ ,  $i = 1, 2, 3$ , into (3.14) and denoting  $W_h(t) = e^{-\lambda t} h^{-\alpha} \|u(t+h) - u(t)\|_X$ , we obtain

$$W_h(t) \leq c + c_1 \int_0^t e^{-\lambda(t-s)} (t-s)^{\alpha-1} W_h(s) ds \leq c + c_1 (T/\lambda)^{\frac{\alpha}{2}} \max_{s \in [0, T]} W_h(s),$$

where the last inequality can be derived in the same way as (3.13). By choosing a sufficiently large  $\lambda$  and taking maximum of the left-hand side with respect to  $t \in [0, T]$ , it implies  $\max_{t \in [0, T]} W_h(t) \leq c$ , which further yields

$$h^{-\alpha} \|u(t+h) - u(t)\|_X \leq ce^{\lambda t} \leq c,$$

where the constant  $c$  is independent of  $h$ . Thus, we have proved  $\|u\|_{C^\alpha([0, T]; X)} \leq c$ .

*Step 3:  $C([0, T]; D)$  regularity.* By applying the operator  $A$  to both sides of (3.12) and using the identity  $AF(t) = \int_0^t AE(t-s) ds$  (cf. Lemma 3.4), we obtain

(3.15)

$$\begin{aligned} Au(t) - Au_0 &= AF(t)Au_0 + \int_0^t AE(t-s) Pf(u(s)) ds \\ &= AF(t)(Au_0 + Pf(u(t))) + \int_0^t AE(t-s)(Pf(u(s)) - Pf(u(t))) ds \\ &= \mathcal{I}_4(t) + \mathcal{I}_5(t). \end{aligned}$$

By Lemma 3.4(iii) and the  $C^\alpha([0, T]; X)$  regularity from Step 2, we have

$$\begin{aligned} \|\mathcal{I}_5(t)\|_X &= \left\| \int_0^t AE(t-s)(Pf(u(s)) - Pf(u(t)))ds \right\|_X \\ &\leq \int_0^t \frac{c\|u(s) - u(t)\|_X}{t-s} ds \leq \int_0^t \frac{c|t-s|^\alpha}{t-s} ds \leq ct^\alpha \quad \forall t \in (0, T]. \end{aligned}$$

Lemma 3.4(iv) implies that  $\mathcal{I}_5(t)$  is continuous for  $t \in (0, T]$ , and the last inequality implies that  $\mathcal{I}_5(t)$  is also continuous at  $t = 0$ . Hence  $\mathcal{I}_5 \in C([0, T]; X)$ . Moreover, Lemma 3.4(ii) gives  $\mathcal{I}_4 \in C([0, T]; X)$  and

$$\|\mathcal{I}_4(t)\|_X \leq c\|Au_0 + Pf(u(t))\|_X \leq c.$$

Substituting the estimates of  $\mathcal{I}_4(t)$  and  $\mathcal{I}_5(t)$  into (3.15) yields  $\|Au\|_{C([0, T]; X)} \leq c$ , which further implies  $\|u\|_{C([0, T]; D)} \leq c$ . The regularity result  $u \in C([0, T]; D)$  together with (3.5) yields  $\partial_t^\alpha u = Au + Pf(u) \in C([0, T]; X)$ .

*Step 4: Estimate of  $\|u'(t)\|_X$ .* By differentiating (3.12) with respect to  $t$ , we obtain

$$\begin{aligned} u'(t) &= F'(t)Au_0 + E(t)Pf(u_0) + \int_0^t E(s)Pf'(u(t-s))u'(t-s)ds \\ &= E(t)(Au_0 + Pf(u_0)) + \int_0^t E(t-s)Pf'(u(s))u'(s)ds. \end{aligned}$$

By multiplying this equation by  $t^{1-\alpha}$ , we get

$$t^{1-\alpha}u'(t) = t^{1-\alpha}E(t)(Au_0 + Pf(u_0)) + \int_0^t t^{1-\alpha}s^{\alpha-1}E(t-s)Pf'(u(s))s^{1-\alpha}u'(s)ds,$$

which together with the  $L^\infty$  stability of  $P_h$  [38, Lemma 6.1] directly implies that

$$\begin{aligned} e^{-\lambda t}t^{1-\alpha}\|u'(t)\|_X &\leq e^{-\lambda t}t^{1-\alpha}\|E(t)\|_{X \rightarrow X}\|Au_0 + Pf(u_0)\|_X \\ &\quad + \int_0^t e^{-\lambda(t-s)}t^{1-\alpha}s^{\alpha-1}(t-s)^{\alpha-1}\|Pf'(u(s))\|_{L^\infty(\Omega)}e^{-\lambda s}s^{1-\alpha}\|u'(s)\|_X ds \\ &\leq ce^{-\lambda t}\|Au_0 + Pf(u_0)\|_X + c(T/\lambda)^{\frac{\alpha}{2}} \max_{s \in [0, T]} e^{-\lambda s}s^{1-\alpha}\|u'(s)\|_X, \end{aligned}$$

where the last line follows similarly as (3.13). By choosing a sufficiently large  $\lambda$  and taking maximum of the left-hand side with respect to  $t \in [0, T]$ , it implies  $\max_{t \in [0, T]} \|e^{-\lambda t}t^{1-\alpha}u'(t)\|_X \leq c$ , which further yields (3.2). The proof of Theorem 3.1 is complete.  $\square$

**4. Error estimates.** Now, we derive error estimates for the numerical solutions of problem (1.1) using the discrete Grönwall's inequality from section 2 and discrete maximal  $\ell^p$ -regularity from [17]. To illustrate the general framework for the numerical analysis of nonlinear time-fractional diffusion equations, we focus on the L1 scheme and backward Euler CQ. Other time-stepping schemes can be analyzed similarly. The convergence rates we show below are sharp (up to a logarithmic factor) with respect to the solution regularity in Theorem 3.1 and are also confirmed by the numerical experiments in section 6.

**4.1. Preliminaries on the linear problem.** First we recall some error estimates for the following linear subdiffusion equation:

$$(4.1) \quad \partial_t^\alpha v(t) - \Delta v(t) = g(t) \quad \forall t \in (0, T],$$

where  $g$  is a given function. The semidiscrete FEM for (4.1) seeks  $v_h(t) \in S_h$  such that

$$(4.2) \quad \partial_t^\alpha v_h(t) - \Delta_h v_h(t) = P_h g(t) \quad \forall t \in (0, T]$$

with  $v_h(0) = R_h v(0)$ , and the fully discrete scheme seeks  $v_h^n \in S_h$ ,  $n = 1, \dots, N$ , such that

$$(4.3) \quad \bar{\partial}_\tau^\alpha (v_h^n - v_h^0) - \Delta_h v_h^n = P_h g(t_n)$$

with  $v_h^0 = v_h(0)$ , where  $\bar{\partial}_\tau^\alpha v_h^n$  denotes either the backward Euler CQ or the L1 scheme.

The semidiscrete solution  $v_h$  satisfies the following error estimate [14, 13, 16].

LEMMA 4.1 (semidiscrete solution of linear problems). *For the semidiscrete solution  $v_h$  to problem (4.2), there holds with  $\ell_h = \log(2 + 1/h)$*

$$\max_{t \in [0, T]} \|v_h(t) - v(t)\|_{L^2(\Omega)} \leq ch^2 \|v(0)\|_{H^2(\Omega)} + ch^2 \ell_h^2 \|g\|_{L^\infty(0, T; L^2(\Omega))}.$$

The solution  $v_h^n$  of the fully discrete scheme (4.3) satisfies the following error estimate. For the backward Euler CQ, it was proved in [16, Theorems 3.5 and 3.6], while the proof for the L1 scheme will be given in section 5.

LEMMA 4.2 (fully discrete solutions of linear problems). *For the fully discrete solutions  $v_h^n$  to problem (4.3) with the L1 scheme or backward Euler CQ, there holds*

$$\begin{aligned} \|v_h(t_n) - v_h^n\|_{L^2(\Omega)} &\leq c\tau t_n^{\alpha-1} (\|\Delta v(0)\|_{L^2(\Omega)} + \|g(0)\|_{L^2(\Omega)}) \\ &\quad + c\tau \int_0^{t_n} (t_n - s)^{\alpha-1} \|g'(s)\|_{L^2(\Omega)} ds. \end{aligned}$$

*Remark 4.3.* If  $1 \leq d \leq 3$  and  $v(0) \in H_0^1(\Omega) \cap H^2(\Omega)$ , then the error estimates in Lemmas 4.1 and 4.2 are still valid if  $v_h(0)$  is the Lagrange interpolation of  $v(0)$ , due to the smoothing property of the solution operator [14, Lemma 3.1]. Consequently, all the results in section 4.2 remain valid in this case.

Lemmas 4.1 and 4.2 will be used below in the analysis of the nonlinear problem.

**4.2. Error estimates for the nonlinear problem.** Now we can present error estimates for problem (1.1). Like in the linear case, we discuss the spatial error and temporal error separately. First, we derive the spatial discretization error.

THEOREM 4.4. *Let  $u_0 \in H_0^1(\Omega) \cap H^2(\Omega)$ , and let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be Lipschitz continuous. Then the semidiscrete problem (1.4) has a unique solution  $u_h \in C([0, T]; L_h^2(\Omega))$ , which satisfies*

$$(4.4) \quad \max_{0 \leq t \leq T} \|u(t) - u_h(t)\|_{L^2(\Omega)} \leq c\ell_h^2 h^2.$$

*Proof.* By Theorem 3.1, the existence and uniqueness of the solution  $u_h$  hold. It remains to establish the estimate (4.4). To this end, we define  $v_h(t)$  as the solution of

$$\partial_t^\alpha v_h(t) - \Delta_h v_h(t) = P_h f(u(t)) \quad \text{with} \quad v_h(0) = u_h(0) = R_h u_0.$$

This together with Lemma 4.1 yields the following estimate for  $t \geq 0$ :

$$(4.5) \quad \|(u - v_h)(t)\|_{L^2(\Omega)} \leq ch^2 \|u(0)\|_{H^2(\Omega)} + ch^2 \ell_h^2 \|f(u)\|_{L^\infty(0,T;L^2(\Omega))} \leq ch^2 \ell_h^2.$$

Meanwhile, we note that  $\rho_h := v_h - u_h$  satisfies the following equation:

$$\partial_t^\alpha \rho_h(t) - \Delta \rho_h(t) = P_h f(u(t)) - P_h f(u_h(t)) \quad \text{with} \quad \rho_h(0) = 0.$$

Then, by the Lipschitz continuity of  $f$  and the maximal  $L^p$ -regularity of fractional evolution equations [5, Corollary 1], we obtain the following estimate for any  $p \in (1, \infty)$ :

$$\begin{aligned} \|\partial_t^\alpha \rho_h\|_{L^p(0,T;L^2(\Omega))} &\leq c \|P_h f(u) - P_h f(u_h)\|_{L^p(0,T;L^2(\Omega))} \\ &\leq c \|u - u_h\|_{L^p(0,T;L^2(\Omega))} \\ &\leq c \|u - v_h\|_{L^p(0,T;L^2(\Omega))} + c \|\rho_h\|_{L^p(0,T;L^2(\Omega))} \\ &\leq ch^2 \ell_h^2 + c \|\rho_h\|_{L^p(0,T;L^2(\Omega))}. \end{aligned}$$

Then by the fractional Grönwall's inequality in Theorem 2.1, we have

$$\max_{t \in [0,T]} \|\rho_h(t)\|_{L^2(\Omega)} \leq ch^2 \ell_h^2.$$

This and (4.5) directly imply the desired result. □

Next we give the temporal discretization error.

**THEOREM 4.5.** *Let  $u_0 \in H_0^1(\Omega) \cap H^2(\Omega)$ , and let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be Lipschitz continuous. Then the fully discrete scheme (1.3), with either the L1 scheme or backward Euler CQ for time discretization, has a unique solution  $u_h^n \in S_h$ ,  $n = 1, \dots, N$ , and the solutions satisfy*

$$(4.6) \quad \max_{1 \leq n \leq N} \|u_h(t_n) - u_h^n\|_{L^2(\Omega)} \leq c\tau^\alpha.$$

*Proof.* For given  $u_h^0, \dots, u_h^{n-1}$ , (1.3) is essentially a linear system with a symmetric positive definite matrix, and thus it has a unique solution  $u_h^n \in S_h$ . It suffices to establish the estimate (4.6). Like before, we decompose the fully discrete solution  $u_h^n$  into two parts,  $u_h^n = v_h^n + \rho_h^n$ , where  $v_h^n$  and  $\rho_h^n$ , respectively, satisfy

$$(4.7) \quad \bar{\partial}_\tau^\alpha (v_h^n - v_h^0) - \Delta_h v_h^n = P_h f(u_h(t_n)),$$

$$(4.8) \quad \bar{\partial}_\tau^\alpha \rho_h^n - \Delta_h \rho_h^n = P_h f(u_h^{n-1}) - P_h f(u_h(t_n)),$$

with  $v_h^0 = u_h(0) = R_h u_0$  and  $\rho_h^0 = 0$ . Equation (4.7) can be viewed as the time discretization of (1.4), with the right-hand side being a given function. Hence, by Lemma 4.2 and using  $\|\partial_s u_h(s)\|_{L^2(\Omega)} \leq cs^{\alpha-1}$  (cf. Theorem 3.1) and Rademacher's theorem, we have

$$\begin{aligned} \|u_h(t_n) - v_h^n\|_{L^2(\Omega)} &\leq ct_n^{\alpha-1} \tau (\|\Delta_h u_h(0)\|_{L^2(\Omega)} + \|f(u_h(0))\|_{L^2(\Omega)}) \\ &\quad + c\tau \int_0^{t_n} (t_n - s)^{\alpha-1} \|f'(u_h(s)) \partial_s u_h(s)\|_{L^2(\Omega)} ds \\ (4.9) \quad &\leq ct_n^{\alpha-1} \tau + c\tau \int_0^{t_n} (t_n - s)^{\alpha-1} s^{\alpha-1} ds \\ &\leq ct_n^{\alpha-1} \tau + ct_n^{2\alpha-1} \tau \leq c\tau^\alpha. \end{aligned}$$

It remains to estimate  $\rho_h^n$ . By applying the discrete maximal  $\ell^p$ -regularity to (4.8) (choosing  $X = L_h^2(\Omega)$  in [17, Theorems 3.1 and 4.1]), we obtain that for all  $1 < p < \infty$ ,

$$\begin{aligned} \|(\bar{\partial}_\tau^\alpha \rho_h^n)_{n=1}^m\|_{\ell^p(L^2(\Omega))} &\leq c\|(f(u_h^{n-1}) - f(u_h(t_n)))_{n=1}^m\|_{\ell^p(L^2(\Omega))} \\ &\leq c\|(f(u_h^{n-1}) - f(u_h(t_{n-1})))_{n=1}^m\|_{\ell^p(L^2(\Omega))} \\ &\quad + c\|(f(u_h(t_{n-1})) - f(u_h(t_n)))_{n=1}^m\|_{\ell^p(L^2(\Omega))}. \end{aligned}$$

By the Lipschitz continuity of  $f$  and the triangle inequality, we arrive at

$$\begin{aligned} &\|(f(u_h^{n-1}) - f(u_h(t_{n-1})))_{n=1}^m\|_{\ell^p(L^2(\Omega))} \\ &\leq c\|(u_h(t_{n-1}) - u_h^{n-1})_{n=1}^m\|_{\ell^p(L^2(\Omega))} \\ &\leq c\|(u_h(t_{n-1}) - v_h^{n-1})_{n=1}^m\|_{\ell^p(L^2(\Omega))} + c\|(\rho_h^{n-1})_{n=1}^m\|_{\ell^p(L^2(\Omega))} \\ &\leq c\tau^\alpha + c\|(\rho_h^n)_{n=1}^{m-1}\|_{\ell^p(L^2(\Omega))}, \end{aligned}$$

where the last inequality follows from (4.9). Similarly, by the Lipschitz continuity of  $f$  and the a priori estimate  $\|u_h\|_{C^\alpha([0,T];L^2(\Omega))} \leq c$  (cf. Theorem 3.1), we deduce

$$\begin{aligned} \|(\|f(u_h(t_{n-1})) - f(u_h(t_n))\|_{L^2(\Omega)})_{n=1}^m\|_{\ell^p} &\leq c\|(\|u_h(t_{n-1}) - u_h(t_n)\|_{L^2(\Omega)})_{n=1}^m\|_{\ell^p} \\ &\leq c\|(c\tau^\alpha)_{n=1}^m\|_{\ell^p}. \end{aligned}$$

Combining the preceding three estimates yields

$$\|(\bar{\partial}_\tau^\alpha \rho_h^n)_{n=1}^m\|_{\ell^p(L^2(\Omega))} \leq c\|(\rho_h^n)_{n=1}^{m-1}\|_{\ell^p(L^2(\Omega))} + c\tau^\alpha.$$

By choosing  $p > 1/\alpha$  and applying the discrete Grönwall's inequality (with  $X = L^2(\Omega)$  in Theorem 2.8), we obtain

$$(4.10) \quad \max_{1 \leq n \leq N} \|\rho_h^n\|_{L^2(\Omega)} \leq c\tau^\alpha.$$

In view of the decomposition  $u_h(t_n) - u_h^n = (u_h(t_n) - v_h^n) - \rho_h^n$ , the two estimates (4.9) and (4.10) imply (4.6), completing the proof of the theorem.  $\square$

*Remark 4.6.* If the nonlinear source  $f$  is not Lipschitz continuous but problem (1.1) has a unique bounded solution  $u$ , then Theorems 4.4 and 4.5 are still valid by proving the boundedness of the semidiscrete solution  $u_h$  and the fully discrete solution  $u_h^n$ . For simplicity, we have assumed  $f$  to be Lipschitz continuous in order to avoid these technicalities.

**5. Proof of Lemma 4.2 for the L1 scheme.** The L1 scheme was analyzed in [15] only for the homogeneous problem. Below we give a proof for the general case.

First, we assume that  $g$  is time-independent, i.e.,  $g(t) \equiv g(0)$ . Then using the Laplace transform, one can derive the error representation (cf. [15, equations (2.7) and (2.9)])

$$\begin{aligned} v_h(t_n) - v_h^n &= \frac{1}{2\pi i} \int_{\Gamma_{\theta,\delta}} e^{zt_n} z^{-1} (z^\alpha - \Delta_h)^{-1} (\Delta_h v_h(0) + P_h g(0)) dz \\ &\quad - \frac{1}{2\pi i} \int_{\Gamma_{\theta,\delta}^\tau} e^{zt_n} \mu(e^{-z\tau})^{-1} (\beta_\tau(e^{-z\tau}) - \Delta_h)^{-1} (\Delta_h v_h(0) + P_h g(0)) dz, \end{aligned}$$

where the contour  $\Gamma_{\theta,\delta}$  is defined in (3.11),  $\Gamma_{\theta,\delta}^\tau = \{z \in \Gamma_{\theta,\delta} : |\text{Im}(z)| \leq 1/\tau\}$ , and

$$\mu(z) = \frac{1 - e^{-z\tau}}{\tau e^{-z\tau}} \quad \text{and} \quad \beta_\tau(e^{-z\tau}) = \frac{(1 - e^{-z\tau})^2}{e^{-z\tau} \tau^\alpha \Gamma(2 - \alpha)} \text{Li}_{\alpha-1}(e^{-z\tau}),$$

which satisfy the following estimates (cf. [15, section 3]):

$$(5.1) \quad c_0|z| \leq |\mu(e^{-z\tau})| \leq c_1|z| \quad \text{and} \quad |\mu(e^{-z\tau}) - z| \leq c\tau|z|^2 \quad \forall z \in \Gamma_{\theta,\delta}^\tau,$$

$$(5.2) \quad |\beta_\tau(e^{-z\tau})| \geq c|z|\tau^{1-\alpha} \quad \text{and} \quad |\beta_\tau(e^{-z\tau}) - z^\alpha| \leq c|z|^2\tau^{2-\alpha} \quad \forall z \in \Gamma_{\theta,\delta}^\tau.$$

By using (5.1)–(5.2), direct calculations yield

$$(5.3) \quad \|z^{-1}(z^\alpha - \Delta_h)^{-1} - \mu(e^{-z\tau})^{-1}(\beta_\tau(e^{-z\tau}) - \Delta_h)^{-1}\|_{L^2(\Omega) \rightarrow L^2(\Omega)} \leq c|z|^{-\alpha}\tau.$$

Now we split the error  $v_h(t_n) - v_h^n$  into two components, i.e.,  $v_h(t_n) - v_h^n = \mathcal{I}_1 + \mathcal{I}_2$ , where

$$\begin{aligned} \mathcal{I}_1 &= \frac{1}{2\pi i} \int_{\Gamma_{\theta,\delta}^\tau} e^{zt_n} (z^{-1}(z^\alpha - \Delta_h)^{-1} \\ &\quad - \mu(e^{-z\tau})^{-1}(\beta_\tau(e^{-z\tau}) - \Delta_h)^{-1})(\Delta_h v_h(0) + P_h g(0)) \, dz, \\ \mathcal{I}_2 &= \frac{1}{2\pi i} \int_{\Gamma_{\theta,\delta} \setminus \Gamma_{\theta,\delta}^\tau} e^{zt_n} z^{-1}(z^\alpha - \Delta_h)^{-1}(\Delta_h v_h(0) + P_h g(0)) \, dz. \end{aligned}$$

By using (5.3) and (3.7), and choosing  $\delta \leq 1/t_n$ , the argument from [15] yields

$$(5.4) \quad \|\mathcal{I}_1\|_{L^2(\Omega)} + \|\mathcal{I}_2\|_{L^2(\Omega)} \leq ct_n^{\alpha-1}\tau \|\Delta_h v_h(0) + P_h g(0)\|_{L^2(\Omega)}.$$

Second, we consider the case  $v(0) = g(0) = 0$ . Then Taylor’s expansion gives

$$(5.5) \quad P_h g(t) = P_h g(0) + 1 * P_h g'(t) = 1 * P_h g'(t).$$

In view of (3.9), the semidiscrete solution  $v_h(t_n)$  can be represented by

$$(5.6) \quad v_h(t_n) = (E * P_h g)(t_n) = (E * (1 * P_h g'))(t_n) = ((E * 1) * P_h g')(t_n).$$

Similarly, we have

$$(\beta_\tau(\xi) - \Delta_h)^{-1} = \sum_{n=0}^{\infty} E_\tau^n \xi^n \quad \text{with} \quad E_\tau^n = \frac{\tau}{2\pi i} \int_{\Gamma_{\theta,\delta}^\tau} e^{zn\tau} (\beta_\tau(e^{-z\tau}) - \Delta_h)^{-1} \, dz.$$

Hence the fully discrete solution  $v_h^n$  can be represented by  $v_h^n = \sum_{j=0}^n E_\tau^{n-j} P_h g(t_j)$ , and the second inequality of (5.2) implies

$$(5.7) \quad \|E_\tau^n\|_{L^2(\Omega) \rightarrow L^2(\Omega)} \leq ct_n^{\alpha-1}\tau.$$

Let  $E_{\tau,\epsilon}(t) = \sum_{n=0}^{\infty} E_\tau^n \delta_{t_n-\epsilon}(t)$ , where  $\delta_{t_n-\epsilon}$  is the Dirac delta function concentrated at  $t_n - \epsilon$  with  $\epsilon \in (0, \tau)$ . Then  $v_h^n$  can be rewritten as

$$(5.8) \quad v_h^n = \lim_{\epsilon \rightarrow 0} (E_{\tau,\epsilon} * P_h g)(t_n) = \lim_{\epsilon \rightarrow 0} (E_{\tau,\epsilon} * (1 * P_h g'))(t_n) = \left( \lim_{\epsilon \rightarrow 0} (E_{\tau,\epsilon} * 1) * P_h g' \right)(t_n).$$

The representations (5.6) and (5.8) yield

$$(5.9) \quad \|v_h(t_n) - v_h^n\|_{L^2(\Omega)} \leq \left\| \lim_{\epsilon \rightarrow 0} ((E - E_{\tau,\epsilon}) * 1) * P_h g' \right\|_{L^2(\Omega)}.$$

Using the Laplace transform and Cauchy's integral formula, we deduce

$$\begin{aligned} \left(\lim_{\epsilon \rightarrow 0} (E - E_{\tau, \epsilon}) * 1\right)(t_n) &= \frac{1}{2\pi i} \int_{\Gamma_{\theta, \delta}} e^{zt_n} z^{-1} (z^\alpha - \Delta_h)^{-1} dz \\ &\quad - \frac{1}{2\pi i} \int_{\Gamma_{\theta, \delta}^\tau} e^{zt_n} \mu(e^{-z\tau})^{-1} (\beta_\tau(e^{-z\tau}) - \Delta_h)^{-1} dz. \end{aligned}$$

Then using the estimate (5.3) we obtain

$$(5.10) \quad \left\| \left(\lim_{\epsilon \rightarrow 0} (E - E_{\tau, \epsilon}) * 1\right)(t_n) \right\|_{L^2(\Omega) \rightarrow L^2(\Omega)} \leq c\tau t_n^{\alpha-1}.$$

It remains to prove the following extension of the estimate (5.10):

$$(5.11) \quad \left\| \left(\lim_{\epsilon \rightarrow 0} (E - E_{\tau, \epsilon}) * 1\right)(t) \right\|_{L^2(\Omega) \rightarrow L^2(\Omega)} \leq c\tau t^{\alpha-1} \quad \forall t \in (0, T).$$

Then this and (5.9) yield the second part on the right-hand side of (4.2), which completes the proof of Lemma 4.2.

To prove (5.11), we consider the Taylor expansion of  $(E(t) - E_{\tau, \epsilon}(t)) * 1$  at  $t = t_n$ , i.e.,

$$(5.12) \quad ((E - E_{\tau, \epsilon}) * 1)(t) = ((E - E_{\tau, \epsilon}) * 1)(t_n) - \int_t^{t_n} (E - E_{\tau, \epsilon})(s) ds.$$

In view of Lemma 3.4(iii), there holds

$$\left\| \int_t^{t_n} E(s) ds \right\|_{L^2(\Omega) \rightarrow L^2(\Omega)} \leq c \int_t^{t_n} s^{\alpha-1} ds \leq c\tau t^{\alpha-1}.$$

Similarly, appealing to (5.7), we have

$$\left\| \lim_{\epsilon \rightarrow 0} \int_t^{t_n} E_{\tau, \epsilon}(s) ds \right\|_{L^2(\Omega) \rightarrow L^2(\Omega)} = \|E_\tau^n\|_{L^2(\Omega) \rightarrow L^2(\Omega)} \leq ct_n^{\alpha-1}\tau.$$

Substituting (5.10) and the last two inequalities into (5.12) yields (5.11).  $\square$

**6. Numerical experiments.** In this section, we present numerical examples to verify the theoretical results in Theorems 4.4 and 4.5. We consider problem (1.1) with a diffusion coefficient 0.1 in the unit square  $\Omega = (0, 1)^2$  with the following two sets of problem data:

- (a)  $u_0(x, y) = xy(1-x)(1-y) \in H_0^1(\Omega) \cap H^2(\Omega)$  and  $f = \sqrt{1+u^2}$ ;
- (b)  $u_0(x, y) = x(1-x)\sin(2\pi y) \in H_0^1(\Omega) \cap H^2(\Omega)$  and  $f = 1 - u^3$ .

In the computation, we divided the domain  $\Omega$  into regular right triangles with  $M$  equal subintervals of length  $h = 1/M$  on each side of the domain. The numerical solutions are computed by using the Galerkin FEM in space and the backward Euler CQ or the L1 scheme in time. To evaluate the convergence, we compute the spatial error  $e_t$  and temporal error  $e_s$ , respectively, defined by

$$e_s = \max_{1 \leq n \leq N} \|u_h(t_n) - u(t_n)\|_{L^2(\Omega)} \quad \text{and} \quad e_t = \max_{1 \leq n \leq N} \|u_h^n - u_h(t_n)\|_{L^2(\Omega)}.$$

Since the exact solution to problem (1.1) is unavailable, we compute reference solutions on a finer mesh, i.e., the continuous solution  $u(t_n)$  with a fixed time step  $\tau = 1/1000$

and mesh size  $h = 1/1280$ , and the semidiscrete solution  $u_h(t_n)$  with  $h = 1/10$  and  $\tau = 1/(64 \times 10^4)$ .

In case (a), since the nonlinearity  $f$  is Lipschitz continuous, the theory in section 4 applies. The numerical results for case (a) are shown in Tables 1 and 2, where the numbers in the brackets in the last column refer to the theoretical predictions from section 4. We observe an  $O(h^2)$  rate for the spatial error  $e_s$  and an  $O(\tau^\alpha)$  rate for the temporal error  $e_t$  for both the backward Euler CQ and the L1 scheme. These observations fully confirm Theorems 4.4 and 4.5.

In case (b), the nonlinear source  $f$  is not Lipschitz continuous. Nonetheless, one observes an  $O(h^2)$  and  $O(\tau^\alpha)$  convergence rate for the spatial and temporal errors, respectively; cf. Tables 3 and 4. This concurs with the discussions in Remarks 3.2 and 4.6. Further, the absolute accuracy of the L1 scheme and backward Euler CQ is comparable with each other for both cases (a) and (b). Interestingly, the spatial error  $e_s$  increases slightly with the fractional order  $\alpha$ , but the temporal error  $e_t$  decreases with  $\alpha$ .

TABLE 1

Numerical results for case (a): the spatial error  $e_s$  with  $T = 1$ , with  $N = 1000$ ,  $h = 1/M$ .

$\alpha \backslash M$	5	10	20	40	80	Rate
0.4	6.89e-2	2.00e-2	5.34e-3	1.37e-3	3.31e-4	$\approx 2.01$ (2.00)
0.6	7.06e-2	2.05e-2	5.58e-3	1.42e-3	3.44e-4	$\approx 2.01$ (2.00)
0.8	7.59e-2	2.18e-2	5.80e-3	1.48e-3	3.57e-4	$\approx 2.01$ (2.00)

TABLE 2

Numerical results for case (a): the temporal error  $e_t$  with  $T = 1$ ,  $\tau = T/N$ ,  $N = k \times 10^4$ , and  $h = 0.1$ . BE = backward Euler.

$\alpha$	$k$	1	2	4	8	16	Rate
0.4	BE	1.16e-3	8.88e-4	6.79e-4	5.19e-4	3.86e-4	$\approx 0.39$ (0.40)
	L1	2.06e-3	1.59e-3	1.22e-3	9.34e-4	7.15e-4	$\approx 0.38$ (0.40)
0.6	BE	1.79e-4	1.18e-4	7.75e-5	5.10e-5	3.36e-5	$\approx 0.60$ (0.60)
	L1	3.05e-4	2.02e-4	1.33e-4	8.80e-5	5.81e-5	$\approx 0.60$ (0.60)
0.8	BE	1.73e-5	9.87e-6	5.65e-6	3.24e-6	1.86e-6	$\approx 0.80$ (0.80)
	L1	3.91e-5	2.24e-5	1.29e-5	7.38e-6	4.24e-6	$\approx 0.80$ (0.80)

TABLE 3

Numerical results for case (b): the spatial error  $e_s$  with  $T = 1$ , with  $N = 1000$ ,  $h = 1/M$ .

$\alpha \backslash M$	5	10	20	40	80	Rate
0.4	5.65e-2	1.68e-2	4.58e-3	1.18e-3	2.87e-4	$\approx 2.00$ (2.00)
0.6	5.90e-2	1.75e-2	4.74e-3	1.22e-3	2.97e-4	$\approx 2.00$ (2.00)
0.8	6.19e-2	1.82e-2	4.93e-3	1.27e-3	3.08e-4	$\approx 2.01$ (2.00)

TABLE 4

Numerical results for case (b): the temporal error  $e_t$  with  $T = 1$ ,  $\tau = T/N$ ,  $N = k \times 10^4$ ,  $h = 0.1$ .

$\alpha$	$k$	1	2	4	8	16	Rate
0.4	BE	1.53e-3	1.17e-3	9.07e-4	6.96e-4	5.33e-4	$\approx 0.38$ (0.40)
	L1	2.73e-3	2.12e-3	1.64e-3	1.26e-3	9.65e-4	$\approx 0.38$ (0.40)
0.6	BE	2.43e-4	1.60e-4	1.05e-4	6.93e-5	4.56e-5	$\approx 0.60$ (0.60)
	L1	4.14e-4	2.74e-4	1.81e-4	1.20e-4	7.89e-5	$\approx 0.60$ (0.60)
0.8	BE	2.35e-5	1.34e-5	7.68e-6	4.40e-6	2.53e-6	$\approx 0.80$ (0.80)
	L1	5.30e-5	3.04e-5	1.75e-5	1.00e-5	5.76e-6	$\approx 0.80$ (0.80)

**Acknowledgment.** The authors are grateful to the anonymous referees for their constructive comments, which were very helpful for improving the presentation of the paper.

## REFERENCES

- [1] G. AKRIVIS AND B. LI, *Maximum norm analysis of implicit-explicit backward difference formulae for nonlinear parabolic equations*, IMA J. Numer. Anal., to appear, <https://doi.org/10.1093/imanum/drx008>.
- [2] G. AKRIVIS, B. LI, AND C. LUBICH, *Combining maximal regularity and energy estimates for time discretizations of quasilinear parabolic equations*, Math. Comp., 86 (2017), pp. 1527–1552.
- [3] A. A. ALIKHANOV, *A new difference scheme for the time fractional diffusion equation*, J. Comput. Phys., 280 (2015), pp. 424–438.
- [4] W. ARENDT, C. J. BATTY, M. HIEBER, AND F. NEUBRANDER, *Vector-valued Laplace Transforms and Cauchy Problems*, 2nd ed., Birkhäuser, Basel, 2011.
- [5] E. BAZHLEKOVA, *Strict  $L^p$  solutions for fractional evolution equations*, Fract. Calc. Appl. Anal., 5 (2002), pp. 427–436.
- [6] B. BERKOWITZ, J. KLAFTER, R. METZLER, AND H. SCHER, *Physical pictures of transport in heterogeneous media: Advection-dispersion, random-walk, and fractional derivative formulations*, Water Resources Res., 38 (2002), pp. 9-1–9-12.
- [7] S. BLUNCK, *Maximal regularity of discrete and continuous time evolution equations*, Stud. Math., 146 (2001), pp. 157–176.
- [8] C. CHEN, V. THOMÉE, AND L. B. WAHLBIN, *Finite element approximation of a parabolic integro-differential equation with a weakly singular kernel*, Math. Comp., 58 (1992), pp. 587–602.
- [9] D. M. CREEDON AND J. J. H. MILLER, *The stability properties of  $q$ -step backward difference schemes*, BIT, 15 (1975), pp. 244–249.
- [10] E. CUESTA, C. LUBICH, AND C. PALENCIA, *Convolution quadrature time discretization of fractional diffusion-wave equations*, Math. Comp., 75 (2006), pp. 673–696.
- [11] P. FLAJOLET, *Singularity analysis and asymptotics of Bernoulli sums*, Theoret. Comput. Sci., 215 (1999), pp. 371–381.
- [12] E. HAIRER, S. P. NØRSETT, AND G. WANNER, *Solving Ordinary Differential Equations. I, Nonstiff Problems*, 2nd ed., Springer-Verlag, Berlin, 2010.
- [13] B. JIN, R. LAZAROV, J. PASCIAK, AND Z. ZHOU, *Error analysis of semidiscrete finite element methods for inhomogeneous time-fractional diffusion*, IMA J. Numer. Anal., 35 (2015), pp. 561–582.
- [14] B. JIN, R. LAZAROV, AND Z. ZHOU, *Error estimates for a semidiscrete finite element method for fractional order parabolic equations*, SIAM J. Numer. Anal., 51 (2013), pp. 445–466.
- [15] B. JIN, R. LAZAROV, AND Z. ZHOU, *An analysis of the  $L_1$  scheme for the subdiffusion equation with nonsmooth data*, IMA J. Numer. Anal., 36 (2016), pp. 197–221.
- [16] B. JIN, R. LAZAROV, AND Z. ZHOU, *Two fully discrete schemes for fractional diffusion and diffusion-wave equations with nonsmooth data*, SIAM J. Sci. Comput., 38 (2016), pp. A146–A170.
- [17] B. JIN, B. LI, AND Z. ZHOU, *Discrete maximal regularity of time-stepping schemes for fractional evolution equations*, Numer. Math., to appear, <https://doi.org/10.1007/s00211-017-0904-8>.
- [18] T. KEMMOCHI, *Discrete maximal regularity for abstract Cauchy problems*, Stud. Math., 234 (2016), pp. 241–263.
- [19] A. A. KILBAS, H. M. SRIVASTAVA, AND J. J. TRUJILLO, *Theory and Applications of Fractional Differential Equations*, Elsevier Science, Amsterdam, 2006.
- [20] S. KOU, *Stochastic modeling in nanoscale biophysics: Subdiffusion within proteins*, Ann. Appl. Stat., 2 (2008), pp. 501–535.
- [21] B. KOVÁCS, B. LI, AND C. LUBICH,  *$A$ -stable time discretizations preserve maximal parabolic regularity*, SIAM J. Numer. Anal., 54 (2016), pp. 3600–3624.
- [22] P. C. KUNSTMANN, B. LI, AND C. LUBICH, *Runge-Kutta time discretization of nonlinear parabolic equations studied via discrete maximal parabolic regularity*, Found. Comput. Math., to appear, <https://doi.org/10.1007/s10208-017-9364-x>.
- [23] P. C. KUNSTMANN AND L. WEIS, *Maximal  $L_p$ -regularity for parabolic equations, Fourier multiplier theorems and  $H^\infty$ -functional calculus*, in Functional Analytic Methods for Evolution Equations, Lecture Notes in Math. 1855, Springer-Verlag, Berlin, 2004, pp. 65–311.
- [24] T. A. M. LANGLANDS AND B. I. HENRY, *The accuracy and stability of an implicit solution method for the fractional diffusion equation*, J. Comput. Phys., 205 (2005), pp. 719–736.

- [25] D. LEYKEKHMAN AND B. VEXLER, *Discrete maximal parabolic regularity for Galerkin finite element methods*, Numer. Math., 135 (2017), pp. 923–952.
- [26] Y. LIN AND C. XU, *Finite difference/spectral approximations for the time-fractional diffusion equation*, J. Comput. Phys., 225 (2007), pp. 1533–1552.
- [27] C. LUBICH, *Discretized fractional calculus*, SIAM J. Math. Anal., 17 (1986), pp. 704–719.
- [28] C. LUBICH, *Convolution quadrature and discretized operational calculus. I*, Numer. Math., 52 (1988), pp. 129–145.
- [29] Y. LUCHKO, W. RUNDELL, M. YAMAMOTO, AND L. ZUO, *Uniqueness and reconstruction of an unknown semilinear term in a time-fractional reaction-diffusion equation*, Inverse Problems, 29 (2013), 065019.
- [30] W. MCLEAN, *Regularity of solutions to a time-fractional diffusion equation*, ANZIAM J., 52 (2010), pp. 123–138.
- [31] W. MCLEAN AND K. MUSTAPHA, *Time-stepping error bounds for fractional diffusion problems with non-smooth initial data*, J. Comput. Phys., 293 (2015), pp. 201–217.
- [32] R. METZLER AND J. KLAFTER, *The random walk's guide to anomalous diffusion: A fractional dynamics approach*, Phys. Rep., 339 (2000), pp. 1–77.
- [33] K. MUSTAPHA, B. ABDALLAH, AND K. M. FURATI, *A discontinuous Petrov-Galerkin method for time-fractional diffusion equations*, SIAM J. Numer. Anal., 52 (2014), pp. 2512–2529.
- [34] K. MUSTAPHA AND H. MUSTAPHA, *A second-order accurate numerical method for a semilinear integro-differential equation with a weakly singular kernel*, IMA J. Numer. Anal., 30 (2010), pp. 555–578.
- [35] R. R. NIGMATULIN, *The realization of the generalized transfer equation in a medium with fractal geometry*, Phys. Stat. Sol. B, 133 (1986), pp. 425–430.
- [36] K. SAKAMOTO AND M. YAMAMOTO, *Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems*, J. Math. Anal. Appl., 382 (2011), pp. 426–447.
- [37] Z.-Z. SUN AND X. WU, *A fully discrete scheme for a diffusion wave system*, Appl. Numer. Math., 56 (2006), pp. 193–209.
- [38] V. THOMÉE, *Galerkin Finite Element Methods for Parabolic Problems*, 2nd ed., Springer-Verlag, Berlin, 2006.
- [39] S. B. YUSTE, *Weighted average finite difference methods for fractional diffusion equations*, J. Comput. Phys., 216 (2006), pp. 264–274.
- [40] F. ZENG, C. LI, F. LIU, AND I. TURNER, *Numerical algorithms for time-fractional subdiffusion equation with second-order accuracy*, SIAM J. Sci. Comput., 37 (2015), pp. A55–A78.