

New ^{234}U - ^{230}Th coral dates from the western Gulf of Corinth: Implications for extensional tectonics

Sarah L. Houghton, Gerald P. Roberts, Ioannis D. Papanikolaou, and John M. McArthur

Research School of Earth Sciences at UCL-Birkbeck, London, UK

Mabs A. Gilmour

Uranium Series Facility, Department of Earth Sciences, The Open University, Milton Keynes, UK

Received 7 July 2003; revised 1 September 2003; accepted 3 September 2003; published 14 October 2003.

[1] We derive rates of uplift of ~ 0.7 – 0.8 mm/yr for the western end of the Gulf of Corinth, Greece, using geomorphic paleashoreline modeling. We calibrate the modeling with new ^{234}U - ^{230}Th dates on the coral *Cladocora caespitosa* collected from raised marine terraces uplifted in the footwall of the active Psathopyrgos fault, the only major active normal fault, reported on published maps controlling the downthrown Rio Straits at the western end of the Gulf of Corinth. In this area of high (15–22 mm/yr) extension rates measured with GPS, the ratio of uplift-rate to extensional velocity is 0.025–0.035, much lower than values of 0.15–0.25 found further east in the gulf. These low values imply that if GPS extension rates are correct then mechanical/kinematic models developed for the eastern and central gulf may not be applicable to the western gulf. **INDEX TERMS:** 1035 Geochemistry: Geochronology; 8109 Tectonophysics: Continental tectonics—extensional (0905); 8107 Tectonophysics: Continental neotectonics. **Citation:** Houghton, S. L., G. P. Roberts, I. D. Papanikolaou, J. M. McArthur, and M. A. Gilmour, New ^{234}U - ^{230}Th coral dates from the western Gulf of Corinth: Implications for extensional tectonics, *Geophys. Res. Lett.*, 30(19), 2003, doi:10.1029/2003GL018112, 2003.

1. Introduction

[2] The Gulf of Corinth in central Greece is a major N-S extensional rift basin in the Hellenic region characterized by intense seismic activity. Continental extension rates of up to 15–22 mm/yr have been measured using global positioning systems in the western Gulf of Corinth, Greece [GPS; Clarke *et al.*, 1997; Briole *et al.*, 2000] with progressively lower rates reported for successively more eastern positions in the central and eastern gulf (Figure 1). Rates of extension measured by GPS in the central and eastern gulf are comparable with rates estimated from offset geology, both over the Quaternary [Armijo *et al.*, 1996], and/or fractions of the Holocene [Collier *et al.*, 1998].

[3] Comparison of uplift rates gained from ^{234}U - ^{230}Th dates on *C. caespitosa* corals from elevated marine terraces with ‘horizontal’ GPS extension rates in the central and eastern gulf (extension/uplift ratios of 0.15–0.25) supports the development of mechanical/kinematic models that are consistent with observed deformation [Armijo *et al.*, 1996]. However, to date, without uplift rates for the Psathopyrgos fault, the only major active normal fault (within its neigh-

boring longitude), reported on published maps controlling the downthrown Rio Straits at the western end of the Gulf of Corinth, it has not been possible to see whether tectonic models developed for the central and eastern gulf are also applicable at its western end.

[4] This paper reports the results of geomorphic paleoshoreline modelling of a flight of uplifted marine terraces in the immediate footwall of the Psathopyrgos fault at the western end of the Gulf of Corinth. We have mapped the terraces at 1:5000 scale and modeled their uplift, calibrating our results with new ^{234}U - ^{230}Th dates corals on *C. caespitosa* corals associated with these terraces. This allows us to identify the ages and elevations of the terraces in order to constrain the uplift rate and compare with the extension rate from GPS data at this site.

2. Existing Data on Deformation Rates in the Gulf of Corinth

[5] Rates of extension across the Gulf, measured with GPS show that relative site velocities between the north and south sides increase smoothly from 2.5 to 10 ± 4 mm/yr across the eastern gulf to 12.7 ± 1 to 22 mm/yr in the west; these are termed GPS extension rates [Figure 1; Briole *et al.*, 2000; Clarke *et al.*, 1997, 1998].

[6] Uplift rates for the footwalls of the faults accommodating the extension on the northern coast of the Peloponnese have been constrained using uplifted and dated *C. caespitosa* corals for a number of sites along the gulf, with the exception of the western gulf [Figure 1; Collier, 1990; Collier *et al.*, 1992; Vita-Finzi, 1993; Dia *et al.*, 1997; Leeder *et al.*, 2003]. Armijo *et al.* [1996] and Morewood and Roberts [1999] have mapped the marine terraces associated with these dated corals, using the results to constrain the development of normal faulting for the gulf. In particular, Armijo *et al.* [1996] use a thick elastic plate model, plus a map of the uplifted marine terraces, to calculate extension rates across the gulf. They show that implied extension rates for portions of the Quaternary are similar to those measured over much shorter time periods using GPS. Extension rates from a palaeoseismic trenching study of Holocene sediments by Collier *et al.* [1998] also agree with those from GPS.

3. Uplifted Marine Terraces and Corals in the Western Gulf

[7] Marine terraces result from the interplay between the rate of global sea-level change and the rate of local tectonic




Figure 1. Map of central Greece showing major on-shore and off-shore faults [Roberts, 1996; Stefatos *et al.*, 2003], plate velocities [Briole *et al.*, 2000] and 'horizontal' GPS velocities with Location map 1 at the western end of the Gulf of Corinth which is enlarged below. Location map 1 shows the Psathopyrgos fault and geomorphic 2D cross section A–A'. HT- Hellenic Trench. See color version of this figure in the HTML.

uplift [e.g. Chappell, 1974; Lajoie, 1986; Valensise and Ward, 1991]. The position of sea-level at the time of terrace formation is defined by the base of palaeo-sea-cliffs; their elevations are precise markers of the amount of uplift from the sea-level at which they formed; we term such a marker an inner edge.

[8] We have constructed topographic profiles across the marine terraces mapped at a 1:5000 scale with contours at 4 metre intervals in order to identify inner edges in the immediate footwall of the Psathopyrgos fault (Figure 1; cross-section A–A'). Inflections in the profiles are interpreted as palaeosea cliffs (palaeoshoreline inner edges), verified by field observations. The elevations of inner edges allow precise estimates to be made of the amount of uplift from knowledge of the elevation of sea level at the time of their formation. The elevation of sea level at such times is determined from a paleosea-level reconstruction based on

the $\delta^{18}\text{O}$ record of sediment cores from the Red Sea [Siddall *et al.*, 2003].

[9] In order to assess the ages of these palaeoshorelines, we have modeled elevations of inner edges, with uplift rate as a variable (Figure 2). We use uplift rates that are constant through time, an approach that is consistent with the results of other workers [Armijo *et al.*, 1996, Morewood and Roberts, 1999]. Present-day elevations are modeled iteratively by assuming a constant uplift rate of given value and attempting to match the modeled terraces elevations with elevations of topographic inner edges. Highstands are not equally-spaced in time and have different initial elevations, so a fit to topographic inflections that are not equally-spaced in elevation can be considered to be robust. Using the sea-level curve [Figure 2b; Siddall *et al.*, 2003], the best fit to heights of terrace inner edges indicates an uplift rate of 0.8 mm/yr. A better fit is achieved if the original elevations




Figure 2. Graphs showing affect of 0.8 mm/yr (a) and 0.7 mm/yr (c) uplift on flight of terraces near the Holy Metamorphoses Church, near Agios Ioannis, given the sea-level curves [Siddall *et al.*, 2003] in (b) and (d). See color version of this figure in the HTML.

Table 1. U/Th Chemical and Isotopic Data for *C. caespitosa* Coral

Sample	^{238}U [ppm]	^{230}Th [ppb]	($^{230}\text{Th}/^{234}\text{U}$)	($^{230}\text{Th}/^{232}\text{Th}$)	^{230}Th AGE [kyr]	$\delta^{234}\text{U}(0)$	$\delta^{234}\text{U}(T)$
AIC wall	2.074 ± 4	0.0299 ± 3	0.813 ± 9	1500 ± 30	175.6 ^{+5.4} _{-5.3}	82.4	135.4
AIC wall	2.065 ± 4	0.0294 ± 3	0.811 ± 9	1460 ± 28	175.2 ^{+5.2} _{-5.1}	74.0	121.4
AIC septa	2.466 ± 12	0.0369 ± 4	0.816 ± 9	524 ± 10	174.7 ^{+6.2} _{-5.9}	120	196.7
MCC1 wall	2.214 ± 5	0.0323 ± 4	0.838 ± 9	1600 ± 30	191.3 ^{+6.5} _{-6.5}	63.1	108.3
MCC2 wall	2.211 ± 5	0.0288 ± 3	0.748 ± 8	1190 ± 22	147.4 ^{+2.9} _{-4.3}	63.8	96.6

Round brackets denotes activity ratio. Contributions from detrital ^{232}Th were too small to require correction.

of highstands at 175 ka and 290 ka are increased by 12 metres (Figure 2d), which is not unreasonable given the uncertainty on the precise elevations of these highstands. This adjustment results in an uplift rate of 0.7 mm/yr. With this estimate of terrace ages, we carried out dating of corals to validate the ages and thus the uplift rates.

4. Samples and Analysis

[10] *C. caespitosa* coral co-occurred with oysters and gastropods within poorly exposed green-yellow marls at a height of ~100–110 m above mean sea level. Coral samples were mechanically and chemically cleaned prior to analysis by TIMS to determine ^{234}U - ^{230}Th ages. Sample preparation and analysis followed standard methods [Edwards *et al.*, 1987; van Calsteren and Schwieter, 1995]. Prepared samples were assessed for alteration using XRD and elemental analysis. The results of the chemical and isotopic analysis are given next and in Table 1.

5. Chemical and Isotopic Data

[11] The elemental composition of the samples is typical of modern corals (Table 1): concentrations of U (2–3 ppm) and Ca (38.5–41.8%) are typical of those reported for recent corals [Stirling *et al.*, 1995]. Concentrations of Sr (5600–7770 ppm) are similar to those given for corals by Stein *et al.* [1993], and concentrations of Mg (830–1180 ppm) and Na (3520–3820 ppm) are within the range given for coral by Bar-Matthews *et al.* [1993].

[12] The $^{234}\text{U}/^{238}\text{U}$ activity ratio of the modern ocean is 1.145 ($\delta^{234}\text{U}(T) = 145$), and samples with an initial $^{234}\text{U}/^{238}\text{U}$ activity ratio close to 1.149 (i.e., $\delta^{234}\text{U}(T) 149 \pm 4$; a limit introduced by Stirling *et al.* [1995] to constrain the 125ka highstand) are considered to give the most reliable ages (*ibid*). Nevertheless, these authors acknowledge that many samples with $\delta^{234}\text{U}(T)$ very different from 149 give reliable numerical ages. For sample AIC, two subsamples of coral wall yield similar ages which, in turn, are similar to the age derived from coral septa of the same sample, despite the latter containing trace calcite identifiable by XRD. We consider these ages as valid because of the concordance in ages, despite the range of $\delta^{234}\text{U}(T)$: 121–197.

[13] Analysis of two other samples (MCC1 & 2) give irreproducible ages and more extreme $\delta^{234}\text{U}(T)$ values, although we note that the mean age for MCC is 169 ± 22 ka, a value within analytical uncertainty of the age of 175 ka we obtain for sample AIC.

[14] We suggest that an age of 175 ka for our terrace is sufficiently robust to link the terrace to the highstand of sealevel at 175 ka noted by Siddall *et al.* [2003], although it might be insufficiently robust to calibrate the time of sealevel highstands. Hence, an age of 175 ka is established for the raised marine terrace at 100–110m elevation, which is

consistent with geomorphic modeling of paleoshorelines for this suite of terraces.

6. Implications for Tectonic Models

[15] Using a constant uplift rate of 0.7 mm/year (Figure 2c), the steps in our topographic profile, confirmed as terraces and paleosea cliffs by field observation, co-incide almost exactly with the expected elevations of terraces formed during sealevel highstands. In particular, the elevation of the terrace we date to 175 ka corresponds closely to the height expected for a terrace of this age formed during the 175 ka sealevel highstand [Siddall *et al.*, 2003].




Figure 3. Graphs and map showing (i) uplift rates determined from ^{234}U - ^{230}Th dating of corals and (ii) ‘Horizontal’ GPS site velocities for the Gulf of Corinth and (iii) (Uplift rate/‘Horizontal’ GPS site velocity). Data points on graph (i) relate to uplift rates gained from the following sources: 1. New uplift rate from this paper; 2. Armijo *et al.* [1996] with position 2 extrapolated 5km further north corresponding to the fault trace; 3. Morewood and Roberts [1999]; 4. Collier *et al.* [1992]; 5. Rigo [1994]. See color version of this figure in the HTML.

[16] The extension rate measured with GPS across the Rio Strait, in the western gulf, is about 20 mm/yr (Figure 3). GPS-derived extension rates for the longitudes of Xylokastron and Alepochori [c. 7–14 mm/yr; 1–3 mm/yr; Clarke *et al.*, 1997] are consistent with slip rates derived from rigorous modeling of uplifted and deformed Quaternary marine terraces [9.2–15.6 mm/yr; 1.8 mm/yr; Armijo *et al.*, 1996]. This implies that mechanical models can be developed which link rates of uplift and horizontal extension which may help understanding of extension elsewhere.

[17] The uplift rates south of Xylokastron, Heraion and Alepochori are 1.28 mm/yr, 0.5–0.74 mm/yr and 0.31 mm/yr respectively. Thus, the ratio of uplift (from raised marine terraces) to extension (from GPS) is 0.26–0.11 for Xylokastron, 0.071–0.11 for Heraion, and 0.31–0.10 for Alepochori. The value for this ratio is much less (0.025–0.035) where we have studied the deformation. If both the GPS extension rates and uplift rates from marine shorelines reported herein are correct then this ratio specifically modeled by Armijo *et al.* [1996] in their attempt to constrain the mechanics and kinematics of the extension for the central and eastern gulf implies that different mechanical/kinematic models for the deformation may apply to the western gulf.

[18] **Acknowledgments.** This study was supported by NERC IP/657/0300 Part 1 & 2 (JMcA), the UCL Graduate School (SH), the Mineralogical Society of Great Britain & Ireland (SH), State Scholarship Foundation of Greece, I.K.Y.(IP), Birkbeck College and Benfield Re-insurance plc.

[19] We thank IGME (Athens) for permission to conduct field studies in Greece and Dr. B. Rosen (Natural History Museum) for identification of *C. caespitosa*. Elemental analyses were done at the NERC ICP-AES facility at Royal Holloway, University of London, with permission from its director Dr. J. N. Walsh and U/Th analyses were carried out at Uranium Series Facility, Department of Earth Sciences, The Open University, Milton Keynes. We also thank S. Hirons for XRD analysis, Dr. P. van Calsteren and J. Rhodes for help with U/Th isotope work. We thank M. Leeder and an anonymous reviewer for their helpful comments.

References

- Armijo, R., B. Meyer, G. C. P. King, A. Rigo, and D. Papanastassiou, Quaternary evolution of the Corinth Rift and its implications for the Late Cenozoic evolution of the Aegean, *Geophys. J. Int.*, **126**, 11–53, 1996.
- Bar-Matthews, M., G. J. Wasserburg, and J. H. Chen, Diagenesis of fossil coral skeletons: Correlation between trace elements, texture and $^{234}\text{U}/^{238}\text{U}$, *Geochim. Cosmochim. Acta.*, **57**, 257–276, 1993.
- Briole, P., A. Rigo, H. Lyon-Caen, J. C. Ruegg, K. Papazissi, C. Mitsakaki, A. Balodimou, G. Veis, D. Hatzfeld, and A. Deschamps, Active deformation of the Corinth rift, Greece: Results from repeated Global Positioning System surveys between 1990 and 1995, *J. Geophys. Res., Solid Earth*, **21**, 25,605–25,625, 2000.
- Chappell, J., Geology of coral terraces, Huon Peninsula, New Guinea: A study of tectonic movements and sea-level changes, *Geol. Soc. Am. Bull.*, **85**, 553–570, 1974.
- Clarke, P. J., R. R. Davies, P. C. England, B. Parsons, H. Billiris, D. Paradissis, G. Veis, P. H. Denys, P. A. Cross, V. Ashkenazi, and R. Bingley, Geodetic estimate of seismic hazard in the Gulf of Korinthis, *Geophys. Res. Lett.*, **24**, 1303–1306, 1997.
- Clarke, P. J., R. R. Davies, P. C. England, B. Parsons, H. Billiris, D. Paradissis, G. Veis, P. A. Cross, P. H. Denys, V. Ashkenazi, R. Bingley, H. G. Kahle, M. V. Muller, and P. Briole, Crustal strain in central Greece from repeated GPS measurements in the interval 1989–1997, *Geophys. J. Int.*, **135**, 195–214, 1998.
- Collier, R. E. L., Eustatic and tectonic controls upon Quaternary coastal sedimentation in the Corinth Basin, Greece, *J. Geol. Soc. London*, **147**, 301–314, 1990.
- Collier, R. E. L., M. R. Leeder, P. J. Rowe, and T. C. Atkinson, Rates of tectonic uplift in the Corinth and Megara basins, Central Greece, *Tectonics*, **11**(6), 1159–1167, 1992.
- Collier, R. E. L., D. Pantosti, G. D'Addezio, P. M. De Martini, E. Masana, and D. Sakellariou, Paleoseismicity of the 1981 Corinth earthquake fault: Seismic contribution to extensional strain in central Greece and implications for seismic hazard, *J. Geophys. Res.*, **103**, 30,001–30,019, 1998.
- Dia, A. N., A. S. Cohen, R. K. O'Nions, and J. A. Jackson, Rates of uplift investigated through ^{230}Th Dating in the Gulf of Corinth (Greece), *Chem. Geol.*, **138**, 171–184, 1997.
- Edwards, L. E., J. H. Chen, and G. J. Wasserburg, ^{238}U - ^{234}U - ^{230}Th - ^{232}Th systematics and the precise measurement of time over the past 500 000 years, *Earth Planet. Sci. Lett.*, **81**, 175–192, 1987.
- Lajoie, K. R., Coastal tectonics, in Active Tectonics, *National Academic*, Washington, DC, 95–124, 1986.
- Leeder, M. R., L. C. McNeill, R. E. L. Collier, C. Portman, W. J. Rowe, J. E. Andrews, and R. L. Gawthorpe, Corinth rift margin uplift: New evidence from Late Quaternary marine shorelines, *Geophys. Res. Lett.*, **30**(12), 1611–1615, 2003.
- Morewood, N. C., and G. P. Roberts, Lateral propagation of the surface trace of the South Alkyonides normal fault segment, central Greece: Its impact on models of fault growth and displacement-length relationships, *J. Struct. Geol.*, **24**, 3081–3084, 1999.
- Roberts, G. P., Variation in fault-slip directions along active and segmented normal fault systems, *J. Struct. Geol.*, **18**, 835–845, 1996.
- Rigo, A., Etude Simotectonique et Géodésique du Golfe de Corinthe (Grèce), Thèse d'Univ. è, Univ., Paris VII, France, 1994.
- Siddall, M., E. J. Rohling, A. Almogi-Labin, Ch. Hemleben, D. Meischner, I. Schmelzer, and D. A. Smeed, Sea-level fluctuations during the last glacial cycle, *Nature*, **423**, 853–858, 2003.
- Stein, M., G. J. Wasserburg, P. Aharon, J. H. Chen, Z. R. Zhu, A. Bloom, and J. Chappell, TIMS U-series dating and stable isotopes of the last interglacial event in Papua New Guinea, *Geochim. Cosmochim. Acta*, **57**, 2541–2554, 1993.
- Stirling, C., T. M. Esat, M. T. McCulloch, and K. Lambeck, High-precision U-series dating of corals from Western Australia and implications for the timing and duration of the Last Interglacial, *Earth Planet. Sci. Lett.*, **135**, 115–130, 1995.
- Stefatos, A., G. Papatheodorou, G. Ferentinos, M. Leeder, and R. Collier, Seismic reflection imaging of active offshore faults in the Gulf of Corinth: Their seismotectonic significance, *Basin Research*, **14**, 487–502, 2003.
- Valensise, G., and S. N. Ward, Long-term uplift of the Santa-Cruz coastline in response to repeated earthquakes along the San-Andreas fault, *Bull. Seismol. Soc. Am.*, **81**(5), 1694–1704, 1991.
- Van Calsteren, P., and J. B. Schwieters, Performance of a thermal ionization mass-spectrometer with a deceleration lens system and post-deceleration detector selection, *Int. J. Mass Spec. Ion Proc.*, **146**, 119–129, 1995.
- Vita-Finzi, C., Evaluation of late Quaternary uplift in Greece and Cyprus, in *Magmatic Processes and Plate Tectonics*, edited by H. M. Pritchard, T. Alabaster, N. B. W. Harris, and C. R. Neary, *Special Pub. Geol. Soc. London*, **76**, 417–424, 1993.

S. L. Houghton, G. P. Roberts, I. D. Papanikolaou, and J. M. McArthur, Research School of Earth Sciences at UCL-Birkbeck, Gower Street, WC1E 6BT, London, UK. (s.houghton@ucl.ac.uk)

M. A. Gilmour, Uranium Series Facility, Department of Earth Sciences, The Open University, Milton Keynes, MK7 6AA, UK.